

Manifolds with reduced holonomy in superstring theories

O.P.Santillan *

Contents

1. Introduction	2
2. Preliminary material	8
2.1 Quaternion Kahler and hyperkahler spaces	8
2.1.1 Quaternionic Kahler spaces in dimension higher than four	9
2.1.2 Quaternion Kahler manifolds in dimension four	12
2.2 Hypercomplex structures	14
2.2.1 Basic concepts	14
2.2.2 Examples of hypercomplex structures	16
2.3 Hyperkahler spaces in four dimensions	18
2.4 Quaternion Kahler and hyperkahler metrics in $d = 4$ with at least one isometry .	21
2.4.1 Hyperkahler metrics with Killing vectors that are not self-dual	21
2.4.2 Integrability of the axial continuum Toda equation	23
2.4.3 Einstein-Weyl structures and hyperkahler metrics	25
2.4.4 Self-dual structures with one Killing vector	27
2.4.5 The Joyce spaces	29
2.4.6 Identification of the quaternionic-Kahler metrics with at least one isometry	31
2.4.7 Examples of toric quaternion Kahler spaces	33
2.5 Higher dimensional hypergeometry	40
2.5.1 Construction of $4n$ hyperkahler manifolds with T^n tri-holomorphic isometry	40
2.5.2 Quaternion Kahler spaces in quaternion notation	42
2.5.3 The Swann extension	44
2.6 Spaces with G_2 holonomy	46
2.6.1 The group G_2 and the octonions	46
2.6.2 G_2 holonomy and self-duality	52
2.6.3 The Bryant-Salamon construction	54
2.6.4 Weak G_2 and Spin(7) holonomy spaces	56
2.7 G_2 holonomy spaces and M-theory compactifications	60

*Bogoliubov Laboratory of Theoretical Physics, JINR, Dubna, Russia, email: firenzecita@hotmail.com and osvaldo@theor.jinr.ru

3. Heterotic geometry without isometries	63
3.1 Introduction	63
3.2 Hyperkahler torsion manifolds	66
3.2.1 Main properties	66
3.2.2 Relation with the Plebanski-Finley conformal structures	68
3.3 The general solution	71
3.4 Discussion	76
4. D-instanton sums for matter hypermultiplets	78
4.1 Ooguri-Vafa solution	78
4.2 Pedersen-Poon Ansatz	80
4.3 D-instanton sums	81
4.4 Discussion	83
5. Hyperkahler spaces with local triholomorphic $U(1) \times U(1)$ isometry and supergravity solutions	84
5.1 Toric hyperkahler metrics of the Swann type	84
5.2 Supergravity solutions related to hyperkahler manifolds	87
6. G_2 toric metrics and supergravity backgrounds	90
7. Conclusion	94
A Supersymmetric sigma models	95

1. Introduction

The Weinberg-Salam theory is a quantum field theory that explain successfully two of the four known interactions of the nature, the electromagnetic and the weak interactions. The strong interaction is widely believed to be described by quantum chromodynamics, which is a theory that fits successfully with deep inelastic lepton nucleon scattering data. The remaining interaction, namely gravity, is believed to be described at classical level, by Einstein's General Theory of Relativity. The Weinberg-Salam model plus quantum chromodynamics and Einstein gravity are incompatible for small distances or, which is the same, for high energy scales of the order of the Planck mass $M_{pl} \approx 10^{19}$ GeV. The physics at this scale of energy is not known yet.

Superstring theory attempts to explain the physical phenomena at this scale. The present work deals with applications of manifolds with reduced holonomy to supersymmetric and superstring theories. The problem of classifying the possible holonomy groups for Riemannian and pseudo Riemannian spaces was posed by Cartan in [2] and partially solved by Berger in [1], who presented a list of all the possible holonomy groups. Two special cases of this list, relevant for our purposes, are the quaternion Kahler and hyperkahler spaces. By definition these spaces are $4n$ -dimensional with holonomies included in the Lie groups $Sp(n) \times Sp(1)$ and $Sp(n)$, respectively. Some of their properties has been investigated in [3], [4], and [5] but they are not completely classified at the present.

It is convenient to explain what we mean when we speak about an holonomy group. Consider a field ψ defined over a given point p on a n -dimensional manifold M , and let us parallel transport it around a given closed curve γ . In general the result of this transport will be a new

field ψ_γ at p that is different than the original one ψ . Let us define the transformation matrix H_γ by

$$\psi_\gamma = H_\gamma \psi. \quad (1.1)$$

If we take into account all the possible closed C^1 -piecewise curves passing through the point p it can be shown that the corresponding matrices $H(p)$ form a group. This is the full holonomy group of the manifold at the point p and is a subgroup of the orthogonal group $O(n)$. If instead we consider only curves γ that are contractible to a point, we will obtain the so called restricted holonomy group $H^0(p)$ which is a subgroup of $SO(n)$.

If we change from the point p to a point q and fix some C^1 -piecewise curve β connecting both point we will have the transformation

$$H(q) = A(\beta)H(p)A(\beta)^{-1}$$

and the same for $H^0(p)$. The matrix A belongs to the holonomy group $H(p)$ and it follows that $H(p)$ and $H(q)$ are isomorphic. Therefore we can talk about holonomy groups without reference to some specific point p . A space is orientable if and only if the holonomy group is in $SO(n)$. If the fundamental group $\pi_1(M) = 0$, then $H = H^0$. We recall that there exist non simply connected manifolds for which $\pi_1(M) = 0$, an example is the flat tori [6].

From the definition of the curvature tensor R_{abcd} [7] it is known the variation of ψ under an infinitesimal closed curve γ is given by

$$\delta\psi = G_{ab}\delta A^{ab}\psi. \quad (1.2)$$

Here A^{ab} is the area spanned by the curve γ and $G_{ab} = R_{abcd}\gamma^{cd}$ being γ^{cd} some generators of $SO(n)$ in the representation of the field (for applications to string theory we deal usually with spinor fields ψ , but this is not important in our discussion). Then it is clear that the "infinitesimal" holonomy group H^i is determined completely by the curvature tensor. Under infinitesimal parallel transport ψ satisfies

$$\partial_i\psi = -\omega_{iab}\gamma^{ab}\psi, \quad (1.3)$$

being ω_b^a the spin connection defined by

$$de^a + \omega_b^a \wedge e^b = 0.$$

From (1.3) we infer that the transformation around a (not necessarily infinitesimal) curve γ is given by the operator

$$U(\gamma) = P \exp\left(-\int_\gamma \omega_{iab}\gamma^{ab}dx^i\right) \quad (1.4)$$

where P denote path-ordering of the exponential [8]. By considering all this curves we obtain the full holonomy group, which could be larger than the infinitesimal one.

If the curve γ is homotopic to the identity then the matrices $U(\gamma)$ are generically in $SO(n)$ with the tangent indices of ω_b^a taking values in the Lie algebra $so(n)$. But there exists infinitely many examples in which the holonomy is reduced to a subgroup of $SO(n)$. A characteristic feature of this reduction is the presence of globally defined tensors α that are invariant under parallel transport. This property is equivalent to the annulation of their covariant derivative, i.e, $D\alpha = 0$ [6]. A globally parallel transported field is an analogous concept of an invariant subspace for the Lie group $SO(n)$. The presence of an invariant subspace implies the reduction

of $SO(n)$ to another Lie group with lower dimension. A common feature of reduced holonomy is in general the presence of covariantly constant p -forms that play an analog role than those invariant subspaces. For instance in $D = 4$ we have the isomorphism $SO(4) \simeq SU(2)_L \times SU(2)_R$, and $SU(2)$ holonomy spaces are characterized by a covariantly constant right-handed (or left-handed) spinor ϵ_R defined over the whole manifold. The presence of ϵ_R implies that the curvature is self-dual and that there exist a rotation of the frame taking the anti-self-dual part of ω_b^a to zero. Examples with self-dual curvature are the Eguchi-Hanson and Taub-Nut spaces. Also for G_2 holonomy manifolds one can choose an orthogonal frame e^i in which the octonionic 3-form

$$\begin{aligned} \Phi = & e^1 \wedge e^2 \wedge e^7 + e^1 \wedge e^3 \wedge e^6 + e^1 \wedge e^4 \wedge e^5 + e^2 \wedge e^3 \wedge e^5 + e^4 \wedge e^2 \wedge e^6 \\ & + e^3 \wedge e^4 \wedge e^7 + e^5 \wedge e^6 \wedge e^7 \end{aligned}$$

and its dual $*\Phi$ are closed [9]. This forms are indeed invariant under the action of G_2 and their closure is equivalent to the presence of a spinor η such that $D_i\eta = 0$. We also recall that G_2 is indeed a subgroup of $SO(7)$ with a one dimensional invariant subspace (as the space generated by η).

During the last twenty years it became clear that quaternion Kahler and hyperkahler geometry is an effective tool in field theory with and without the presence of supersymmetries. For example the moduli space of magnetic monopoles and the moduli space of Yang-Mills instantons in flat space are hyperkahler [10], [11]. Hyperkahler geometry also describe the couplings of supersymmetric sigma models with $N=4$ rigid supersymmetries [12],[13] and [44]. If a Wess-Zumino type term is included, then the resulting geometry will be hyperkahler torsion (HKT) [95]. When the supersymmetry is local the hypermultiplets appears coupled to gravity and the resulting target space is a quaternionic Kahler manifold [14]. Quaternionic manifolds also characterize the hypermultiplet geometry of classical and perturbative moduli spaces of type II strings compactified on a Calabi-Yau manifold [137], and even taking into account non perturbative corrections due to D-instantons [173]. Therefore there appears an interesting problem to construct quaternion Kahler spaces with singularities [132]. Other applications of this geometry can be found in [15]-[20] and references therein.

One of the latest achievements for constructing this type of manifolds is the hyperkahler quotient [12], [21], a simple example of this quotient is realized in the ADHM construction [11]. This is a construction of hyperkahler manifolds of a given dimension taking the quotient of a higher dimensional hyperkahler one by certain group generating tri-holomorphic isometries, and it have a physical origin related to dualities in supersymmetric sigma models [12]. A sort of inverse method is due to Swann [22] who shows how a quaternionic Kahler metric in $D=4n$ can be extended to a quaternionic Kahler and hyperkahler examples in $D=4(n+1)$. The Swann construction was applied recently to construct hyperkahler cones in [23], relevant in theories with $N=2$ rigid supersymmetries, and to construct certain scalar manifolds in M-theory on a Calabi-Yau threefold in the vicinity of a flop transition [24].

Hypergeometry (that is quaternion Kahler and hyperkahler geometry) is not the only case of the Berger list with applications in modern theoretical physics. Spaces of $SU(3)$, G_2 and $Spin(7)$ holonomy play a crucial role as internal spaces of heterotic string and M-theory preserving certain amount of supersymmetries [25], [26]. The main feature of such manifolds is the presence of parallel spinors fields η_i . The number of supersymmetries preserved by the usual Kaluza-Klein reduction is related to the number of such spinors over the internal space. In general, if a given Riemannian metric with dimension n admits at least one covariantly constant spinor η satisfying $D_i\eta = 0$ the holonomy group will be $SU(\frac{n}{2})$, $Sp(\frac{n}{4})$, G_2 or $Spin(7)$. The last two

cases corresponds to seven and eight dimensions. For G_2 (and $Spin(7)$) holonomy manifolds there is exactly one η .

In fact after the completion of the Berger work it remained as an open question the existence of metrics with exceptional holonomies G_2 and $Spin(7)$. This problem was solved by Bryant and Salamon thirty years after the appearance of the Berger list [27]. They showed that such spaces indeed exist by providing a family of examples [29]. The construction of this family was based on an interesting link between quaternion kahler and special holonomy spaces. This link allows to find a large class of spaces with special holonomy as extensions of four dimensional quaternion Kahler manifolds. This is called the Bryant-Salamon extension, and has certain analogies with the Swann one.

Spaces with special holonomy has vanishing Ricci tensor, i.e, they are Ricci-flat. Their curvature tensor R_{abcd} satisfies a generalization of self-duality condition to $D = 7$ namely

$$R_{ab} = \pm \frac{c_{abcd}}{2} R_{cd}.$$

The octonion constants c_{abcd} in $D = 7$ play an analogous role of the Kronecker symbols ε_{abcd} in $D = 4$. Self-duality implies Ricci-flatness and G_2 restricted holonomy [93] (analogous considerations hold in eight dimensions for $Spin(7)$). Although self-duality gives a non linear system of equations they have been solved in cases with suitable symmetries [9].

After the Bryant work explicit compact and non-compact G_2 holonomy metrics were constructed in [28] and [29], and complete ones in [30] and [31]. Recently, new examples have been found in [32], [33], [34], [36], [37], [38] and [35]. There are in the literature examples of "weak G_2 holonomy" [26], which are again backgrounds of the M-theory that give rise to N=1 supersymmetry in $D = 4$. In this case, there is a spinor field η which is not covariantly constant but satisfies $D_i \eta \sim \lambda \gamma_i \eta$. The Ricci flatness condition is replaced by $R_{ij} \sim \lambda g_{ij}$. In the limit $\lambda \rightarrow 0$ one obtain G_2 as restricted holonomy group. Hitchin has shown that under certain conditions is possible to construct this kind of manifolds starting with an $Spin(7)$ holonomy one [39]. Physically weak G_2 spaces are supersymmetric backgrounds in presence of fluxes.

Although there is a big progress in the study of special holonomy spaces, there are also some open problems, in particular related to Kaluza-Klein theories. A realistic compactification of M-theory should give rise to chiral matter and this can not be obtained by applying Kaluza-Klein procedure over G_2 smooth manifolds [40]. In the smooth case the harmonic Kaluza-Klein decomposition of the eleven-dimensional supergravity is the N=1 four dimensional supergravity coupled Abelian vector multiplets. But the chiral matter fields can emerge only if the manifold develops a singularity, as pointed out by Witten and Acharya in [41] and [42]. It turns out that to obtain a realistic model one should investigate the dynamics of the M-theory over orbifolds. Another request is that the internal space should be compact. The existence of compact spaces with special holonomy and orbifold singularities was proved rigorously by Joyce in [32]. But no explicit metric over such spaces is known yet.

Fortunately, some examples with weak G_2 holonomy that are compact and admitting certain kind of singularities are known [43]. Moreover Witten has shown that the physics near a singularity is local in essence and independent on the global properties of the internal space [95]. For this reason there is still much interest in construct special holonomy manifolds with conical singularities without the restriction to be compact.

It should be mentioned that the study of explicit metrics with exceptional holonomy has also importance in the context of dualities of string theory and M-theory [100]-[103]. Also generalizations of the work of Acharya and Witten on singular G_2 spaces were investigated recently

in [96], and over complete spaces with torus symmetry and admitting only orbifold singularities in [97] and [98]. Moreover compact G_2 -holonomy spaces with orbifold singularities are believed to arise as quotients of a certain conical hyperkahler manifold in $D=8$ by one of its isometries [42], thus providing a new link between special holonomy manifolds and hypergeometry. The range of applications of this topic is very wide; some of them can be found in [106]-[124].

The purpose of this dissertation

This dissertation is based on the works of the author during his studies in order to get the degree of Candidate in Sciences [189],[190], [191] and [192] and is related to applications of reduced holonomy manifolds and hyperkahler torsion geometry to string and M-theory. Our aim is the following:

a) To show that in four dimensions weak hyperkahler torsion structures are the same than hypercomplex structures, and also equivalent to certain self-dual structures considered by Plebanski and Finley [192].

b) To find the most general form for a weak hyperkahler torsion metric in four dimensions and discuss some particular examples [192].

c) To construct the hyperkahler target space metric corresponding to several matter hypermultiplets of type IIA superstring compactified on a Calabi-Yau threefold, even taking into account the D-instanton contributions, by imposing the physical requirements due to Vafa and Ooguri to a generic $4n$ dimensional hyperkahler metric [191].

d) To construct a family of toric hyperkahler spaces with tri-holomorphic killing vectors in eight dimensions. To lift the result to new eleven dimensional backgrounds not related to D-brane solutions. To find new IIA backgrounds by reduction along one of the isometries and new IIB backgrounds by use of dualities [190].

e) To construct a family of toric G_2 holonomy manifold and $SU(3)$ torsion structures with two commuting isometries [189]. To lift the result to new eleven dimensional backgrounds. To find IIA backgrounds by reduction along one of the isometries and IIB backgrounds by use of dualities [190].

The structure of this dissertation

In the preliminary material we explain in detail the elementary features about hyperkahler geometry, hypercomplex structures, quaternion Kahler spaces, G_2 and $Spin(7)$ holonomy manifolds. The results of this section are rather mathematical and not original at all, but the presentation is new. We consider of great importance to make clear certain mathematical features in simple language. In section 2.1.1 we present the most important features about quaternion Kahler and hyperkahler geometry in dimension higher than four. In section 2.1.2 we show that in four dimensions any quaternion Kahler space is self-dual Einstein. In section 2.2 we give definition of an hypercomplex structures. In this context we review in section 2.3 four dimensional hyperkahler geometry, in particular the Ashtekar-Jacobson-Smolín and Plebanski formulations. This part is importance for the points a) and b) mentioned above. Hyperkahler

spaces with isometries are presented in section 2.4, in particular it is reviewed the Boyer-Finley classification of the possible Killing vectors for this geometry. It is also remarked a geometrical interpretation for the integrability of the axial $SU(\infty)$ Toda equation that naturally appears in this context. In section 2.4.3 it is presented a one to one link between self-dual conformal structures and Einstein-Weyl structures, which is known as the Jones-Tod correspondence. We also present in section 2.4.4 the most general self-dual structures with two commuting isometries that are non conformal to an hyperkahler space namely, the Joyce spaces. The other part of section 2.4 present the subcases of the Joyce spaces that are Einstein, and therefore quaternion Kahler. This are the Calderbank-Pedersen spaces, which play an important role in the present dissertation, in particular for the points d) and e).

Section 2.5 concerns with certain aspects of higher dimensional quaternion Kahler and hyperkahler geometry. In section 2.5.1 are presented the most general $4n$ dimensional hyperkahler spaces with n commuting tri-holomorphic isometries, which is a higher dimensional generalization of the Gibbons-Hawking metrics. The metrics are written in the momentum map system, which is the most suitable form for the purposes of this dissertation. This result will be used in the point c) and d). We also present in section 2.5.3 the Swann extension, which is a construction of a higher dimensional quaternion kahler space starting with other with less dimensions. This part concerns mainly with the point d).

Section 2.6 presents some features about the groups G_2 and $Spin(7)$ and the relation with the octonion algebra. We show the link between G_2 and $Spin(7)$ holonomy manifolds, the presence of globally defined covariantly constant spinor fields and octonionic self-duality. We present the Bryant-Salamon construction, which extend any quaternion kahler manifold in four dimensional to a seven dimensional one with holonomy included in G_2 and to an eight dimensional one with holonomy in $Spin(7)$. We also present spaces with weak G_2 holonomy as the cones of certain $Spin(7)$ spaces, and half-flat six dimensional manifolds as the cones of certain G_2 spaces. We emphasize the relation between G_2 holonomy manifolds and compactifications of M theory preserving certain amount of supersymmetries. This part is related to the point e).

The remaining sections are devoted to solve the tasks a)-e) enumerated above. The preliminary material plays an important role in order to understand this section. In the conclusions we mention the main results of the present work. We have included an appendix that explain the relations between hyperkahler torsion geometry and supersymmetric sigma models. At the end it is included the bibliography and the publications of the author.

2. Preliminary material

2.1 Quaternion Kahler and hyperkahler spaces

By definition, a quaternion Kahler manifold is an euclidean $4n$ dimensional manifold with holonomy group Γ included into the Lie group $Sp(n) \times Sp(1) \subset SO(4n)$ [1]. This statement is non trivial if $D > 4$, but in $D = 4$ we have the isomorphism $SO(4) \simeq SU(2)_L \times SU(2)_R \simeq Sp(1) \times Sp(1)$ related to the orthogonal decomposition of spinors in left and right states of chirality, and so the statement $\Gamma \subseteq Sp(1) \times Sp(1)$ is trivial and should be modified. The first subsection of this section concerns with spaces with dimension at least 8, the case $D = 4$ will be treated separately. We will show that a quaternionic Kahler metric has the following properties [4], [3].

1) There exists three automorphism J^i ($i = 1, 2, 3$) of the tangent space TM_x at a given point x such that $J^i \cdot J^j = -\delta_{ij} + \epsilon_{ijk} J^k$, and for which the metric g is quaternion hermitian, that is

$$g(X, Y) = g(J^i X, J^i Y), \quad (2.5)$$

being X and Y arbitrary vector fields.

2) The structures J^i satisfy the fundamental relation

$$\nabla_X J^i = \epsilon_{ijk} J^j \omega_-^k, \quad (2.6)$$

with ∇_X the Levi-Civita connection of the manifold and ω_-^i its $Sp(1)$ part. As a consequence of hermiticity of g stated above the tensor $\bar{J}_{ab}^i = (J^i)_a^c g_{cb}$ is antisymmetric and therefore there exists an associated 2-form

$$\bar{J}^i = \bar{J}_{ab}^i e^a \wedge e^b$$

for which

$$d\bar{J}^i = \epsilon_{ijk} \bar{J}^j \wedge \omega_-^k, \quad (2.7)$$

being d the usual exterior derivative.

3) Corresponding to the $Sp(1)$ connection we can define the 2-form

$$F^i = d\omega_-^i + \epsilon_{ijk} \omega_-^j \wedge \omega_-^k.$$

Then for a quaternion Kahler manifold

$$R_-^i = \Lambda \bar{J}^i, \quad (2.8)$$

$$F^i = \Lambda' \bar{J}^i. \quad (2.9)$$

being Λ and Λ' certain constants. The tensor R_-^a is the $Sp(1)$ part of the curvature. The last two conditions implies that g is Einstein with non zero cosmological constant $R_{ij} \sim g_{ij}$.

4) In a quaternion Kahler space the globally defined $(0, 4)$ and $(2, 2)$ tensors

$$\Theta = \bar{J}^1 \wedge \bar{J}^1 + \bar{J}^2 \wedge \bar{J}^2 + \bar{J}^3 \wedge \bar{J}^3,$$

$$\Xi = J^1 \otimes J^1 + J^2 \otimes J^2 + J^3 \otimes J^3$$

are covariantly constant with respect to the usual Levi Civita connection.

5) Any quaternion Kahler space is orientable.

6) If there exists a local frame in which ω_-^i is equal to zero, then the metric is called hyperkahler and is Kahler with respect to any of the complex structures. The metric g is in this case Ricci-flat and the holonomy is reduced to a subgroup of $Sp(n)$.

7) In four dimensions quaternion Kahler spaces are the same than Einstein spaces and with self-dual Weyl tensor.

This section is devoted to explain this concepts in more detail. By taken granted properties 1-7 the reader can go directly to the next section.

2.1.1 Quaternionic Kahler spaces in dimension higher than four

It is well known that the generators J^i of the Lie algebra $sp(1)$ of $Sp(1) \simeq SU(2)$ have the multiplication rule

$$J^i \cdot J^j = -\delta^{ij} I + \epsilon_{ijk} J^k, \quad (2.10)$$

which implies the $so(3) \simeq su(2)$ commutation rule

$$[J^i, J^j] = \epsilon_{ijk} J^k. \quad (2.11)$$

We see that $J^i J^i = -I$ and therefore J^i will be called almost complex structures. An useful $4n \times 4n$ representation is

$$J^1 = \begin{pmatrix} 0 & -I_{n \times n} & 0 & 0 \\ I_{n \times n} & 0 & 0 & 0 \\ 0 & 0 & 0 & -I_{n \times n} \\ 0 & 0 & I_{n \times n} & 0 \end{pmatrix}, \quad J^2 = \begin{pmatrix} 0 & 0 & -I_{n \times n} & 0 \\ 0 & 0 & 0 & I_{n \times n} \\ I_{n \times n} & 0 & 0 & 0 \\ 0 & -I_{n \times n} & 0 & 0 \end{pmatrix}$$

$$J^3 = J^1 J^2 = \begin{pmatrix} 0 & 0 & 0 & -I_{n \times n} \\ 0 & 0 & -I_{n \times n} & 0 \\ 0 & I_{n \times n} & 0 & 0 \\ I_{n \times n} & 0 & 0 & 0 \end{pmatrix}. \quad (2.12)$$

The group $SO(4n)$ is a Lie group and this means in particular that for any $SO(4n)$ tensor A_b^a the commutator $[A, J^i]$ will take also values in $SO(4n)$. We will say that A belong to the subgroup $Sp(n)$ of $SO(4n)$ if and only if

$$[A, J^i] = 0. \quad (2.13)$$

Condition (2.13) together with (2.11) implies that a tensor B_b^a belongs to the subgroup $Sp(n) \times Sp(1)$ if and only if

$$[B, J^i] = \epsilon_{ijk} J^j B_-^k,$$

being B_-^k the components of B in the basis J^k . Both conditions are independent of the representation.

We will write a metric over a $4n$ dimensional manifold M as $g = \delta_{ab}e^a \otimes e^b$, being e^a the $4n$ -bein basis for which g is diagonal. Let us define the triplet of $(1, 1)$ tensors

$$J^i = (J^i)_b^a e_a \otimes e^b, \quad (2.14)$$

defined by the matrices (3.300). If the holonomy is in $Sp(n) \times Sp(1)$, then from the beginning ω_b^a will take values on its lie algebra $sp(n) \oplus sp(1)$. As we saw above, this implies that

$$[\omega, J^i] = \epsilon_{ijk} J^j \omega_-^k. \quad (2.15)$$

As usual, the connection ω_b^a is defined through

$$\nabla_X e^a = -\omega_b^a(X) e^b,$$

together with the Levi-Civita conditions $\nabla g = 0$ and $T(X, Y) = 0$. Using the chain rule $\nabla(A \otimes B) = (\nabla A) \otimes B + A \otimes (\nabla B)$ for tensorial products show us that in the einbein basis

$$[\omega, J^i] = \nabla_X J^i. \quad (2.16)$$

Comparing (2.15) and (2.16) we see that quaternionic Kahler manifold are defined by the relation

$$\nabla_X J^i = \epsilon_{ijk} J^j \omega_-^k,$$

which is independent on the election of the frame e^a . This proves that (2.6) describe quaternion Kahler metrics [4] in dimension higher than four.

The basis e^a for a metric g is defined up to an $SO(4n)$ rotation. Under this $SO(4n)$ transformation the tensors (2.14) are also transformed, but it can be shown that the multiplication (2.10) is unaffected. In other words, given the tensors J^i one can construct a new set of complex structures

$$J'^i = C_j^i J^j, \quad J'^i \cdot J'^j = -\delta^{ij} I + \epsilon_{ijk} J'^k \quad \Longleftrightarrow \quad C_k^i C_j^k = \delta_j^i \quad (2.17)$$

This can be paraphrased by saying that a quaternionic Kahler manifold has a bundle V of complex structures parameterized by the sphere S^2 . Using the textbook properties of ∇ it can be seen that (2.6) is unaltered under such rotations.

Let us define three new tensors $(\bar{J}^i)_{ab}$ by $(\bar{J}^i)_{ab} = (J^i)_a^c \delta_{cb}$. From (3.300) it follows that

$$(J^i)_b^a = -(J^i)_a^b \quad \Longleftrightarrow \quad (\bar{J}^i)_{ab} = -(\bar{J}^i)_{ba}$$

This show that $(\bar{J}^i)_{ab}$ are the components of the two-forms $\bar{J}^i$ defined by

$$\bar{J}^i = (\bar{J}^i)_{ab} e^a \wedge e^b. \quad (2.18)$$

The forms (3.330) are known as the hyperkahler forms. From (2.6) it is obtained that

$$\nabla_X J^i = \epsilon_{ijk} J^j \omega_-^k \quad \Longrightarrow \quad d\bar{J}^i = \epsilon_{ijk} \omega_-^j \wedge \bar{J}^k,$$

being d the usual exterior derivative. The last implication proves relation (2.7).

If we change the frame e^a to a new one x_μ then the definition $(\bar{J}^i)_{ab} = (J^i)_a^c \delta_{cb}$ should be modified by the covariant one $(\bar{J}^i)_{\alpha\beta} = (J^i)_\alpha^\gamma g_{\gamma\beta}$. Here the greek index indicates the components in the new basis and $g_{\gamma\beta}$ are the corresponding components of the metric. Therefore

$$(\bar{J}^i)_{ab} = -(\bar{J}^i)_{ba} \quad \Longleftrightarrow \quad (J^i)_\alpha^\gamma g_{\gamma\beta} = (J^i)_\beta^\gamma g_{\gamma\alpha}$$

The last relation is equivalent to

$$g(J^i X, Y) = g(X, J^i Y) \iff g(X, Y) = g(J^i X, J^i Y)$$

for arbitrary vector fields X and Y in TM . Then the metric g will be always quaternion hermitian with respect to the complex structures. Relation (2.5) is also invariant under the automorphism of the complex structures.

In general, if in a given manifold there exist three complex structures satisfying (2.10), and we take intersecting coordinate neighborhoods U and U' , then we have two associated basis J^i and J'^i . Both basis should be related by an $SO(3)$ transformation in order to satisfy (2.10). This means that any quaternion Kahler space is orientable [4]. Consider now the fundamental 4-form

$$\Theta = \overline{J}^1 \wedge \overline{J}^1 + \overline{J}^2 \wedge \overline{J}^2 + \overline{J}^3 \wedge \overline{J}^3, \quad (2.19)$$

and the globally defined $(2, 2)$ tensor

$$\Xi = J^1 \otimes J^1 + J^2 \otimes J^2 + J^3 \otimes J^3. \quad (2.20)$$

By means of the formula (2.17) it follows that both tensors (2.19) and (2.20) are globally defined on the manifold M . For a quaternionic Kahler manifold it is obtained directly from (2.6) and (2.7) that [4]

$$\nabla \Theta = 0, \quad \nabla \Xi = 0.$$

In $D = 8$ for a quaternion Kahler manifold $d\Theta = 0$ and if the manifold is of dimension at least 12 then $d\Theta$ determines completely $\nabla \Theta$. In particular $d\Theta = 0$ implies $\nabla \Theta = 0$ [22].

One of the most important consequences of (2.6) is that quaternionic Kahler spaces are always Einstein with cosmological constant [3]. The proof is briefly as follows. From the definition of the curvature tensor $R(X, Y) = [\nabla_X, \nabla_Y] - \nabla_{[X, Y]}$ together with (2.6) it follows in the einbein basis that

$$R_{ijm}^l (J^a)_k^m - R_{ijk}^m (J^a)_m^l = \epsilon_{abc} (F^b)_{ij} (J^c)_k^l. \quad (2.21)$$

where R_{ijm}^l are the components of the curvature tensor and the two form F^a was defined as ¹

$$F^a = d\omega_-^a + \epsilon_{abc} \omega_-^b \wedge \omega_-^c.$$

We can rewrite (2.21) as a commutator

$$[R(X, Y), J^a] = \epsilon_{abc} F^b J^c,$$

being X and Y arbitrary vector fields. Multiplying (2.21) by $(J^a)_l^s$ and contracting indices, and then multiplying by $(J^b)_l^k$ and using the identity ²

$$(J^a)_l^s (J^b)_s^l = 4n \delta^{ab},$$

gives the formula

$$F_{ij}^a = \frac{1}{2n} R_{ijk}^l (J^a)_l^k. \quad (2.22)$$

¹In the physical literature sometimes the three components ω_- are referred as an $SU(2)$ vector potential and F^a as the corresponding strength tensor

²Which is clearly true in the representation (3.300)

Inserting (2.22) into (2.21) yields

$$R_{ijk}^l (J^a)_l^k = \frac{2n}{2+n} R_{im} (J^a)_j^m,$$

which can also be expressed as

$$R_-^i = \frac{2n}{2+n} R \bar{J}^i, \quad (2.23)$$

being R is the scalar curvature and R_-^i are the $Sp(1)$ components of the curvature tensor. The second Bianchi identities together with (2.23) shows that R is constant and thus $R_{ij} \sim g_{ij}$ [3]. Thus, in any dimension, quaternionic Kahler spaces are always Einstein with non zero cosmological constant λ .

Because R is a constant we see from (2.23) that

$$R_-^i = \Lambda \bar{J}^i,$$

being Λ certain constant. We also have from (2.22) that

$$F^i = \Lambda' \bar{J}^i,$$

being Λ' another constant. This condition are the same as (2.8) and (2.9) given above. In the limit $\lambda \rightarrow 0$ the constants Λ and Λ' goes simultaneously to zero.

If there exists a rotation of the local frame for which $\omega_- = 0$ then the complex structures are locally covariantly constant, that is

$$\nabla_X J^i = 0. \quad (2.24)$$

In this case $R_-^i = F^i = 0$ thus the space has self-dual curvature, which implies Ricci flatness. This spaces are called hyperkahler, and (2.24) shows that they are Kahler with respect of any of the complex structures. Condition (2.24) implies that the holonomy is in $Sp(n)$ and that

$$d\bar{J}^a = 0 \quad (2.25)$$

together with the annulation of the Niejenhuis tensor given by

$$N(X, Y) = [X, Y] + J[X, JY] + J[JX, Y] - [JX, JY], \quad (2.26)$$

[60]. A complex structure for which $N(X, Y) = 0$ is called integrable. We will explain this relations in more detail in section 4.

2.1.2 Quaternion Kahler manifolds in dimension four

As we saw starting the previous section, in four dimensions the statement that the holonomy is $\Gamma \subseteq Sp(n) \times Sp(1)$ is trivial due to the isomorphism $SO(4) \simeq SU(2)_L \times SU(2)_R \simeq Sp(1) \times Sp(1)$. We will modify this definition and we will say that a four dimensional manifold M is quaternionic Kahler if (2.8) holds. This condition is not trivial, we will show below that quaternion Kahler spaces in $d = 4$ are Einstein (as in the higher dimensional case) and with self-dual Weyl tensor.

Let us consider a four dimensional metric $g = \delta_{ab} e^a \otimes e^b$ and the connection ω_b^a given by the first Cartan equation

$$de^a + \omega_b^a \wedge e^b = 0, \quad \omega_{ib}^a = -\omega_{bi}^a.$$

The notation $SU(2)_\pm$ denote the $SU(2)_L$ and $SU(2)_R$ groups respectively. The $SU(2)_\pm$ components of the spin connection are explicitly

$$\omega_\pm^a = \omega_0^a \pm \epsilon_{abc}\omega_c^b. \quad (2.27)$$

The curvature tensor is given by the second Cartan equation

$$R_b^a = d\omega_b^a + \omega_s^a \wedge \omega_b^s = R_{b,st}^a e^s \wedge e^t$$

and the $SU(2)$ parts are

$$R_\pm^a = R_0^a \pm \epsilon_{abc}R_c^b. \quad (2.28)$$

The Ricci tensor is defined in the diagonal basis by $R_{ij} = R_{i,aj}^a$ and the scalar curvature is $R_{ii} = R$.

Instead of use the basis $e^a \wedge e^b$ we can use the basis $\bar{J}_\pm^a = e^0 \wedge e^a \pm \epsilon_{abc}e^b \wedge e^c$. Then it follows that $\bar{J}_\pm^a$ are separately complex structure with definite self-duality properties, that is

$$*\bar{J}_\pm^a = \pm \bar{J}_\pm^a.$$

In this basis

$$R_+^a = A_{ab}\bar{J}_+^b + B_{ab}\bar{J}_-^b, \quad R_-^a = B_{ab}^t\bar{J}_+^b + C_{ab}\bar{J}_-^b \quad (2.29)$$

where the matrices A and C are symmetric. The components of the Ricci tensor are

$$R_{00} = Tr(A + B), \quad R_{0a} = \frac{\epsilon_{abc}}{2}(B_{bc}^t - B_{bc}), \quad R_{ab} = Tr(A - B)\delta_{ab} + B_{ab} + B_{ab}^t, \quad (2.30)$$

and the scalar curvature is

$$R = 4Tr(A) = 4Tr(C). \quad (2.31)$$

It is clearly seen from (2.30) that the Einstein condition $R_{ij} = \Lambda\delta_{ij}$ is equivalent to $B = 0$ and $Tr(A) = Tr(C) = \Lambda$.

The components of the Weyl tensor in the diagonal basis are given by

$$W_{bcd}^a = R_{bcd}^a - \frac{1}{2}(\delta_{ac}R_{bd} - \delta_{ad}R_{bc} + \delta_{bd}R_{ac} - \delta_{bc}R_{ad}) + \frac{R}{6}(\delta_{ac}\delta_{bd} - \delta_{ad}\delta_{bc}). \quad (2.32)$$

The tensor W is invariant under a conformal transformation $g \rightarrow \Omega^2 g$ and the associated two form is

$$W_b^a = W_{bcd}^a e^c \wedge e^d.$$

An explicit calculation shows that the $SU(2)_\pm$ of W are

$$W_+^a = W_0^a + \epsilon_{abc}W_c^b = (A_{ab} - \frac{1}{3}Tr(A)\delta_{ab})\bar{J}_+^b,$$

$$W_-^a = W_0^a - \epsilon_{abc}W_c^b = (C_{ab} - \frac{1}{3}Tr(C)\delta_{ab})\bar{J}_-^b.$$

From this expressions we see that to say that an Einstein space is self dual (i.e, $W_-^a = 0$) is equivalent to

$$C_{ab} = \frac{\Lambda}{3}\delta_{ab} \iff R_-^a = \frac{\Lambda}{3}\bar{J}_-^a. \quad (2.33)$$

The second (2.33) is the same as (2.8) in four dimensions. Thus we conclude then that in $D = 4$ quaternionic Kahler is the same as self-dual Einstein.

The self-duality condition $W_- = 0$ is conformal invariant and then if a metric g is self-dual, then the family $[g]$ constructed by g through conformal transformations $g \rightarrow \Omega^2 g$ is also self-dual. But Einstein condition is not conformal invariant in general.

2.2 Hypercomplex structures

As we have seen, the problem of classifying the possible quaternion Kahler spaces appearing in $d = 4$ is equivalent to find all the Einstein spaces with self-dual Weyl tensor. In this section we will focus our attention in four dimensional self-dual spaces. Let us consider a metric g and denote with $[g]$ the family of metrics obtained from g by arbitrary conformal transformation $g \rightarrow \Omega^2 g$. Because the Weyl tensor is conformally invariant it follows that the self-duality of g implies the self-duality of $[g]$. The converse of this statement is also true.

An important example of self-dual families are the hypercomplex structures [60], [61], that we will introduce in this section. We will show in the following sections that hypercomplex structures and weak torsion hyperkahler structures are the same concept in four dimensions.

2.2.1 Basic concepts

As it follows from equation (2.7), for hyperkahler spaces there exist a frame for which

$$d\overline{\mathcal{J}}^i = 0, \quad (2.34)$$

that is, for which the Kahler triplet is *closed*. Conversely, it was shown in [76] that any space in $d = 4$ satisfying (2.34) is automatically hyperkahler. The curvature tensor R_{bcd}^a of an hyperkahler space is automatically self-dual, and the almost complex structures are integrable, thus complex structure.

For a given hyperkahler structure $(g, \overline{\mathcal{J}}^i)$ one can define an new structure $(\tilde{g}, \tilde{\mathcal{J}}^i)$ being $\tilde{g} = \Omega^2 g$ and $\tilde{\mathcal{J}}^i = \Omega^2 \overline{\mathcal{J}}^i$. This new metric is quaternion hermitian with respect of the $\tilde{\mathcal{J}}^i$. But from (2.34) one obtains that

$$d\tilde{\mathcal{J}}^i + 2d \log(\Omega) \wedge \tilde{\mathcal{J}}^i = 0. \quad (2.35)$$

This means that $(\tilde{g}, \tilde{\mathcal{J}}^i)$ is not hyperkahler. A conformally invariant generalization of the hyperkahler condition (2.34) is

$$d\tilde{\mathcal{J}}^i + \alpha \wedge \tilde{\mathcal{J}}^i = 0, \quad (2.36)$$

being α an arbitrary 1-form. Under conformal transformation we have $\alpha + 2d \log(\Omega)$ and (2.36) is unaltered [61]. This property define a conformal family $[g]$ with all the element sharing the property (2.36).

If α is a gradient ($\alpha = \nabla \phi$, being ϕ certain function depending on all the coordinates), then it will exist a representative g of $[g]$ with self-dual curvature and therefore hyperkahler. From the expression (2.32) of the Weyl tensor it is possible to check that if R is self-dual, then W is also self-dual. Thus the family $[g]$ corresponding to an hyperkahler metric will have self-dual Weyl tensor (2.32). But the explicit calculation of the W_{abcd} for a metric satisfying (2.36) shows that α disappear in its calculation. Therefore the condition to be gradient is not necessary to ensure $W_- = 0$ and condition (2.36) defines a self-dual structure in general [61]. We will denote it as $([g], [\overline{\mathcal{J}}^i])$.

A family of structures $([g], [\overline{\mathcal{J}}^i])$ satisfying (2.36) are the hypercomplex structures in four dimensions [60]. For a given representative $(g, \overline{\mathcal{J}}^i)$ of an hypercomplex structure, the metric g is quaternion hermitian with respect to the $\overline{\mathcal{J}}^i$ and the Niejenhuis tensor

$$N(X, Y) = [X, Y] + J[X, JY] + J[JX, Y] - [JX, JY] \quad (2.37)$$

corresponding to every $\overline{J}^i$ vanish for every pair of vector fields X, Y . Such an almost complex structures are called integrable, and they turn into complex structures. The metric g will be complex in this case. The definition of hypercomplex structures is consistent due to the fact that the integrability condition is conformally invariant. In four dimensions we can select the self-dual complex structures

$$J^i = e_4 \otimes e^i - e_i \otimes e^4 + \epsilon_{ijk} e_j \otimes e^k \quad (2.38)$$

up to an $SU(2)$ transformation. The action of (2.38) over the tangent space TM_x is defined by

$$\begin{aligned} J^1(e_1) &= e_2, & J^1(e_2) &= -e_1, & J^1(e_3) &= e_4, & J^1(e_4) &= -e_3, \\ J^2(e_1) &= e_3, & J^2(e_2) &= -e_4, & J^2(e_3) &= -e_1, & J^2(e_4) &= e_2, \\ J^3(e_1) &= e_4, & J^3(e_2) &= e_3, & J^3(e_3) &= -e_2, & J^3(e_4) &= -e_1. \end{aligned} \quad (2.39)$$

Then the annulation of the tensor $N^i(X, Y)$ will be equivalent to the conditions

$$\begin{aligned} [e_1, e_2] + [e_3, e_4] &= -A_2 e_1 + A_1 e_2 + A_3 e_4 - A_4 e_3 \\ [e_1, e_3] + [e_4, e_2] &= -A_3 e_1 + A_4 e_2 + A_1 e_3 - A_2 e_4 \\ [e_1, e_4] + [e_2, e_3] &= -A_4 e_1 - A_3 e_2 + A_2 e_3 + A_1 e_4 \end{aligned} \quad (2.40)$$

for some set of functions A_1, A_2, A_3, A_4 [59]. This is a direct consequence of (2.39) and (2.37). We immediately recognize that in the case $A_a = 0$ we recover the Ashtekar-Jacobson-Smolín quadratic equations for self-dual spaces [153]. If we consider the conformal transformation $g \rightarrow \Omega^2 g$, then it is seen that (2.40) is also satisfied with $A_a \rightarrow A_a + 2e_a \log(\Omega)$. This means that condition (2.40) define a conformal family of metrics $[g]$ and therefore the definition of hypercomplex structures is consistent.

Hypercomplex condition implies, but is not implied by, that the family $[g]$ is self-dual [59]-[60]. This follows from the fact that, as we will show now, condition (2.40) implies (2.36). Let us consider the connection ω given by

$$de^a + \omega_b^a \wedge e^b = 0.$$

The antisymmetric part $\omega_{[bc]}^a$ is related to the structure functions defined by the Lie bracket $[e_a, e_b] = c_{ab}^c e_c$ by

$$\omega_{[bc]}^a = \frac{1}{2} c_{ab}^c.$$

It is convenient to define the one-form

$$\alpha = A - \chi$$

where

$$A = A_a e^a, \quad \chi = c_{ab}^b e^a, \quad (2.41)$$

and the self-dual two-form $\overline{J}^1 = e^1 \wedge e^2 + e^3 \wedge e^4$. Then

$$\begin{aligned} d\overline{J}^1 &= d(e^1 \wedge e^2 + e^3 \wedge e^4) \\ &= -\omega_{[ab]}^1 e^a \wedge e^b \wedge e^2 + \omega_{[ab]}^2 e^a \wedge e^b \wedge e^1 - \omega_{[ab]}^3 e^a \wedge e^b \wedge e^4 + \omega_{[ab]}^4 e^a \wedge e^b \wedge e^3 \end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{2}c_{ab}^1 e^a \wedge e^b \wedge e^2 + \frac{1}{2}c_{ab}^2 e^a \wedge e^b \wedge e^1 - \frac{1}{2}c_{ab}^3 e^a \wedge e^b \wedge e^4 + \frac{1}{2}c_{ab}^4 e^a \wedge e^b \wedge e^3 \\
&= e^1 \wedge e^2 \wedge (c_{ab}^a e^b + A_3 e^3 + A_4 e^4) + e^3 \wedge e^4 \wedge (c_{ab}^a e^b + A_1 e^1 + A_2 e^2) \\
&= (e^1 \wedge e^2 + e^3 \wedge e^4) \wedge (A - \chi)
\end{aligned}$$

and therefore

$$d\bar{J}^1 = \bar{J}^1 \wedge (A - \chi) = \bar{J}^1 \wedge \alpha \quad (2.42)$$

This is equivalent to (2.36). The same formula holds for $\bar{J}^2$ and $\bar{J}^3$. Therefore we reach to the conclusion that hypercomplex structures satisfies (2.36) and therefore are self-dual [59]. It is straightforward to prove that under $g \rightarrow \Omega^2 g$ the forms α , A and χ transform as

$$\begin{aligned}
\alpha_a &\rightarrow \alpha_a + 2e_a \log(\Omega) \\
A_a &\rightarrow A_a - 2e_a \log(\Omega) \\
\chi_a &\rightarrow \chi_a - 3e_a \log(\Omega).
\end{aligned} \quad (2.43)$$

Hypercomplex structures are not the only self-dual structures in four dimensions. Counterexamples are for instance the Joyce spaces that will be introduced in another section.

2.2.2 Examples of hypercomplex structures

Let us look now for hypercomplex structures $[g]$ over a manifold M with two commuting $U(1)$ Killing vectors. The representatives $[g]$ of such structures will be of Gowdy form

$$g = g_{ab} dx^a dx^b + g_{\alpha\beta} dx^\alpha dx^\beta. \quad (2.44)$$

The latin and greek indices takes values 1 and 2. Both g_{ab} and $g_{\alpha\beta}$ are supposed to be independent of the coordinates $x^\alpha = (\theta, \varphi)$. Then the Killing vectors are $\partial/\partial\theta$ and $\partial/\partial\varphi$ and are commuting, so there is a $U(1) \times U(1)$ action on the manifold.

By Gauss theorem there exists a local scale transformation $g \rightarrow \Omega^2 g$ which reduce (2.44) to

$$\tilde{g} = \Omega^2 (d\rho^2 + d\eta^2) + \tilde{g}_{\alpha\beta} dx^\alpha dx^\beta, \quad (2.45)$$

being $\tilde{g}_{\alpha\beta}$ function of the new coordinates (ρ, η) . We can express this metric as

$$\tilde{g} = \Omega^2 (d\rho^2 + d\eta^2) + (\bar{A}_0 d\theta + \bar{A}_1 d\varphi)^2 + (\bar{B}_0 d\theta + \bar{B}_1 d\varphi)^2, \quad (2.46)$$

where $\bar{A}_i$ and $\bar{B}_i$ are unknown functions of (ρ, η) and the factor ρ was introduced by convenience. But instead to work with the local form (2.46) we will work with the following equivalent expression

$$\tilde{g} = \Omega^2 (d\rho^2 + d\eta^2) + \frac{(A_0 d\theta - B_0 d\varphi)^2 + (A_1 d\theta - B_1 d\varphi)^2}{(A_0 B_1 - A_1 B_0)^2}. \quad (2.47)$$

Ansatz (2.47) looks more complicated than (2.46), but has the advantage that the dual basis e_i takes the simple form

$$e_1 = \Omega \partial_\rho, \quad e_2 = \Omega \partial_\eta \quad (2.48)$$

$$e_3 = B_1 \partial_\theta + A_1 \partial_\varphi, \quad e_4 = B_0 \partial_\theta + A_0 \partial_\varphi. \quad (2.49)$$

Inserting (2.48) and (2.49) into the quadratic system (2.40) gives $\Omega = 1$ and the Cauchy Riemman equations

$$(A_0)_\rho = (A_1)_\eta, \quad (A_0)_\eta = -(A_1)_\rho \quad (2.50)$$

and the same equations for B_0 and B_1 . This means that the metric (2.47) is described in terms of two holomorphic functions

$$F(z) = A_0 + iA_1, \quad G(z) = B_0 + iB_1, \quad (2.51)$$

being $z = \rho + i\eta$.

We can find the same result in another way. Let us consider four functions $f_1, \dots, f_4$ and $g_1, \dots, g_4$ depending on the coordinates $x^1 = \rho$ and $x^2 = \eta$ and let us define the vector fields

$$e_a = f_a \partial_\theta + g_a \partial_\varphi + \partial_{x^a} \quad (a = 1, 2),$$

$$e_a = f_a \partial_\theta + g_a \partial_\varphi, \quad (a = 3, 4).$$

Clearly e_a are the most general vector fields for a metric with two commuting isometries up to a conformal scaling. Introducing this expressions into (2.55) gives the system of equations

$$\begin{aligned} (f_3)_\eta - (f_4)_\rho &= 0, & (g_3)_\eta - (g_4)_\rho &= 0, \\ (f_2)_\eta + (f_1)_\rho &= 0, & (g_2)_\eta + (g_1)_\rho &= 0, \\ (f_1)_\eta - (f_2)_\rho &= 0, & (g_1)_\eta - (g_2)_\rho &= 0. \end{aligned} \quad (2.52)$$

From the first we see that $f_3 = (H)_\rho$ and $f_4 = (H)_\eta$, being $H(\rho, \eta)$ certain function. Therefore we can make the coordinate change $(\rho, \eta, \theta, \varphi) \rightarrow (\rho - H, \eta, \theta, \varphi)$ and eliminate f_3 and f_4 . The same holds for g_3 and g_4 and therefore we are dealing with the case described in (2.49). The corresponding hyperkahler metric is then conformal to

$$\tilde{g} = d\rho^2 + d\eta^2 + \frac{(A_0 d\theta - B_0 d\varphi)^2 + (A_1 d\theta - B_1 d\varphi)^2}{(A_0 B_1 - A_1 B_0)^2}, \quad (2.53)$$

which is the same metric as above with $\Omega = 1$. We have redefined $e_1 = A_0$, $e_2 = B_0$, $f_1 = A_1$, $f_2 = B_1$ in (2.53).

There exists another family of hypercomplex structures that can be constructed in terms of holomorphic functions. It is direct to check that Ashtekar equations (2.55) can be cast in the following complex form

$$[e_1 + ie_2, e_1 - ie_2] - [e_3 + ie_4, e_3 - ie_4] = 0, \quad [e_1 + ie_2, e_3 - ie_4] = 0 \quad (2.54)$$

Let M be a complex surface with holomorphic coordinates (z_1, z_2) and let us define four vector fields e_i as

$$\begin{aligned} e_1 + ie_2 &= f_1 \frac{\partial}{\partial z_1} + f_2 \frac{\partial}{\partial z_2} \\ e_3 + ie_4 &= f_3 \frac{\partial}{\partial z_1} + f_4 \frac{\partial}{\partial z_2} \end{aligned}$$

being f_j a complex function on M . Then (2.54) implies that $\partial f_j / \partial \bar{z}_k = 0$ and therefore we can construct an hypercomplex structure using four arbitrary holomorphic functions or two holomorphic vector fields [147].

2.3 Hyperkahler spaces in four dimensions

In a well known work [153] Ashtekar, Jacobson and Smolin introduced a formulation for self-dual manifolds in which they reduced the problem to solve certain quadratic equations for the dual vector fields and a volume form preserving condition. Here we review their construction in the context of hypercomplex structures.

Proposition 3 *Consider an oriented manifold M and four vector fields e_1, e_2, e_3 and e_4 forming an oriented basis for TM at each point. Let us suppose that the fields satisfies the quadratic equations*

$$[e_1, e_2] + [e_3, e_4] = 0, \quad [e_1, e_3] + [e_4, e_2] = 0, \quad [e_1, e_4] + [e_2, e_3] = 0, \quad (2.55)$$

and the volume preserving condition $\mathcal{L}_{e_a}\Theta = 0$ for some 4-form Θ . Then the vectors e_a are conformal to an orthonormal frame of a self-dual Ricci-flat metric.

Before to explain Proposition 3 let us note that

$$[e_1, e_2] + [e_3, e_4] = 0, \quad [e_1, e_3] + [e_4, e_2] = 0, \quad [e_1, e_4] + [e_2, e_3] = 0,$$

is the system (2.55) but with the A_a 's equal to zero. Then (2.43) implies that $\chi = -\alpha$. Then if χ were a gradient, the form α also would be a gradient and there will exist a conformal change taking the condition

$$d\bar{J}^i = \alpha \wedge \bar{J}^i$$

to $d\bar{J}^i = 0$. The corresponding metric will be conformal to an hyperkahler one, which is automatically self-dual Ricci-flat. As we will see, this is essentially the content of the volume preserving condition of proposition 3.

In general the covariant divergence of an arbitrary vector field V is obtained by the formula

$$\mathcal{L}_V\Theta = (\nabla \cdot V)\Theta. \quad (2.56)$$

The divergence of a tetrad $\bar{e}_a$ is given by

$$\nabla \cdot \bar{e}_a = \bar{c}_{ab}^b, \quad (2.57)$$

being $\bar{c}_{ab}^c$ defined by

$$[\bar{e}_b, \bar{e}_c] = \bar{c}_{bc}^a \bar{e}_a.$$

Let us define a factor Ω through

$$\Omega^2 = \Theta(e_1, e_2, e_3, e_4) \quad (2.58)$$

and take the derivative of (2.58) along $\bar{e}_a = \Omega^{-1}e_a$. Using the condition $\mathcal{L}_{e_a}\Theta = 0$ together with (2.56) and (2.57) gives

$$\bar{c}_{ab}^b = -\bar{e}_a(\log \Omega) \quad \Longleftrightarrow \quad \chi = -\bar{e}_a(\log \Omega)\bar{e}^a.$$

The last implication is just the definition (3.326). The transformation rule (2.43) implies that under $g \rightarrow \Omega^{-2}g$ we have

$$\chi_a \rightarrow \chi_a + 3\bar{e}_a \log(\Omega) \quad \Longleftrightarrow \quad \chi = 2\bar{e}_a(\log \Omega)e^a. \quad (2.59)$$

If (2.55) also hold we have

$$\chi = -\alpha \quad \Longleftrightarrow \quad \alpha_a = -2\bar{e}_a(\log \Omega) = -2e_a\left(\frac{1}{\Omega}\right) \quad (2.60)$$

and therefore α is a gradient. Then g is conformal to an hyperkahler metric. The converse of this theorem is also true, more details of this assertions can be found in the original references [153].

It is also possible to cast into this formalism the Plebanski formulation for hyperkahler spaces [76]. By defining the complex vector fields

$$u = e_1 + ie_2, \quad v = e_1 - ie_2, \quad w = e_3 + ie_4, \quad x = e_3 - ie_4,$$

the system (2.55) takes the form

$$[u, w] = 0, \quad [v, x] = 0, \quad [u, v] + [w, x] = 0. \quad (2.61)$$

It follows from the first (2.61) that we can introduce coordinates (t, y) on each of these surfaces such that

$$u = \partial_t, \quad w = \partial_y.$$

Let us select the vector fields

$$v = a_y \partial_x - b_y \partial_z, \quad x = -a_t \partial_x + b_t \partial_z.$$

being a and b unknown functions. Equations (2.61) give us

$$\{a, a_x\} = \{b, a_z\}, \quad \{a, b_x\} = \{b, b_z\}, \quad (2.62)$$

where we have defined the Poisson Bracket by

$$\{f, g\} = f_t g_y - f_y g_t.$$

Let us suppose that the vector fields e_i are divergence free with respect to the volume form $\Theta = dt \wedge dx \wedge dy \wedge dz$. Then it exists a function Ω such that $a = K_z$ and $b = K_y$. By integrating equations (2.64) once, and scaling the coordinates (y, z) , one finds that K satisfies the equation

$$\{K_y, K_z\} = 1, \quad (2.63)$$

which is the First Heavenly equation [76]. This equation can be expressed as

$$\begin{vmatrix} K_{yt} & K_{yx} \\ K_{zt} & K_{zx} \end{vmatrix} = 1.$$

The corresponding hyperkahler metric is

$$g = 4(b_t a_y - a_t b_y)^{-1} [dt \otimes (a_t dz + b_t dx) + dy \otimes (a_y dz + b_y dx)],$$

or in terms of K ,

$$g = K_{ty} dt dy + K_{tz} dt dz + K_{xy} dx dy + K_{xz} dx dz. \quad (2.64)$$

The metrics (2.64) are known as the Plebanski heavens [76] and are the most general hyperkahler examples. An hyperkahler space is also Kahler, and the key function K plays the role of the Kahler potential.

Let us consider metrics with one isometry and introduce a the dual basis $e_a = f_a \partial / \partial x_1 + \partial / \partial x_a$. Then (2.55) reduce to the system

$$\frac{\partial f_1}{\partial x_2} - \frac{\partial f_3}{\partial x_4} + \frac{\partial f_4}{\partial x_3} = 0, \quad \frac{\partial f_1}{\partial x_3} - \frac{\partial f_4}{\partial x_2} + \frac{\partial f_2}{\partial x_4} = 0, \quad \frac{\partial f_1}{\partial x_4} - \frac{\partial f_2}{\partial x_3} + \frac{\partial f_3}{\partial x_2} = 0.$$

Let us assume that $\partial / \partial x^1$ is a Killing vector and let us denote $x_1 = t$. This implies that the functions f_i are independent of t and this system reduces to

$$\nabla V = \nabla \times A \quad \Longleftrightarrow \quad dU = *dA \quad (2.65)$$

which implies that $\Delta V = 0$ and we have defined $V = f_1$ and $A = (f_2, f_3, f_4)$. The corresponding metric is

$$g = V^{-1}(dt - A)^2 + V dx \cdot dx, \quad (2.66)$$

and it can be checked that is indeed Ricci-flat.

A particular solution of (2.65) is

$$(V(x))^{-1} = a + \sum_{i=1}^n 2m_i |r - r_i|^{-1}$$

being a and m_i constants and r_i the position of certain points on the manifold. If we select $m_i = M$ and t periodic in the range $0 \leq t \leq \frac{8\pi M}{n}$ then the singularities in $r = r_i$ are removable. Such spaces have finite energy and therefore are gravitational instantons. The metrics corresponding $a = n = 1$ are known as the Taub-Nut metrics. If $a = 0$ the metrics are known as multi-instantons and if $n = 2$ we obtain the Eguchi-Hanson instanton [126].

Metrics (2.66) are known as Gibbons-Hawking metrics [154], although they also were considered independently by Kloster, Son and Das [73]. They are the four dimensional self-dual examples that are Ricci-flat with a $U(1)$ -isometry for which condition $\mathcal{L}_{\partial_t} J^a = 0$ is satisfied. Such isometries are called tri-holomorphic. Another characterization of (2.66) is that they have self-dual Killing vector K_i , i.e, the tensor $K_{ij} = \nabla_i K_j$ is self-dual [60], [74]. We will return to this point in the next subsection.

2.4 Quaternion Kahler and hyperkahler metrics in $d = 4$ with at least one isometry

This section present the most general local form of a metric belonging to a self-dual conformal structure and at least one Killing vector. The spaces in consideration are four dimensional. Although this results are not new, we present them in an slightly different way than the standard one. We show the relation between self-dual structures with a Killing vector and Einstein-Weyl ones, which satisfy a generalization of the Einstein equations including conformal transformations, i.e, Einstein-Weyl equations are invariant under the group $CO(n) = R_+ \times SO(n)$ [150], [157]. A geometric interpretation for the integrability of the non linear axial $SU(\infty)$ Toda equation will be also presented. The argument is that the axial Toda equation describe an hyperkahler space of the Gibbons-Hawking type (which are entirely described in terms of a linear equation) in a coordinate system that hides its linear nature.

We should remark that $U(1) \times U(1)$ quaternion Kahler metrics are important for the goals of the present work. By extending them to spaces of G_2 , $Spin(7)$ holonomy and higher dimensional hyperkahler ones, we constructed certain classical solutions of M-theory preserving some amount of supersymmetries. The presence of isometries allows to find type IIA and IIB backgrounds by use of dualities. We will discuss it in the next sections.

2.4.1 Hyperkahler metrics with Killing vectors that are not self-dual

The most general form of an hyperkahler metric is given by (2.64) and defined in terms of a Kahler potential K satisfying the non linear equation (2.63). It will be convenient for the following to introduce two complex coordinates v, w and write the heavenly metrics (2.64) in complex form as

$$g = K_{vw}dvdw + K_{v\bar{w}}dv d\bar{w} + K_{\bar{v}w}d\bar{v}dw + K_{\bar{v}\bar{w}}d\bar{v}d\bar{w} \quad (2.67)$$

where bar indicate complex conjugation. Because Plebanski metrics are general it follows that the Gibbons-Hawking metrics (2.66) should arise as a subcase of (2.67). Such reduction goes as follows. Select the key function K as

$$K(v, w, \bar{v}, \bar{w}) = K(i(\bar{w} - w), v, \bar{v}) \quad (2.68)$$

If one define $u = i(\bar{w} - w)$, then equation (2.63) becomes

$$K_{uu}K_{v\bar{v}} - K_{uv}K_{u\bar{v}} = 1. \quad (2.69)$$

Consider the new independent variables $x, v, \bar{v}$ with $x = K_u$. By setting

$$H(x, v, \bar{v}) = xu - K \quad (2.70)$$

the heavenly equation (2.75) reduce to

$$H_{xx} + H_{yy} + H_{zz} = 0 \quad (2.71)$$

being $y = v + \bar{v}$ and $iz = v - \bar{v}$. The metric (2.67) in this coordinate system take the form

$$g = H_{xx}(dx^2 + dy^2 + dz^2) + H_{xx}^{-1}(dt + H_{xy}dz - H_{xz}dy)^2, \quad (2.72)$$

being $it = \bar{v} - v$. By selecting $H_x = G$ we can rewrite (2.72) in the more familiar form

$$g = G_x(dx^2 + dy^2 + dz^2) + G_x^{-1}(dt + G_ydz - G_zdy)^2, \quad (2.73)$$

By identifying $G_x = V$ and $A = G_y dz - G_z dy$ we find that $\nabla \times A = \nabla V$. Then (2.73) are the Gibbons-Hawking metrics. The Killing vector for such metric is $\partial/\partial t$ and it is called translational in the literature [60].

There exist another type of hyperkahler metrics with one Killing vector, and related to certain limit of a Toda equation. Let us select the key function K as

$$K(v, w, \bar{v}, \bar{w}) = K(\log |w|^2, v, \bar{v}) \quad (2.74)$$

By choosing $F = \log |w|^2$ equation (2.63) becomes

$$K_{uu}K_{v\bar{v}} - K_{uv}K_{u\bar{v}} = e^F. \quad (2.75)$$

Consider the new independent variables $t, v, \bar{v}$ with $\tau = K_F$. If we set

$$H(\tau, v, \bar{v}) = \tau F - K \quad (2.76)$$

then equation (2.75) is converted into

$$H_{xx} + H_{yy} + (e^{H_\tau})_\tau = 0 \quad (2.77)$$

with $x = v + \bar{v}$ and $iy = v - \bar{v}$. Defining $u = H_\tau$ and $z = \tau$ and differentiating (2.77) with respect to z gives

$$(e^u)_{zz} + u_{yy} + u_{xx} = 0. \quad (2.78)$$

Equation (2.78) is known as the continuum limit of the $sl(n)$ Toda equation. The metric (2.67) is expressed in terms of the new coordinates as

$$g = u_z[e^u(dx^2 + dy^2) + dz^2] + u_z^{-1}[dt + (u_x dy - u_y dx)]^2, \quad (2.79)$$

being u a solution of the $SU(\infty)$ Toda equation (2.78). We have defined the coordinate $it = \log(\frac{w}{\bar{w}})$ and the Killing vector is $\partial/\partial t = \partial/\partial(\arg w)$. This explain why this Killing vectors are called rotational in the literature.

A result due to Boyer and Finley [75] show that both (2.79) and (2.66) exhaust all the possible metrics with at least one isometry. The first are the most general ones with self-dual Killing vector, the second are those with a non self-dual isometry. Metrics (2.66) are related to a linear equation and therefore are more easy to construct. Instead the continuum Toda equation (2.78) is integrable but not many explicit solutions are known. One interesting solution is those that is factorized by $u = f(x, y) + g(z)$, then there appears three solutions given by

$$\begin{aligned} e^{u_1} &= \frac{1}{2}(z^2 - a^2)[1 + \frac{1}{8}(x^2 + y^2)]^{-2}, & z^2 &\geq a^2 \\ e^{u_2} &= \frac{1}{2}(z^2 + a^2)[1 + \frac{1}{8}(x^2 + y^2)]^{-2}, & z^2 &\geq a^2 \\ e^{u_3} &= \frac{1}{2}(a^2 - z^2)[1 + \frac{1}{8}(x^2 + y^2)]^{-2}, & z^2 &\leq a^2 \end{aligned}$$

The first of this solutions corresponds to the Eguchi-Hanson metric. This can be visualized defining $\sigma = 1 + \frac{1}{8}(x^2 + y^2)$ and the coordinate system (r, θ, φ) given by

$$x\sigma^{-1} = \sqrt{2}\cos(\varphi)\sin(\theta),$$

$$y\sigma^{-1} = \sqrt{2}\sin(\varphi)\sin(\theta),$$

$$4z = r^2,$$

then the metric (2.79) becomes the standard Eguchi-Hanson, namely

$$g = (1 - \frac{\alpha^4}{r^4})^{-1}dr^2 + \frac{r^2}{4}(d\theta^2 + \sin(\theta)^2d\varphi^2) + \frac{r^2}{4}(1 - \frac{\alpha^4}{r^4})(\cos(\theta)d\varphi + dt)^2 \quad (2.80)$$

being $\alpha^4 = 16a^2$. It is seen that there is another Killing vector $\partial/\partial\varphi$. There exists a coordinate change that bring this metric to the form of Hawking multi-instanton solution. Therefore $\partial/\partial\varphi$ is self-dual but $\partial/\partial t$ is not. The metrics corresponding to the other solutions u_2 and u_3 have singular points for the scalar curvature which cannot be excluded from the space time. Therefore this spaces are not gravitational instantons [74].

2.4.2 Integrability of the axial continuum Toda equation

The Eguchi-Hanson metric is an example possessing simultaneously a self-dual Killing vector and a not self-dual one (2.80). Such examples are described simultaneously by the non linear equation (2.78) for certain election of coordinates, and by the linear equation (2.65) for other coordinates. The transformation from one of such coordinates into the other is a mapping between the axial $SU(\infty)$ Toda equation and the so called Ward monopole equation, which is linear. Therefore although the continuum Toda equation is not linear, in the axial case it is completely equivalent to a linear one. By axial Toda we mean the equation

$$(e^u)_{zz} + u_{xx} = 0. \quad (2.81)$$

The geometrical origin of the equivalence stated above is that the axial Toda equation describe Gibbons-Hawking metrics in a coordinate system which hide its linear nature [129].

To see this statement more concretely let us take a Gibbons-Hawking metric (2.66) and write the flat 3-dimensional part in cylindrical polar coordinates (η, ρ, φ) . The result is

$$g_f = dx^2 + dy^2 + dz^2 = d\eta^2 + d\rho^2 + \rho^2 d\varphi^2.$$

Let us also suppose that the generator of the axial symmetry $\partial/\partial\varphi$ is also a Killing vector, then $V(\rho, \eta)$ in (2.66) is independent on the φ coordinate and the equation (2.66) becomes the Ward monopole equation, namely

$$\rho^{-1}(\rho V_\rho)_\rho + V_{\eta\eta} = 0. \quad (2.82)$$

Note that if $\tilde{V}$ is a solution of (2.82), so is $\tilde{V}_\eta$, and $\tilde{V}_\eta$ determines $\tilde{V}$ up to the addition of $C_1 \log(C_2\rho)$ for some constants C_1 and C_2 . This provides a way of integrating the equation $d*dV = 0$ to give $*dV = dA$. If we choose $V = \tilde{V}_\eta$, then we can take $A = \rho\tilde{V}_\rho d\phi$. The corresponding hyperkahler metric is

$$g = V_\eta(d\eta^2 + d\rho^2 + \rho^2 d\phi^2) + V_\eta^{-1}(d\theta + \rho V_\rho d\phi)^2, \quad (2.83)$$

where have reexpressed $\tilde{V}$ as V in order to simplify the notation.

The metric (2.83) has $U(1) \times U(1)$ isometry and the two commuting Killing vectors are $\partial/\partial\varphi$ and $\partial/\partial\theta$. By construction $\partial/\partial\theta$ is self-dual. But $\partial/\partial\varphi$ is not self-dual in general. Therefore

there exist a coordinate system (x, y, z, φ) for which (2.83) take the form (2.79). We can rewrite (2.83) by completing squares as

$$g = V_\eta \left(d\rho^2 + d\eta^2 + \frac{1}{V_\eta^2 + V_\rho^2} d\theta^2 \right) + \frac{\rho^2(V_\eta^2 + V_\rho^2)}{V_\eta} \left(d\varphi + \frac{V_\rho}{\rho(V_\eta^2 + V_\rho^2)} d\theta \right)^2.$$

or equivalently as

$$\begin{aligned} g = & \frac{V_\eta}{\rho^2(V_\eta^2 + V_\rho^2)} [\rho^2(V_\eta^2 + V_\rho^2)(d\rho^2 + d\eta^2) + \rho^2 d\psi^2] \\ & + \frac{\rho^2(V_\eta^2 + V_\rho^2)}{V_\eta} \left(d\varphi + \frac{V_\rho}{\rho(V_\eta^2 + V_\rho^2)} d\theta \right)^2. \end{aligned} \quad (2.84)$$

It is extremely important to recall that (2.83) and (2.84) are the *same* metric. Comparison between (2.79) and (2.81) shows that

$$u_z = \frac{V_\eta}{\rho^2(V_\eta^2 + V_\rho^2)} \quad (2.85)$$

and that there it should exists a coordinate system (x, y, z) for which

$$\rho^2(V_\eta^2 + V_\rho^2)(d\rho^2 + d\eta^2) + \rho^2 d\theta^2 = e^u(dx^2 + dy^2) + dz^2 \quad (2.86)$$

with u satisfying (2.81). The system (x, y, z) is obtained as follows. There are two exact differentials constructed with V , namely

$$dV = V_\rho d\rho + V_\eta d\eta,$$

$$dG = \rho V_\rho d\eta - \rho V_\eta d\rho.$$

The equation $d^2V = 0$ is identically satisfied and $d^2G = 0$ is a consequence of the equation (2.82), therefore dV and dG are truly differentials. The function G is defined by V through the Backlund transformation

$$\rho V_\rho = G_\eta, \quad \rho V_\eta = -G_\rho.$$

It is elementary to show that

$$\rho^2(V_\eta^2 + V_\rho^2)(d\rho^2 + d\eta^2) + \rho^2 d\theta^2 = \rho^2(dV^2 + d\psi^2) + dG^2.$$

Therefore

$$\rho^2(dV^2 + d\theta^2) + dG^2 = e^u(dx^2 + dy^2) + dz^2$$

This means that $y = \theta$ and a Toda solution is defined through the equation $e^u = \rho^2$. We have obtained the following proposition [129].

Proposition 4 *Any solution V of the equation $V_\eta + \rho^{-1}(\rho V_\rho)_\rho = 0$ defines locally the coordinate system (x, z)*

$$x = V_\eta, \quad z = \rho V_\rho, \quad (2.87)$$

in terms of (ρ, η) and conversely (2.87) defines implicitly (ρ, η) as functions of (x, z) . Then the function $u(x, z) = \log(\rho^2)$ is a solution of the axially symmetric Toda equation

$$(e^u)_{zz} + u_{xx} = 0. \quad (2.88)$$

This procedure can be inverted in order to find a Ward monopole V starting with a given Toda solution.

Proposition 4 gives a method to find solutions of a non linear equation (the continuum Toda one) starting with a solution of a linear one (the Ward equation) and shows the equivalence that we were looking for. Proposition 4 can be checked by an elementary (but lengthy) chain rule exercise. It is difficult in practice to find explicit solutions of (2.88) and usually proposition 4 gives implicit solutions.

2.4.3 Einstein-Weyl structures and hyperkahler metrics

An important feature about the three dimensional part

$$\gamma = e^u(dx^2 + dy^2) + dz^2, \quad (2.89)$$

is that it is an Einstein-Weyl structure. Einstein-Weyl condition is a generalization of the ordinary Einstein condition in which the Lorentz group $SO(4)$ is extended to the conformal group $CO(4) = R_+ \times SO(4)$ [150]³. The Einstein equations are generalized in this picture in order to include invariance under coordinate rotations plus dilatations. In this section some important facts about them are sketched following [150] and [157].

It is known that for a given space $\overline{W}$ endowed with a metric $g_{\mu\nu}$ the Levi-Civita connection ∇ is uniquely defined by

$$\nabla g = 0, \quad T(\nabla) = 0 \quad (2.90)$$

where $T(\nabla)$ is the torsion. A Weyl-structure is defined by the manifold $\overline{W}$ together with:

(a) A class of conformal metrics $[g]$, whose elements are related by the conformal scaling (or gauge transformation)

$$g_{\mu\nu} \rightarrow \Omega^2 g_{\mu\nu}, \quad (2.91)$$

together with $SO(n)$ -coordinate transformations. Ω^2 is an smooth, positive real function over $\overline{W}$.

(b) A torsion-free connection D_μ which acts over a representative g_{ab} of the conformal class $[g]$ as

$$D_\mu g_{\nu\alpha} = \omega_\mu g_{\nu\alpha}, \quad (2.92)$$

for certain functions ω_μ defining an one form ω . Then it is said that D preserves $[g]$.

The conformal group in n -dimensions is $CO(n) = R_+ \times SO(n)$ and includes rotations plus general scale transformations. The structure $[g]$ is called $CO(n)$ -structure over $\overline{W}$. The connection D is uniquely determined by (2.92) in terms of ω and g . This can be seen in a coordinate basis ∂_k in which the system (2.92) takes the form

$$g_{\mu\nu,\alpha} = g_{\beta\nu} \Upsilon_{\mu\alpha}^\beta + g_{\mu\beta} \Upsilon_{\nu\alpha}^\beta + \omega_\mu g_{\nu\alpha},$$

³Usually the conformal group is defined by the angle-preserving (conformal) transformations. We are not using this definition here. We are considering only $SO(4)$ transformations and dilatations

where the symbols $\Upsilon_{\mu\alpha}^\beta$ denotes the connection coefficients of D , which are symmetric in the lower indices by the torsionless condition. Thus a series of steps analogous to those needed to determine the Levi-Civita connection shows that $\Upsilon_{\mu\alpha}^\beta$ is

$$\Upsilon_{\mu\alpha}^\beta = \Gamma_{\mu\alpha}^\beta + \gamma_{\mu\alpha}^\beta \quad (2.93)$$

where $\Gamma_{\mu\alpha}^\beta$ are the Christoffel symbols and the add $\gamma_{\mu\alpha}^\beta$ is

$$2\gamma_{\mu\alpha}^\beta = (\delta_\mu^\beta \omega_\alpha + \delta_\alpha^\beta \omega_\mu + g_{\mu\alpha} \omega^\beta). \quad (2.94)$$

The form ω is not invariant under (2.95), its transformation law can be obtained from (2.92) and (2.94) and is

$$\omega_\mu \rightarrow \omega_\mu + 2\partial_\mu \log(\Omega). \quad (2.95)$$

It follows from (2.94) and (2.95) that the Levi-Civita of any g of $[g]$ preserves the conformal structure.

If for a Weyl-structure the symmetric part of the Ricci tensor $\tilde{R}_{(\mu\nu)}$ constructed with D_μ satisfies

$$\tilde{R}_{(\mu\nu)} = \Lambda g_{\mu\nu} \quad (2.96)$$

for certain Λ , then will be called Einstein-Weyl. If in addition the antisymmetric part of $\tilde{R}_{[\mu\nu]}$ vanish there exists a gauge in which (2.96) reduces to the vacuum Einstein equation with cosmological constant Λ . To see this it is needed to calculate

$$\tilde{R} = D_X D_Y - D_Y D_X - D_{[X,Y]}$$

using the formula (2.93) for D . The relation $CO(4) = R_+ \times SO(4)$ decompose $\tilde{R}$ into a real component R_0 and into an $SO(4)$ -component R that is equal to the curvature tensor constructed with ∇ . After contracting indices it is obtained

$$\tilde{R}_{\mu\nu} = R_{\mu\nu} + \nabla_\mu \omega_\nu - \frac{1}{2} \nabla_\mu \omega_\nu - \frac{1}{4} \omega_\mu \omega_\nu + g_{\mu\nu} \left(\frac{1}{2} \nabla_\alpha \omega^\alpha + \frac{1}{4} \omega_\alpha \omega^\alpha \right),$$

where $R_{\mu\nu}$ is the Ricci tensor found with ∇ . The antisymmetric part is originated by the R_0 component and is determined in terms of ω as

$$\tilde{R}_{[\mu\nu]} = \frac{3}{2} \nabla_{[\mu} \omega_{\nu]}. \quad (2.97)$$

If (2.97) is zero, then ω is the gradient of certain function Ψ over $\overline{W}$. The conformal rescaling (2.95) with $\Omega = -e^\Psi$ set $\omega = 0$. This reduce the symmetric part

$$\tilde{R}_{(\mu\nu)} = R_{\mu\nu} - \frac{1}{2} \nabla_{(\mu} \omega_{\nu)} - \frac{1}{4} \omega_\mu \omega_\nu$$

to $R_{\mu\nu}$ and (2.96) is the Einstein equation with Λ , thus $[g]$ contains an Einstein metric.

As we recalled in the last subsection, the metric

$$\gamma = e^u(dx^2 + dy^2) + dz^2,$$

is the representative of an Einstein-Weyl structure with the factor ω given by

$$\omega = -u_z dz. \quad (2.98)$$

Therefore the Gegenberg and Das hyperkahler metrics (2.79) are constructed with a three dimensional basis that is Einstein-Weyl. It can also be seen that if we identify

$$V = u_z, \quad A = u_x dy - u_y dx,$$

then (2.79) take the form

$$g = V[e^u(dx^2 + dy^2) + dz^2] + \frac{(dt + A)^2}{V},$$

and it is seen from (2.78) that the pair (V, A) satisfy

$$*dA = dV + \frac{\omega}{2}V. \quad (2.99)$$

The Hodge star $*$ is taken with respect to the Einstein-Weyl structure. Equation (2.99) is the conformal generalization of the Gibbons-Hawking one $*dA = dV$ and therefore it will be called generalized monopole equation.

2.4.4 Self-dual structures with one Killing vector

In this subsection we show that the generalized monopole equation (2.99) is conformally invariant and that it describe all the self-dual conformal structures with one conformal Killing vector. Let us consider a metric with the form

$$g = Vh + \frac{(dt + A)^2}{V},$$

being h an Einstein-Weyl structure and the functions (V, A) satisfying (2.99). Then if we perform the conformal change

$$\tilde{h} = \Omega^{-2}h,$$

we can express g as

$$g = Vh + \frac{(dt + A)^2}{V} = \Omega\tilde{g} \quad (2.100)$$

where $\tilde{V} = V\Omega$. We want to prove that if (2.99) holds then

$$*_h dA = d\tilde{V} + \frac{\tilde{\omega}}{2}\tilde{V}$$

being $\tilde{\omega}$ the conformal factor associated to $\tilde{h}$. If we express the conformal transformation (2.95) for ω as

$$\Omega\tilde{\omega} = \Omega\omega - 2d\Omega$$

then it is easily seen that

$$d\tilde{V} + \frac{\tilde{\omega}}{2}\tilde{V} = \Omega(dV + \frac{\omega}{2}V)$$

and from (2.99) and the last expression we have

$$d\tilde{V} + \frac{\tilde{\omega}}{2}\tilde{V} = \Omega *_h dA. \quad (2.101)$$

By another side we can put

$$dA = A_{ab}e^a \wedge e^b \implies *_h dA = \epsilon_{abc}A_{ab}e^c,$$

where we denote as A_{ab} the coefficients of dA in the basis $e^a \wedge e^b$, being e^a the einbein corresponding to h . Then

$$\Omega *_h dA = \epsilon_{abc}A_{ab}\Omega e^c = \epsilon_{abc}A_{ab}\Omega^2 \tilde{e}^c$$

being $\tilde{e}^c$ the einbein corresponding to $\tilde{h}$, namely $\tilde{e}^c = e^c/\Omega$. Therefore

$$*_h(\Omega *_h dA) = A_{ab}\Omega^2 \tilde{e}^a \wedge \tilde{e}^b = A_{ab}e^a \wedge e^b = dA.$$

Using the fact that $*^2 = 1$ from the last formula we obtain

$$\Omega *_h dA = *_h dA. \quad (2.102)$$

From (2.101) and (2.102) we see that (2.99) implies

$$*_h dA = d\tilde{V} + \frac{\tilde{\omega}}{2}\tilde{V}$$

which implies that (2.99) is conformally invariant.

We have also seen that equation (2.99) also describe metrics with self-dual curvature as in (2.79) or (2.66). It is evident from the result of this subsection that conformal family $[g]$ associated to (2.79) and (2.66) have self-dual Weyl tensor and a conformal Killing vector, and that all their elements are described with an equation of the form (2.99). This raise the question whether or not *any* self-dual family with one conformal Killing vector has the form

$$g = Vh + \frac{(dt + A)^2}{V},$$

being h an *arbitrary* Einstein-Weyl structure and (V, A) a pair satisfying (2.99). The answer to this question is positive, it was proved by Jones and Tod and is stated in the following proposition [157].

Proposition 5 *a) Consider an Einstein-Weyl structure $[h]$ in $D=3$ and a representative h . Then the four dimensional metric*

$$g = Vh + \frac{(dt + A)^2}{V} \quad (2.103)$$

is self-dual with one Killing vector ∂_t if the pair of functions (V, A) satisfies the generalized monopole equation

$$dA = *_h(dV - V\omega). \quad (2.104)$$

*The Hodge star $*_h$ is taken with respect to $h_{\mu\nu}$ and ω is the conformal factor corresponding to h .*

b) Conversely given a metric g that is self-dual and has one conformal Killing vector K^a then a conformal transformation can be performed in order that K^a becomes a Killing vector ∂_t and there exists a system of coordinates in which g takes the form (2.103), being h a representative of an Einstein-Weyl structure. The factor ω will be obtained in this case through

(2.104).

Some comments are in order. The Gegenberg and Das metrics (2.79) and (2.66) generates all the families conformal to an hyperkahler metric. For all the space over a Toda base we have

$$dV - \omega V = V_x dx + V_y dy + (V_z + u_z V) dz = V_x dx + V_y dy + e^{-u} (e^u V)_z dz,$$

and (2.104) is then explicitly

$$dA = *_h(dV - \omega V) = V_x dz \wedge dy + V_y dx \wedge dz + (Ve^u)_z dy \wedge dx.$$

Therefore the integrability condition for the existence of A is

$$(Ve^u)_{zz} + V_{yy} + V_{xx} = 0. \quad (2.105)$$

In the next subsection we will show an application of this correspondence, the construction of all the self-dual structures with two commuting Killing vectors that are surface orthogonal. This are the Joyce spaces.

2.4.5 The Joyce spaces

Let us focus in self-dual families $[h]$ with two commuting $U(1)$ isometries. Consider an structure $[g]$ over M with representatives g that locally takes the form (2.47), or equivalently

$$g = (A_0 B_1 - A_1 B_0)(d\rho^2 + d\eta^2) + \frac{(A_0 d\theta - B_0 d\varphi)^2 + (A_1 d\theta - B_1 d\varphi)^2}{(A_0 B_1 - A_1 B_0)}. \quad (2.106)$$

We made by convenience a conformal transformation in (2.47) with factor $\Omega = A_0 B_1 - A_1 B_0$, we can always do this if we are dealing with a self-dual structure due to the conformal invariance the Weyl tensor. The unknown functions A_i and B_i are supposed to be independent on the coordinates (θ, φ) . By construction $\partial/\partial\theta$ and $\partial/\partial\varphi$ are commuting Killing vectors of the metric (2.106).

We will fix the functions A_i and B_i in order that (2.47) will be self-dual. The Jones-Tod correspondence insure that there should exists a coordinate system (x, y, z, t) defined in terms of the old one $(\rho, \eta, \theta, \varphi)$ for which (2.106) is conformal to

$$g = Vh + \frac{1}{V}(dt + A)^2 \quad (2.107)$$

according to (2.103). To find this system of coordinates we write (2.106) by completing squares as

$$\begin{aligned} g = & \frac{A_0 B_1 - A_1 B_0}{\rho^2 (A_0^2 + A_1^2)} ((A_0^2 + A_1^2)(d\rho^2 + d\eta^2) + \rho^2 d\varphi^2) \\ & + \frac{A_0^2 + A_1^2}{A_0 B_1 - A_1 B_0} \left(d\theta - \frac{(A_0 B_0 + A_1 B_1) d\varphi}{A_0^2 + A_1^2} \right)^2 \end{aligned} \quad (2.108)$$

and is seen that after scaling by ρ and identifying $t = \theta$ that it takes the form (2.103) with a metric h and a monopole (V, A) given by

$$h = (A_0^2 + A_1^2)(d\rho^2 + d\eta^2) + \rho^2 d\varphi^2, \quad V = \frac{A_1 B_0 - A_0 B_1}{\rho(A_0^2 + A_1^2)}, \quad (2.109)$$

$$A = -\frac{(A_0B_0 + A_1B_1)}{A_0^2 + A_1^2}d\varphi. \quad (2.110)$$

Comparison of the first (2.109) with (2.86) and Proposition 4 shows the identification

$$A_0 = \rho\tilde{V}_\eta, \quad A_1 = -\rho\tilde{V}_\rho$$

being $\tilde{V}$ a solution of the Ward monopole equation (2.82). Therefore A_i should satisfy

$$(A_0)_\eta - (A_1)_\rho = 0,$$

and from the identity $\tilde{V}_{\rho\eta} = \tilde{V}_{\eta\rho}$ we have the integrability condition

$$\left(\frac{A_0}{\rho}\right)_\rho = -\left(\frac{A_1}{\rho}\right)_\eta$$

or equivalently

$$(A_0)_\rho + (A_1)_\eta = A_0/\rho.$$

Clearly the same reasoning holds for the B_i and therefore we have obtained part of the following result due to Joyce [147].

Proposition 6 *Consider a set of functions A_i and B_i with $i = 1, 2$, satisfying*

$$(A_0)_\rho + (A_1)_\eta = A_0/\rho, \quad (2.111)$$

$$(A_0)_\eta - (A_1)_\rho = 0, \quad (2.112)$$

and the same equations for B_i . Then the family $[g_j]$ associated to the metric g_j given by

$$g_j = (A_0B_1 - A_1B_0)\frac{d\rho^2 + d\eta^2}{\rho^2} + \frac{(A_0d\theta - B_0d\varphi)^2 + (A_1d\theta - B_1d\varphi)^2}{(A_0B_1 - A_1B_0)}, \quad (2.113)$$

is self-dual. All self-dual metrics with two commuting surface orthogonal Killing vectors arise locally in this way.

Equation (2.112) implies the existence of a potential function G such that

$$A_0 = G_\rho; \quad A_1 = G_\eta. \quad (2.114)$$

Then (2.111) implies that $G_{\rho\rho} + G_{\eta\eta} = G_\rho/\rho$. Conversely, as we have seen above, (2.111) implies that

$$A_0 = -\rho V_\eta; \quad A_1 = \rho V_\rho, \quad (2.115)$$

and (2.112) gives the Ward monopole equation [129]

$$V_{\eta\eta} + \rho^{-1}(\rho V_\rho)_\rho = 0. \quad (2.116)$$

The relations

$$G_\rho = -\rho V_\eta; \quad G_\eta = \rho V_\rho, \quad (2.117)$$

constitute a Backlund transformation allowing to find a monopole V starting with a known G or viceversa. The functions B_i can be also expressed in terms of another potential functions G' and V' satisfying the same equations than V and G .

It should be reminded that we have presented till now two different types of self-dual families with two commuting Killing vectors, namely (2.113) and (2.47). The equations (2.111) and (2.112) are different to the Cauchy-Riemann one (2.50), and is simple to check that there does not exist a coordinate change converting (2.111) and (2.112) into (2.50). Therefore (2.113) and (2.47) are not related by a coordinate transformation. The first are the most general self-dual structures with two commuting Killing vector that are surface-orthogonal [147], [132]. This mean that the manifold M corresponding to (2.113) is of the form $M = N \times T^2$, being T^2 the two dimensional torus. Instead the families (2.47) admits a T^2 action but has no compatible product structure $M = N \times T^2$.

There exist a theorem given in [132] that shows that the Killing vector of a quaternion Kahler space are surface orthogonal. Therefore in order to find the toric quaternion Kahler spaces we should impose the Einstein condition on (2.113) and not on (2.47). This is the purpose of the next subsection.

2.4.6 Identification of the quaternionic-Kahler metrics with at least one isometry

We have shown in the previous subsection a notable one to one correspondence between self-dual structures with one isometry and 3-dimensional Einstein-Weyl structures. A generalization of this result to the Einstein self-dual metrics was obtained independently in [148], [82]. The result is stated in the following proposition.

Proposition 7 *For any self-dual Einstein metric g with one Killing vector in $D=4$ there exist a system of coordinates (x, y, z, t) for which takes the form*

$$g = \frac{1}{z^2} [V(e^u(dx^2 + dy^2) + dz^2) + \frac{1}{V}(dt + A)^2]. \quad (2.118)$$

The functions (V, A, u) are independent of the variable t and satisfies

$$(e^u)_{zz} + u_{yy} + u_{xx} = 0, \quad (2.119)$$

$$dA = V_x dz \wedge dy + V_y dx \wedge dz + (Ve^u)_z dy \wedge dx, \quad (2.120)$$

$$V = 2 - zu_z. \quad (2.121)$$

Conversely, any solution of (2.119), (2.119) and (2.121) define by (2.118) a self-dual Einstein metric.

If the condition (2.121) is relaxed then proposition 7 reduces to the proposition 5 up to an scaling by $1/z^2$. Then (2.121) is the condition to be satisfied in order to have an Einstein metric. It is sufficient because the integrability condition

$$(Ve^u)_{zz} + V_{yy} + V_{xx} = 0,$$

is always satisfied for $V = 2 - zu_z$. In other words, every $SU(\infty)$ Toda solution define a self-dual metric by (2.118).

Proposition 7 is describe the local form of a quaternion Kahler space with one isometry. Quaternion Kahler spaces with two commuting Killing are a subcase of (2.118). Equation (2.119) show us that the family $[g]$ associated to such quaternion Kahler metrics are of the Toda type. Therefore in order to find the quaternion Kahler metrics with $U(1) \times U(1)$ isometry we should apply proposition 7 to the metrics of proposition 6 and not to (2.47).

To determine which g among the Joyce metrics (2.113) satisfy $R_{ij} \sim g_{ij}$ we apply condition (2.121) to them. For Joyce spaces we can calculate the factor $\omega = -u_z dz$ by use of $*dA = dV - \omega V$, (2.109) and (2.110). The result is

$$u_z = \frac{A_0}{\rho(A_0^2 + A_1^2)}.$$

Then we have

$$2 - zu_z = \frac{\rho(A_0^2 + A_1^2) - GA_0}{\rho(A_0^2 + A_1^2)}. \quad (2.122)$$

Insertion of the expression for V (2.109) into the Einstein condition $V = 2 - zu_z$ and taking into account (2.122) gives

$$A_1 B_0 - A_0 B_1 = \rho(A_0^2 + A_1^2) - GA_0. \quad (2.123)$$

From (2.123) it is obtained that $B_0 = \rho A_1 + \xi_0$ and $B_1 = G - \rho A_0 + \xi_1$ with $A_1 \xi_0 = A_0 \xi_1$. The functions ξ_i are determined by the requirement that B_i also satisfy the Joyce system (2.111) and (2.112), the result is $\xi_0 = -\eta A_0$ and $\xi_1 = -\eta A_1$. Therefore the metric $g_j/\rho z^2$ is Einstein if and only if

$$A_0 = G_\rho; \quad A_1 = G_\eta \quad (2.124)$$

$$B_0 = \eta G_\rho - \rho G_\eta; \quad B_1 = \rho G_\rho + \eta G_\eta - G. \quad (2.125)$$

Defining $G = \sqrt{\rho}F$ it follows that F satisfies

$$F_{\rho\rho} + F_{\eta\eta} = \frac{3F}{4\rho^2}.$$

Then inserting (2.124) and (2.125) expressed in terms of F into $g_j/\rho z^2$ and making the identification $z = G$ gives the following assertion due to Calderbank and Pedersen [132].

Proposition 8 *For any Einstein-metric with self-dual Weyl tensor and nonzero scalar curvature possessing two linearly independent commuting Killing fields there exists a coordinate system in which the metric g has locally the form*

$$g = \frac{F^2 - 4\rho^2(F_\rho^2 + F_\eta^2)}{4F^2} \frac{d\rho^2 + d\eta^2}{\rho^2} + \frac{[(F - 2\rho F_\rho)\alpha - 2\rho F_\eta \beta]^2 + [(F + 2\rho F_\rho)\beta - 2\rho F_\eta \alpha]^2}{F^2[F^2 - 4\rho^2(F_\rho^2 + F_\eta^2)]}, \quad (2.126)$$

where $\alpha = \sqrt{\rho}d\theta$ and $\beta = (d\varphi + \eta d\theta)/\sqrt{\rho}$ and $F(\rho, \eta)$ is a solution of the equation

$$F_{\rho\rho} + F_{\eta\eta} = \frac{3F}{4\rho^2}. \quad (2.127)$$

on some open subset of the half-space $\rho > 0$. On the open set defined by $F^2 > 4\rho^2(F_\rho^2 + F_\eta^2)$ the metric g has positive scalar curvature, whereas for $F^2 < 4\rho^2(F_\rho^2 + F_\eta^2)$ $-g$ is self-dual with negative scalar curvature.

The Einstein condition $R_{ij} = \kappa g_{ij}$ is not invariant under scale transformations, so Proposition 8 gives all the quaternionic-Kahler metrics with T^2 isometry up to a constant multiple. The problem to find them is reduced to find an F satisfying the linear equation (2.127), that is, an eigenfunction of the hyperbolic laplacian with eigenvalue $3/4$.

The equation for V (2.116) and has solutions of the integral form

$$V_1(\rho, \eta) = W(\eta, i\rho) + c.c., \quad V_2(\rho, \eta) = W(i\eta, \rho) + c.c.$$

$$W(\eta, \rho) = \frac{1}{2\pi} \int_0^{2\pi} H(\rho \sin(\theta) + \eta) d\theta \quad (2.128)$$

where $H(z)$ is an arbitrary function of one variable [129]. The Backlund relations (2.117) define V in terms of G , and consequently in terms of F , and viceversa. For instance, non trivial eigenfunctions F can be constructed selecting an arbitrary $H(z)$, performing the integration (2.128) and finding G through (2.117), then $F = G/\sqrt{\rho}$.

2.4.7 Examples of toric quaternion Kahler spaces

The following are quaternionic metrics constructed starting with an arbitrary function $G(x)$.

A. The trivial four dimensional toric metric

$$ds^2 = d\rho^2 + d\eta^2 + \rho^2 d\phi^2 + d\psi^2$$

corresponds to the monopole

$$V = \eta$$

(This AHF holds using $G(x) = x \log(x)$ in the integral expression (2.128)). The equations (2.117) are in this case

$$H_\rho = -\rho; \quad H_\eta = 0,$$

and the eigenfunction F is given by

$$F = \frac{\rho^{3/2}}{2}.$$

The insertion of the last eigenfunction in (2.126) gives the following metric

$$g = -\frac{2d\rho^2 + 2d\eta^2}{\rho^2} - \frac{\rho^2(1 + 3\rho) + \eta^2(9 + 7\rho)}{2\rho^5} d\phi^2 - \frac{8}{\rho^4} d\psi^2 - \frac{16\eta}{\rho^4} d\phi d\psi. \quad (2.129)$$

The inequality $F^2 < 4\rho^2(F_\rho^2 + F_\eta^2)$ holds for $\rho > 0$. Invoking the Taldersbank-Pedersen theorem, we see that for $\rho > 0$ the quaternionic kahler metric is -g. In this case $\kappa = -1$ and the three 1-forms A^i are

$$A^1 = \frac{2d\eta}{\rho}; \quad A^2 = \frac{2d\phi}{\rho}; \quad A^3 = \frac{2\eta}{\rho^2} d\phi + \frac{2}{\rho^2} d\psi.$$

The metric constructed in this example is defined for all the positive values of ρ .

B. With the function $G(x) = \log(x)x^3$ it is found the monopole

$$V = 3\eta\rho^2 - 2\eta^3$$

and the hyperkahler metric

$$ds^2 = (3\rho^2 - 6\eta^2)(d\rho^2 + d\eta^2 + \rho^2 d\phi^2) + \frac{1}{3\rho^2 - 6\eta^2} (d\psi + 6\eta\rho^2 d\phi)^2.$$

The Backlund transformed F results

$$F = \frac{3}{4}\rho^{3/2}(4\eta^2 - \rho^2)$$

and the corresponding metric is

$$g = g_{\rho\rho}(d\rho^2 + d\eta^2) + g_{\phi\phi}d\phi^2 + g_{\psi\psi}d\psi^2 + 2g_{\phi\psi}d\phi d\psi, \quad (2.130)$$

where the components of the metric tensor are

$$\begin{aligned} g_{\rho\rho} &= -\frac{4(8\eta^4 + 6\eta^2\rho^2 + 3\rho^4)}{(\rho^3 - 4\eta^2\rho)^2} \\ g_{\phi\phi} &= -\frac{8\eta^4\rho^2(19 + 5\rho) + 16\eta^6(9 + 7\rho) + \rho^6(1 + 35\rho) + 3\eta^2\rho^4(35 + 61\rho)}{9\rho^5(\rho^2 - 4\eta^2)^2(8\eta^4 + 6\eta^2\rho^2 + 3\rho^4)} \\ g_{\psi\psi} &= -\frac{64(4\eta^4 + \rho^4)}{9\rho^4(\rho^2 - 4\eta^2)^2(8\eta^4 + 6\eta^2\rho^2 + 3\rho^4)} \\ g_{\phi\psi} &= -\frac{32(8\eta^5 - \rho^2\eta^3 + 3\eta\rho^4)}{9\rho^4(\rho^2 - 4\eta^2)^2(8\eta^4 + 6\eta^2\rho^2 + 3\rho^4)}. \end{aligned}$$

As in the example A, $F^2 < 4\rho^2(F_\rho^2 + F_\eta^2)$ for $\rho > 0$, -g is a quaternionic kahler metric and $\kappa = -1$. The three forms A^i are given by

$$\begin{aligned} A^1 &= \frac{8\eta}{\rho^2 - 4\eta^2}d\rho + \frac{4(\rho^2 - 2\eta^2)}{\rho(\rho^2 - 4\eta^2)}d\eta \\ A^2 &= \frac{4}{3\rho(\rho^2 - 4\eta^2)}d\phi; \quad A^3 = \frac{4\eta}{3\rho(\rho^2 - 4\eta^2)}d\phi + \frac{4}{3\rho(\rho^2 - 4\eta^2)}d\psi. \end{aligned}$$

The metric (2.130) is singular at $\rho \rightarrow 0$ and at the lines $2|\rho| = |\eta|$.

C. The powers $G(x) = x^n$ and $G(x) = \log(x)x^{2n+1}$ can be integrated out giving polynomial solutions of higher degree. For instance $G(x) = \text{Log}(x)x^5$ gives

$$V = 3\eta\rho^2 - 2\eta^3; \quad F = \frac{1}{\eta}(8\eta^4 + 40\eta^2\rho^2 + 15\rho^4).$$

The even powers $G(x) = x^{2n} \log(x)$ can be integrated too, but the expressions of the metrics are more complicated by the appearance of logarithm terms. For example, with $G(x) = \log(x)x^2$ it is obtained

$$\begin{aligned} V &= 6\eta^2 - \rho^2 - 6\eta^2\sqrt{1 + \frac{\rho^2}{\eta^2}} + 2(\rho^2 - 2\eta^2)\log\left(\frac{2}{\eta + \sqrt{\eta^2 + \rho^2}}\right) \\ F &= \frac{4}{3}\eta^3 - 4\eta\rho^2 - \frac{4}{3}\eta^2(\eta^2 + \rho^2) + \frac{8}{3}\rho^2(\eta^2 + \rho^2) + 4\eta\rho^2\log\left(\frac{2}{\eta + \sqrt{\eta^2 + \rho^2}}\right). \end{aligned}$$

The expression for the quaternionic metric and the hyperkahler one corresponding to such solutions is very large and difficult to simplify.

D. The function $G(x) = e^x$ gives

$$V = e^{\pm i\eta}I_0(\rho) + c.c$$

where $I_n(\rho)$ denotes the modified Bessel function of the first kind, which are solutions of the equation

$$\rho^2 H''(\rho) + \rho H'(\rho) - (\rho^2 + n^2)H(\rho) = 0.$$

The hyperkahler space that corresponds to this monopole is:

$$ds^2 = \rho I_1(\rho)(d\rho^2 + d\eta^2 + \rho^2 d\phi^2) \cos(\eta) + \frac{1}{\rho I_1(\rho) \cos(\eta)} \{d\psi + \rho[I_1(\rho) + \frac{\rho}{2}(I_0(\rho) + I_2(\rho))]\sin(\eta)d\phi\}^2. \quad (2.131)$$

The Backlund transformed eigenfunction F is given by

$$F = \sqrt{\rho} I_1(\rho) e^{\pm i\eta} + c.c.$$

and from this solutions it follows the quaternionic metric

$$ds_{qk} = \frac{\Theta(\rho, \eta)}{4\rho I_1(\rho)^2} (d\rho^2 + d\eta^2) + \frac{[2\rho^{3/2} I_1(\rho) \beta \cos(\eta) - \rho^{3/2} (I_0(\rho) + I_2(\rho)) \alpha \sin(\eta)]^2}{\Phi(\rho, \eta)} + \frac{[2\rho^{3/2} I_1(\rho) \alpha \cos(\eta) + \sqrt{\rho} (\rho I_0(\rho) + 2I_1(\rho) + \rho I_2(\rho)) \beta \sin(\eta)]^2}{\Phi(\rho, \eta)}, \quad (2.132)$$

where it has been defined

$$\Theta(\rho, \eta) = \rho I_0(\rho)^2 + 2I_1(\rho) I_2(\rho) + \rho I_2(\rho)^2 + 4\rho I_1(\rho)^2 \left(\frac{\cos(\eta)}{\sin(\eta)}\right)^2 + 2I_0(\rho) (I_1(\rho) + \rho I_2(\rho))$$

and

$$\Phi(\rho, \eta) = -\rho^2 I_1(\rho)^2 \sin(\eta)^2 [4\rho^2 I_1(\rho)^2 \cos(\eta)^2 - I_1(\rho)^2 \sin(\eta)^2 + (\rho I_0(\rho) + I_1(\rho) + \rho I_2(\rho))^2 \sin(\eta)^2].$$

It will be quaternionic for the regions of the plane (ρ, η) in which $F^2 < 4\rho^2 (F_\rho^2 + F_\eta^2)$. In those regions $k = -1$. The three one forms A^i are

$$A^1 = \frac{1}{2} [1 + \tan(\eta) + \frac{\rho}{I_1(\rho)} \tan(\eta) (I_0(\rho) + I_2(\rho))] \frac{d\eta}{\rho} - \tan(\eta) d\rho$$

$$A^2 = \frac{\alpha}{\sqrt{\rho} I_1(\rho) \cos(\eta)}; \quad A^3 = \frac{\beta}{\sqrt{\rho} I_1(\rho) \cos(\eta)}.$$

The radial component of the metric (2.132) shows that some of the singularities are the zeros of $I_1(\rho)$.

By completeness we will also discuss the spaces corresponding to the m -pole solutions investigated in [155] and [132]. The dynamics of the M-theory on toric G_2 cones constructed with m -pole spaces as base manifolds have been analyzed in [97] and [98]. Such examples leads to toric Einstein self-dual spaces of positive scalar curvature which are complete (thus compact) and admitting only orbifold singularities. As such, they seem to be the best candidates with $U(1) \times U(1)$ isometry for which the physical analysis given in [42] can be applied. In the first reference of [132] it has been described the moduli space corresponding to the 3-pole solutions and has been shown that they encode some well known examples appearing in the physics, like the Bianchi type spaces. It will be shown that the hyperkahler metrics corresponding to the 3-pole solution are those discussed in [131] which gives rise to the 3-dimensional Eguchi-Hanson

like Einstein-Weyl metrics after the quotient by one of the isometries. The following exposition follows closely those given in the references [132].

The basic eigenfunctions F of (2.127) which we need to consider are

$$F(\rho, \eta, y) = \frac{\sqrt{(\rho)^2 + (\eta - y)^2}}{\sqrt{\rho}} \quad (2.133)$$

where the parameter y takes arbitrary real values. Using the Backlund transformation it is found the basic monopole

$$V(\eta, \rho, y) = -\log[\eta - y + \sqrt{\rho^2 + (\eta - y)^2}]. \quad (2.134)$$

Being the equations for F and V linear, for any set of real numbers w_i the functions

$$F = \sum_{j=0}^{k+1} w_j F(\rho, \eta, y_j). \quad (2.135)$$

$$V = \sum_{j=0}^{k+1} w_j V(\rho, \eta, y_j) \quad (2.136)$$

will be solutions too. For this reason the 2-pole functions given by

$$F_1 = \frac{1 + \sqrt{\rho^2 + \eta^2}}{\sqrt{\rho}}; \quad F_2 = \frac{\sqrt{(\rho)^2 + (\eta + 1)^2}}{\sqrt{\rho}} - \frac{\sqrt{(\rho)^2 + (\eta - 1)^2}}{\sqrt{\rho}},$$

are eigenfunctions of the hyperbolic laplacian. The first one gives rise to the spherical metric, while the second one gives rise to the hyperbolic metric

$$ds^2 = (1 - r_1^2 - r_2^2)^{-2} (dr_1^2 + dr_2^2 + r_1^2 d\theta_1^2 + r_2^2 d\theta_2^2).$$

The relation between the coordinates (r_1, r_2) and (ρ, η) can be extracted from the relation

$$(r_1 + ir_2)^2 = \frac{\eta - 1 + i\rho}{\eta + 1 + i\rho}.$$

The hyperkahler metrics corresponding to both cases are

$$ds^2 = -\frac{1}{\sqrt{\rho^2 + \eta^2}} (d\rho^2 + d\eta^2 + \rho^2 d\phi^2) - \sqrt{\rho^2 + \eta^2} (d\psi + \frac{\eta}{\sqrt{\rho^2 + \eta^2}} d\phi)^2, \quad (2.137)$$

and

$$ds^2 = \frac{\sqrt{\rho^2 + (\eta - 1)^2} - \sqrt{\rho^2 + (\eta + 1)^2}}{\sqrt{\rho^2 + (\eta + 1)^2} \sqrt{\rho^2 + (\eta - 1)^2}} (d\rho^2 + d\eta^2 + \rho^2 d\phi^2) + \frac{\sqrt{\rho^2 + (\eta + 1)^2} \sqrt{\rho^2 + (\eta - 1)^2}}{\sqrt{\rho^2 + (\eta - 1)^2} - \sqrt{\rho^2 + (\eta + 1)^2}} [d\psi + (\frac{\eta + 1}{\sqrt{\rho^2 + (\eta + 1)^2}} - \frac{\eta - 1}{\sqrt{\rho^2 + (\eta - 1)^2}}) d\phi]^2. \quad (2.138)$$

The general "3-pole" solutions are

$$F = \frac{1}{\sqrt{\rho}} + \frac{b + c/m}{2} \frac{\sqrt{\rho^2 + (\eta + m)^2}}{\sqrt{\rho}} + \frac{b - c/m}{2} \frac{\sqrt{\rho^2 + (\eta - m)^2}}{\sqrt{\rho}}.$$

By definition $-m^2 = \pm 1$, which means that m can be imaginary or real. The corresponding solutions are denominated type I and type II respectively. It is convenient to introduce the Eguchi-Hanson like coordinate system defined by

$$\rho = \sqrt{R^2 \pm 1} \cos(\theta), \quad \eta = R \sin(\theta),$$

where θ takes values in the interval $(-\pi/2, \pi/2)$. In this coordinates

$$\sqrt{\rho}F = 1 + bR + c \sin(\theta), \quad (2.139)$$

$$\rho^{-1}[\frac{1}{4}F^2 - \rho^2(F_\rho^2 + F_\eta^2)] = \frac{b(R \mp b) + c(\sin(\theta) + c)}{R^2 \pm \sin^2(\theta)}. \quad (2.140)$$

and the family of self-dual metrics corresponding to the 3-pole are expressed as

$$\begin{aligned} ds^2 = & \frac{b^2 - c^2 + (bR - cS)}{(1 + bR + cS)^2} \left(\frac{dR^2}{R^2 - 1} + \frac{dS^2}{1 - S^2} \right) \\ & \frac{1}{(1 + bR + cS)^2 (b^2 - c^2 + (bR - cS))(R^2 - S^2)} \\ & *((R^2 - 1)(1 - S^2)((bR - cS)d\varphi + (cR - bS)d\psi)^2 \\ & + ((b(R^2 - 1)S + c(1 - S^2)R)d\varphi + (c(R^2 - 1)S + b(1 - S^2)R + (R^2 - S^2)d\psi)^2) \end{aligned} \quad (2.141)$$

It has been denoted $S = \sin(\theta)$ here. The expression (2.141) includes some well known metrics. Let us focus in the type I case. The formulas (2.139) and (2.140) allows to determine the domain of definition of the metric (2.126). When b is nonzero for a given value of θ , $F = 0$ if $R = -(1 + c \sin(\theta))/b$ and $(\frac{1}{4}F^2 - \rho^2(F_\rho^2 + F_\eta^2)) = 0$ if $R = (b^2 + c^2 + c \sin(\theta))/b$. The case $c = 0$ correspond to a bi-axial Bianchi IX metric [135]. The domains of definition are $(-\infty, R_\infty)$, $(R_\infty, R_\pm)$, and $(R_\pm, \infty)$. In the first two cases the curvature is negative, and in the last one positive, and in the two last cases there is an unremovable singularity at $R = R_\pm$. In the case $b = 0$ for $c > 1$ and $c < 1$ the metric will be of Bianchi VIII type [133]. The case $c = 1$ corresponds to the Bergmann metric on CH^2 .

For the type II case, the range of R is $(1, \infty)$ but the moduli space is more complex than in the type I case. For the lines $b = \pm c$ it is obtained the hyperbolic metric if $b < 0$ and the spherical metric if $b > 0$. If $(b, c) = (1, 0)$ it is obtained the Fubini-Study metric on CP^2 whereas the points $(0, 1)$, $(-1, 0)$ and $(0, -1)$ yield again the Bergmann metric on CH^2 . Along the lines joining $(1, 0)$ with others we have bi-axial Bianchi metric IX, while along the lines between $(0, 1)$, $(-1, 0)$ and $(0, -1)$ the metric is Bianchi VIII. A more complete description is given in [132].

The triplet of one forms corresponding to this family of metrics is

$$\begin{aligned} A^1 &= A_+^1 + A_-^1, \\ A^2 &= \frac{\sqrt{(R^2 \pm 1)(1 - S^2)}}{(1 + bR + cS)} d\phi \quad A^3 = \frac{d\psi + \eta d\phi}{(1 + bR + cS)}, \end{aligned}$$

where it has been defined

$$A_\pm^1 = A_{1\pm}^1 + A_{2\pm}^1 + A_{3\pm}^1,$$

with

$$A_{1\pm}^1 = \frac{(b \pm c/m)(SR \pm m)}{(1 + bR + cS)\sqrt{(1 - S^2)(R^2 \pm 1) + (SR \pm m)^2}} * \\ \left(\frac{S\sqrt{R^2 \pm 1}}{2\sqrt{1 - S^2}} dS + \frac{R\sqrt{1 - S^2}}{\sqrt{R^2 \pm 1}} dR \right)$$

and

$$A_{2\pm}^1 = \frac{(b \pm c/m)\sqrt{(1 - S^2)(R^2 \pm 1) + (SR \pm m)^2}}{2(1 + bR + cS)\sqrt{R^2 \pm 1}} \left(R \frac{dS}{\sqrt{1 - S^2}} + S dR \right)$$

and

$$A_{3\pm}^1 = \left[\frac{(b \pm c/m)(R^2 \pm 1)^{1/2}(1 - S^2)^{1/2}}{2\sqrt{(1 - S^2)(R^2 \pm 1) + (SR \pm m)^2}} - \frac{(b \pm c/m)\sqrt{(1 - S^2)(R^2 \pm 1) + (SR \pm m)^2}}{4(1 - S^2)^{1/2}(R^2 \pm 1)^{1/2}} \right] * \\ \frac{\sqrt{(1 - S^2)(R^2 \pm 1)}}{(1 + bR + cS)\sqrt{R^2 \pm 1}} \left(R \frac{dS}{\sqrt{1 - S^2}} + S dR \right)$$

(The sign $\pm$ in $(R^2 \pm 1)$ depends only on the metric in consideration, it is $+$ for type I and $-$ for type II.)

The Backlund transformed function V reads

$$V = \log(\rho) + \frac{(b + c/m)}{2} \log \left[\frac{\eta - m + \sqrt{(\eta - m)^2 + \rho^2}}{\rho} \right] \\ + \frac{(b - c/m)}{2} \log \left[\frac{\eta + m + \sqrt{(\eta + m)^2 + \rho^2}}{\rho} \right].$$

For the type I case this is the potential for an axially symmetric circle of charge, while the type II case corresponds to two point sources on the axis of symmetry. The hyperkahler metrics obtained are encoded in the following expression

$$ds^2 = \frac{bR + c\sqrt{1 - S^2}}{R^2 \pm (1 - S^2)} (d\rho^2 + d\eta^2 + \rho^2 d\phi^2) \\ + \frac{R^2 \pm (1 - S^2)}{bR + c\sqrt{1 - S^2}} \left[d\psi + \frac{R^2 \pm (1 - S^2) - b(R^2 \pm 1)\sqrt{1 - S^2} + cRS^2}{R^2 \pm (1 - S^2)} d\phi \right]^2. \quad (2.142)$$

This manifolds have been investigated recently in [131] and it has been shown that the quotient of (2.142) with $\frac{\partial}{\partial \phi}$ gives the Eguchi-Hanson type Einstein-Weyl metrics in D=3.

The continuum limit of the expressions (2.135) and (2.136) are

$$F(\rho, \eta) = \int w(y) F(\rho, \eta, y) dy. \quad (2.143)$$

$$V(\rho, \eta) = \int w(y) V(\rho, \eta, y) dy. \quad (2.144)$$

where $w(y)$ is a distribution with compact support in R . A choice of $w(y)$ for which at least one of the integrals (2.143) and (2.144) converges gives rise to an smooth solution. For instance for $w(y) = y/(y^2 + 1)^2$ it is obtained the following non-trivial monopole

$$V(\rho, \eta) = \frac{\cos(\frac{1}{2} \text{Arg}(1 - 2i\eta - \eta^2 - \rho^2)) \log(\frac{|1-i\eta-\sqrt{(1-i\eta)^2+\rho^2}|}{|1+i\eta-\sqrt{(1-i\eta)^2+\rho^2}|})}{\sqrt{|(1-i\eta)^2 + \rho^2|}} \\ + \frac{\sin(\frac{1}{2} \text{Arg}(1 - 2i\eta - \eta^2 - \rho^2)) \text{Arg}(\frac{1-i\eta-\sqrt{(1-i\eta)^2+\rho^2}}{1+i\eta-\sqrt{(1-i\eta)^2+\rho^2}})}{\sqrt{|(1-i\eta)^2 + \rho^2|}},$$

and from (2.83) follows an hyperkahler metric. But (2.143) is divergent for this distribution.

To conclude this subsection it should be mentioned that higher m -pole solutions have been considered in [97] and [98], and that quaternionic spaces with torus symmetry have been investigated recently in [136] using the harmonic space formalism.

2.5 Higher dimensional hypergeometry

The previous sections were concerned with quaternion Kahler and hyperkahler manifolds in $d = 4$. In this section we deal with higher dimensional ones. We present the most general $4n$ -dimensional hyperkahler metrics with n commuting tri-holomorphic Killing vectors [146], namely the Pedersen-Poon metrics. This result generalize the Gibbons-Hawking one (2.66) to $d = 4n$. We also present a construction allowing to extend an arbitrary quaternion Kahler metric in $d = 4n$ to other quaternion Kahler and an hyperkahler ones in $d = 4(n + 1)$. This is the Swann construction [22]. As we will see, the Swann extension of the Calderbank-Pedersen metrics (2.126) preserves the Killing commuting Killing vectors and they are tri-holomorphic with respect to the new metric. Therefore the extension is of a Pedersen-Poon type. Once this is understood it is straightforward to construct certain classical supergravity solutions. This is the subject of the next section.

2.5.1 Construction of $4n$ hyperkahler manifolds with T^n tri-holomorphic isometry

Let us remind that in section 3 we have described all the four dimensional hyperkahler spaces with at least one self-dual Killing vector namely, the Gibbons-Hawking metrics. We have seen that for such spaces there always exist a coordinate system for which the metric take the form

$$g = V^{-1}(dt - \omega)^2 + V dx \cdot dx. \quad (2.145)$$

The vector $K = \partial_t$ satisfies

$$\mathcal{L}_K g = 0 \quad \implies \quad K_{\mu;\nu} + K_{\nu;\mu} = 0 \quad (2.146)$$

and is therefore a Killing vector. The tensor $K_{\mu\nu} = \nabla_\mu K_\nu$ satisfies

$$*_g K_{\mu\nu} = K_{\mu\nu} \quad (2.147)$$

and for this reason K is called self-dual in the terminology of the references [75]-[74].

The vector $K = \partial_t$ also preserve the hyperkahler triplet $\overline{\mathcal{J}}^k$ of (2.145). Explicitly $\overline{\mathcal{J}}^k$ is given by

$$\overline{\mathcal{J}}^k = (dt + A) \wedge dx^k - \epsilon_{rs}^k U(dx^r \wedge dx^s), \quad (2.148)$$

and from (2.148) its is direct to check that

$$\mathcal{L}_K \overline{\mathcal{J}}^i = 0. \quad (2.149)$$

In general a vector preserving $\overline{\mathcal{J}}^i$ as in (2.150) is called tri-holomorphic. If instead we have that

$$\mathcal{L}_K J^i = 0, \quad (2.150)$$

then K preserve the complex structures and is called tri-hamiltonian.

It can be shown that any two of the conditions given above for K implies the third. This mean in particular that for the metric (2.145) the vector $K = \partial_t$ is Killing, tri-holomorphic and tri-hamiltonian. For the spaces (2.79) this condition does not hold, the Gibbons-Hawking metrics (2.145) are the only one with this properties in $d = 4$.

The higher dimensional analogs of (2.145) to $d = 4n$ dimensions are the hyperkahler spaces with n commuting tri-holomorphic Killing vectors ∂_{t_i} . A plausible anatz is obtained by introducing a set on $3n$ coordinates $x^i = \{x_r^i, \ i = 1, \dots, n \ r = 1, 2, 3\}$ and the metric

$$\overline{g} = U_{ij} dx_i \cdot dx_j + U^{ij} (dt_i + A_i)(dt_j + A_j), \quad (2.151)$$

where the symmetric matrix U_{ij} and then its inverse U^{ij} are independent of the coordinates t^i . The n one forms A_j have the form

$$A_i = dx^j \cdot \Sigma_{ij} \quad (2.152)$$

where the matrix Σ_{ij} also do not depend on the t_i coordinates. The hyperkahler triplet corresponding to (2.151) is given by

$$\overline{\mathcal{J}}^k = (dt_r + A_r) \wedge dx_k^r - U_{ij}(dx^i \times dx^j)_k \quad (2.153)$$

Here $\times$ denote the exterior product of forms (for instance the 3-component of $dx^i \times dx^j$ is $dx_1^i \times dx_2^j$). Then it is directly seen that $\mathcal{L}_{\partial_i} \overline{\mathcal{J}}^k = 0$, therefore the Killing vectors ∂_{t_i} are tri-holomorphic and tri-hamiltonian. For $n = 1$ (2.151) reduce to the Gibbons-Hawking metrics (2.145).

Let us now find the conditions should be satisfied for (2.151) to be an hyperkahler metric. In order to use the $4n$ -bein formalism we can express the matrix U as

$$U = K^T K \quad (2.154)$$

for a non singular matrix K , which is defined by (2.154) up to an $SO(n)$ rotation. Then we can select the $4n$ -bein as

$$\begin{aligned} e_a &= K_{aj} dx^j \quad (i = 1, \dots, 3n) \\ e_a &= (dt_j + A_j) K^{aj} \end{aligned} \quad (2.155)$$

being K^{aj} the inverse of K_{aj} . Then if (2.151) is hyperkahler, then $N^i(X, Y) = 0$ for every of the complex structures J^i corresponding to (3.300). In $4n$ dimensions the conditions (2.99) are generalized to a system of equations with a $4n$ vector potential $\tilde{A}_1, \dots, \tilde{A}_{4n}$ that we will not write for simplicity. But introducing (2.155) into such system gives finally the following system of equations

$$\begin{aligned} F_{x_k^i x_l^j} &= \epsilon_{klm} \nabla_{x_m^i} U_j, \\ \nabla_{x_m^i} U_j &= \nabla_{x_m^j} U_i, \\ U_i &= (U_{i1}, \dots, U_{in}), \end{aligned} \quad (2.156)$$

being $F_{x_k^i x_l^j}$ the strength tensor corresponding to the "vector potential" A_a . If the last two conditions holds then (2.151) is hyperkahler.

It was also shown that (2.151) are unique. In other words, for any T^n hyperkahler metrics there exists a coordinate system in which they can be cast in the form (2.151) [146]-[171]. Such system is the momentum map system. In general momentum maps are related to a compact Lie group G acting over an hyperkahler manifold M by tri-holomorphic Killing vectors X , i.e, Killing vectors satisfying

$$\mathcal{L}_X J^k = 0.$$

The last condition implies that X preserves the hyperkahler forms $\overline{\mathcal{J}}^k$, that is

$$\mathcal{L}_X \overline{\mathcal{J}}^k = 0 = i_X d\overline{\mathcal{J}}^k + d(i_X \overline{\mathcal{J}}^k).$$

Here $i_X \overline{\mathcal{J}}^k$ denotes the contraction of X with the hyperkahler forms. By supposition M is hyperkahler, then $d\overline{\mathcal{J}}^k = 0$ and

$$d(i_X \overline{\mathcal{J}}^k) = 0.$$

This mean that $i_X \overline{\mathcal{J}}^k$ can be expressed locally as the differential of certain function x_k^X , namely

$$dx_k^X = i_X \overline{\mathcal{J}}^k. \quad (2.157)$$

The functions x_k^X are called *momentum maps* and are defined up to a constant.

In the case (2.151) the isometries are $\partial/\partial t_i$ and the hyperkahler form corresponding to (2.160) is [138]

$$\overline{\mathcal{J}}^k = (dt_i + A_i)dx_k^i - U_{ij}(dx^i \wedge dx^j)_k. \quad (2.158)$$

From the last expression it follows that

$$dx_k^i = i_{t_i} \overline{\mathcal{J}}^k \quad (2.159)$$

and therefore $(x_i^1, \dots, x_i^n)$ are really the momentum maps of the isometries. All the results of this section are stated in the following proposition [146].

Proposition 9 *For any hyperkahler metric in $D = 4n$ with n commuting tri-holomorphic $U(1)$ isometries there exists a coordinate system in which takes the form*

$$\overline{g} = U_{ij}dx^i \cdot dx^j + U^{ij}(dt_i + A_i)(dt_j + A_j), \quad (2.160)$$

where (U_{ij}, A_i) are solutions of the generalized monopole equation

$$\begin{aligned} F_{x_\mu^i x_\nu^j} &= \epsilon_{\mu\nu\lambda} \nabla_{x_\lambda^i} U_j, \\ \nabla_{x_\lambda^i} U_j &= \nabla_{x_\lambda^j} U_i, \\ U_i &= (U_{i1}, \dots, U_{in}), \end{aligned} \quad (2.161)$$

and the coordinates $(x_i^1, \dots, x_i^n)$ with $i = 1, 2, 3$ are the momentum maps of the tri-holomorphic vector fields $\partial/\partial t_i$.

The spaces in the form (2.160) are known as the Pedersen-Poon metrics. Equations (2.161) are known as monopole equations by interpreting $F_{x_\mu^i x_\nu^j}$ as a vector potential and U_i as certain Higgs fields. Some Pedersen-Poon examples will be constructed in the following section.

2.5.2 Quaternion Kahler spaces in quaternion notation

It is convenient to introduce a notation for quaternion Kahler manifolds in terms of quaternions, that has the advantage to be more compact [22]. It is fundamental to observe that the holonomy of a quaternion Kahler space is in $Sp(n) \times Sp(1)$ and therefore from the very beginning the connection ω will take values in the algebra $sp(n) \oplus sp(1)$ with splitting $\omega = \omega_+ + \omega_-$. The Lie algebra $sp(n)$ of $Sp(n)$ can be expressed in terms of quaternion valued matrices A with the property $A + \overline{A}^t = 0$.

Let us consider now two arbitrary $\mathbf{H}$ -valued 1-forms

$$\mu = \mu_0 + \mu_1 I + \mu_2 J + \mu_3 K, \quad \nu = \nu_0 + \nu_1 I + \nu_2 J + \nu_3 K,$$

being I, J and K unit quaternions. We define the quaternionic wedge product as

$$\mu \wedge_q \nu = (\mu_0 \wedge \nu_1 - \mu_2 \wedge \nu_3)I + (\mu_0 \wedge \nu_2 - \mu_3 \wedge \nu_1)J + (\mu_0 \wedge \nu_3 - \mu_1 \wedge \nu_2)K \quad (2.162)$$

$$+\mu_0 \wedge \nu_0 + \mu_1 \wedge \nu_1 + \mu_2 \wedge \nu_2 + \mu_3 \wedge \nu_3,$$

and in particular for two pure quaternionic forms we have

$$\bar{\mu} \wedge_q \mu = (\mu_0 \wedge \mu_1 - \mu_2 \wedge \mu_3)I + (\mu_0 \wedge \mu_2 - \mu_3 \wedge \mu_1)J + (\mu_0 \wedge \mu_3 - \mu_1 \wedge \mu_2)K, \quad (2.163)$$

with components only in $\text{Im } \mathbf{H}$. This product can be extended to $\mathbf{H}^n$ valued 1-forms, with $\mathbf{H}$ -components given by

$$\mu^i = \mu_0^i I + \mu_1^i J + \mu_2^i K, \quad \nu^i = \nu_0^i I + \nu_1^i J + \nu_2^i K. \quad (2.164)$$

The wedge product $\bar{\wedge}$ of the forms (2.164) is defined as

$$\mu \bar{\wedge} \nu = \mu^i \wedge_q \nu^i,$$

and is clearly seen that $\mu \bar{\wedge} \nu$ is $\mathbf{H}$ -valued.

Consider now a four dimensional metric

$$g = e^1 \otimes e^1 + e^2 \otimes e^2 + e^3 \otimes e^3 + e^4 \otimes e^4.$$

with an orthonormal tetrad e^i . We can extend such tetrad to a quaternionic valued one

$$e = e_1 + e_2 I + e_3 J + e_4 K, \quad \bar{e} = e_1 - e_2 I - e_3 J - e_4 K,$$

for which the distance g is expressed as $g = e \otimes \bar{e}$. The hyperkahler triplet associated to g

$$\begin{aligned} \bar{J}^1 &= e_4 \otimes e^1 - e_1 \otimes e^4 + e_2 \otimes e^3 - e_3 \otimes e^2 \\ \bar{J}^2 &= e_4 \otimes e^2 - e_2 \otimes e^4 + e_3 \otimes e^1 - e_1 \otimes e^3 \\ \bar{J}^3 &= e_4 \otimes e^3 - e_3 \otimes e^4 + e_1 \otimes e^2 - e_2 \otimes e^1 \end{aligned} \quad (2.165)$$

is extended in this notation to a quaternion valued $(1,1)$ tensor

$$\bar{J} = \bar{J}^1 I + \bar{J}^2 J + \bar{J}^3 K. \quad (2.166)$$

From the definition (2.163) it follows that

$$\bar{J} = e \wedge_q \bar{e}. \quad (2.167)$$

In $4n$ dimensions we select the $\mathbf{H}^n$ -valued einbein e defined by

$$e^a = e_0^a + e_1^a I + e_2^a J + e_3^a K. \quad (2.168)$$

The metric is then $g = e^a \otimes \bar{e}^a$. Taking into account the representation of $Sp(n)$ in terms of quaternions we can write $\omega = \omega_+ + \omega_-$ also as a $\mathbf{H}^n$ one-form. Then it can be shown that the first Cartan equation translates into

$$de = -\omega \bar{\wedge} e \quad (2.169)$$

and that the hyperkahler triplet is

$$\bar{J} = e \bar{\wedge} \bar{e}. \quad (2.170)$$

The quaternionic expression of the fundamental relations (2.9) and (2.7) in $D=4n$ are

$$d\omega_- - \omega_- \bar{\wedge} \omega_- = \Lambda \bar{J}, \quad d\bar{J} - \omega_- \bar{\wedge} \bar{J} + \bar{J} \bar{\wedge} \omega_- = 0. \quad (2.171)$$

Introducing the second (2.171) into the first to give

$$\Lambda d\bar{J} + \omega_- \bar{\wedge} d\omega_- - d\omega_- \bar{\wedge} \omega_- = 0, \quad d\omega_- - \omega_- \bar{\wedge} \omega_- + \Lambda \bar{J} = 0, \quad (2.172)$$

expressing $\bar{J}$ and $d\bar{J}$ entirely in terms of the $sp(1)$ connection ω_- and its differentials.

2.5.3 The Swann extension

The Swann construction extend any quaternionic Kahler in dimension $D = 4n$ to another $4(n+1)$ -dimensional quaternion Kahler one [22]. Before to present it, let us consider an $4n$ dimensional hyperkahler manifold M with metric $\bar{g} = \delta_{ab}e^a \otimes e^b$. From the definition $\bar{g}$ has holonomy in $Sp(n)$. We make the trivial extension to a $4(n+1)$ -dimensional metric given by

$$g = \bar{g} + (du_0)^2 + (du_1)^2 + (du_2)^2 + (du_3)^2, \quad (2.173)$$

being (u_0, u_1, u_2, u_3) four new coordinates. The metric (2.173) is the direct sum of $\bar{g}$ plus a flat metric. Then the holonomy Γ of (2.173) lies in $Sp(n) \subset Sp(n+1)$, and thus (2.173) is a trivial extension of $\bar{g}$ to a higher dimensional hyperkahler one.

Now let us generalize the trivial ansatz (2.173) by extending a quaternionic Kahler metric $\bar{g} = \delta_{ab}e^a \otimes e^b$ to another one in $D = 4(n+1)$ of the form

$$g_s = g|u|^2\bar{g} + f|du + u\omega_-|^2. \quad (2.174)$$

We have introduced the quaternions

$$u = u_0 + u_1I + u_2J + u_3K, \quad \bar{u} = u_0 - u_1I - u_2J - u_3K$$

$$\omega_- = \omega_-^1I + \omega_-^2J + \omega_-^3K$$

and the radius

$$|u|^2 = u\bar{u} = (u_0)^2 + (u_1)^2 + (u_2)^2 + (u_3)^2.$$

being I, J, K the unit quaternions. It is clear that if we consider an hyperkahler base $\bar{g}$ then $\omega_-^i = 0$ and we recover (2.173) from (2.174) if the functions $f(|u|^2)$ and $g(|u|^2)$ are simply constants. In general the explicit form of the metric is

$$g_s = g|u|^2\bar{g} + f[(du_0 - u_i\omega_-^i)^2 + (du_i + u_0\omega_-^i + \epsilon_{ijk}u_k\omega_-^k)^2] \quad (2.175)$$

although the expression (2.174) will be better for the following.

If g_s is quaternion Kahler space then Θ is closed (see the discussion below (2.19)) and this gives a system of equation defining f and g . The fundamental four form (2.19) for g_s is explicitly

$$\begin{aligned} \Theta &= fg(\alpha \wedge \bar{\alpha} \wedge u\bar{e}^t \wedge e\bar{u} + u\bar{e}^t \wedge e\bar{u} \wedge \alpha \wedge \bar{\alpha}) \\ &\quad + g^2|u|^4\bar{e}^t \wedge e \wedge \bar{e}^t \wedge e + f^2\alpha \wedge \bar{\alpha} \wedge \alpha \wedge \bar{\alpha} \end{aligned} \quad (2.176)$$

being $\alpha = du + u\omega_-$. If we want to impose $d\Theta = 0$ to (2.174) we need the identities

$$d(\bar{e}^t \wedge e \wedge \bar{e}^t \wedge e) = 0, \quad (2.177)$$

$$dr^2 = u d\bar{u} + \bar{u} du = u\bar{\alpha} + \alpha\bar{u}. \quad (2.178)$$

$$d(u\bar{e}^t \wedge e\bar{u}) = \alpha \wedge \bar{e}^t \wedge e\bar{u} + u\bar{e}^t \wedge e \wedge \bar{\alpha}, \quad (2.179)$$

$$d(\alpha \wedge \bar{\alpha}) = -c(u\bar{e}^t \wedge e \wedge \bar{\alpha} + \alpha \wedge \bar{e}^t \wedge e\bar{u}). \quad (2.180)$$

An straightforward calculation by using (2.177)-(2.180) gives

$$d\Theta = A\alpha \wedge \bar{\alpha} \wedge u\bar{e}^t \wedge e\bar{u} + Au\bar{e}^t \wedge e\bar{u} \wedge \alpha \wedge \bar{\alpha} + B|u|^4 dr^2 \wedge \bar{e}^t \wedge e \wedge \bar{e}^t \wedge e, \quad (2.181)$$

with the coefficients A and B given explicitly by

$$A = -\frac{3c}{|u|^2}f^2 + \frac{3}{|u|^2}fg + (fg)_{|u|},$$

$$B = -\frac{2c}{|u|^2}fg + \frac{2}{|u|^2}g^2 + 2g(g)_{|u|}.$$

The closure of Θ implies $A = B = 0$ and this gives two differential equations for f and g with solution

$$f = \frac{a}{c(b|u|^2 + d)^2}, \quad (2.182)$$

$$g = \frac{1}{b|u|^2 + d}, \quad (2.183)$$

with a , b , c and d constants. Introducing (2.182)-(2.183) into (2.175) gives the metric

$$g_s = \frac{|u|^2}{b|u|^2 + d}\bar{g} + \frac{a}{c(b|u|^2 + d)^2}[(du_0 - u_i\omega_-^i)^2 + (du_i + u_0\omega_-^i + \epsilon_{ijk}u_k\omega_-^k)^2]. \quad (2.184)$$

The closure of Θ in principle is not enough to prove that (2.174) is quaternionic Kahler. But it is possible to show that the other necessary and sufficient conditions (2.8) and (2.9) are indeed satisfied for (2.184). Therefore (2.184) gives a family of quaternion Kahler metrics, known as the Swann metrics. The case $b = 0$ corresponds to an hyperkahler metric.

2.6 Spaces with G_2 holonomy

The Lie group G_2 was considered by Berger [1] as one of the possible holonomy groups of a Riemannian Ricci-flat space. He also proved that G_2 holonomy is possible only in seven dimensions. The existence of G_2 holonomy metrics was proved rigorously in [27] 30 years after the Berger work. In particular explicit non compact examples in [29].

These spaces have direct application to Kaluza-Klein compactifications of M-theory. This is due to the fact that G_2 holonomy manifolds have globally defined one Killing spinor η satisfying $D_i\eta = 0$ and therefore as internal spaces they preserve certain amount of supersymmetries after compactification. In fact the presence of such spinor is the reason for the reduction of the holonomy from $SO(7)$ to G_2 .

One of the main requirements in order to obtain a realistic four dimensional theory after compactification is that the internal space should be compact [26]. Also the appearance of chiral fermions in the effective theory is possible only if the internal space has certain singularities [42]. Although the existence of compact G_2 holonomy spaces was rigorously proved recently by Joyce in [32], no explicit metrics over them are known. Although this problem complicates the analysis of the effective theory in $d = 4$ all the physical phenomena near of the singularity is local, that is, independent on the global properties of the manifold [95]. Non compact metrics possessing conical singularities are therefore suitable for analyzing this kind of phenomena.

There also exist seven dimensional spaces that admit conformal Killing spinors that is, satisfying $D_i\eta \sim \eta$. These spaces are called weak G_2 holonomy spaces although their holonomy is generically in $SO(7)$. Weak G_2 holonomy metrics also preserve certain amount of supersymmetries and there are known compact conical examples [43]. In general weak G_2 holonomy spaces are internal spaces with the presence of fluxes and can be constructed as the conical part of cohomogeneity one $Spin(7)$ spaces.

In this section we present certain features about G_2 , weak G_2 and $Spin(7)$ holonomy metrics. In particular we will show that they are defined by certain self-duality conditions analog to the four dimensional case but with the octonion constants replacing the Kronecker symbols. The relation with octonion algebra is explained in certain detail.

2.6.1 The group G_2 and the octonions

In the following we make a brief description about the Lie group G_2 . It is known that $G_2 \subset SO(7)$ is one of the exceptional simple Lie groups and that is simply connected. Its Lie algebra has 14 generators G^{ab} and is the subgroup of $SO(7)$ that is isomorphic to the automorphism group of the octonions O [125].

The octonions constitute the only non associative division algebra and can be constructed as the doubling of the quaternions H (the associative ones are R , C and H). An arbitrary octonion $x \in O$ can be written as $x = x^0 + x^i e_i$, where the set e_i is a basis of 7 unit octonions with the multiplication rule

$$e_i \cdot e_j = c_{ijk} e_k; \quad e_i \cdot 1 = 1 \cdot e_i = e_i \quad (2.185)$$

and the components x^i are real numbers. The constants c_{ijk} that define the multiplication table (2.185) are totally antisymmetric and defined by

$$c_{123} = c_{246} = c_{435} = c_{367} = c_{651} = c_{572} = c_{714} = 1, \quad (2.186)$$

up to an index permutation. The constants corresponding to another set of indices are zero. The conjugated octonion basis is $\bar{e}_i = -e_i$ and it is clear that $\bar{x} = x^0 - x^i e_i$.

It is extremely important to recall that the octonions itself are not an associative algebra. From (2.185) and (2.192) it is seen that $(e^3 e^7) e^5 - e^3 (e^7 e^5) = -e^1$, which clearly shows non associativity. For this reason the octonions itself cannot be represented as a matrices with complex components and do not satisfy the Jacobi identities. In other words, the algebra of O is not a Lie algebra.

An automorphism is an isomorphism of an algebra A into itself. Under an automorphism transformation the multiplication table

$$x.y = z$$

is left invariant, in other words

$$T(x).T(y) = T(z) \quad \text{where } T \in \text{Aut} A$$

The following characterization of the group G_2 as the group of automorphism of the octonion algebra O is due to Zorn [169]. Consider three arbitrary elements of O , for instance, e_1 , e_2 and e_4 , and the transformation σ given by

$$\begin{aligned} \sigma(e_1) &= e_1 \\ \sigma(e_2) &= \cos(\alpha)e_2 + \sin(\alpha)e_3 \\ \sigma(e_4) &= \cos(\beta)e_4 + \sin(\beta)e_7 \end{aligned} \tag{2.187}$$

The images of the other elements are defined by the relations

$$\begin{aligned} \sigma(e_2).\sigma(e_4) &= \sigma(e_6), & \sigma(e_1).\sigma(e_2) &= \sigma(e_3) \\ \sigma(e_1).\sigma(e_4) &= \sigma(e_7), & \sigma(e_4).\sigma(e_3) &= \sigma(e_5) \end{aligned} \tag{2.188}$$

Explicitly we obtain that

$$\begin{aligned} \sigma(e_1) &= e_1 \\ \begin{pmatrix} \sigma(e_2) \\ \sigma(e_3) \end{pmatrix} &= \begin{pmatrix} \cos(\alpha) & \sin(\alpha) \\ -\sin(\alpha) & \cos(\alpha) \end{pmatrix} \begin{pmatrix} e_2 \\ e_3 \end{pmatrix} \\ \begin{pmatrix} \sigma(e_4) \\ \sigma(e_7) \end{pmatrix} &= \begin{pmatrix} \cos(\beta) & \sin(\beta) \\ -\sin(\beta) & \cos(\beta) \end{pmatrix} \begin{pmatrix} e_4 \\ e_7 \end{pmatrix} \\ \begin{pmatrix} \sigma(e_6) \\ \sigma(e_5) \end{pmatrix} &= \begin{pmatrix} \cos(\alpha + \beta) & \sin(\alpha + \beta) \\ -\sin(\alpha + \beta) & \cos(\alpha + \beta) \end{pmatrix} \begin{pmatrix} e_6 \\ e_5 \end{pmatrix} \end{aligned} \tag{2.189}$$

Relation (2.189) can also be written as

$$\begin{pmatrix} \sigma(e_2) + i\sigma(e_3) \\ \sigma(e_4) + i\sigma(e_7) \\ \sigma(e_6) + i\sigma(e_5) \end{pmatrix} = e^{(a\lambda_3 + b\lambda_8)e_1} \begin{pmatrix} e_2 + ie_3 \\ e_4 + ie_7 \\ e_6 + ie_5 \end{pmatrix} \tag{2.190}$$

where λ_3 and λ_8 are the Gell-Mann traceless matrices

$$\lambda_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \lambda_8 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix} \tag{2.191}$$

and a and b are related to the angles by

$$\alpha = a + \frac{b}{\sqrt{3}}, \quad \beta = -a + \frac{b}{\sqrt{3}}. \quad (2.192)$$

Relations (2.188) and (1.437) implies that $\sigma(e_i)$ satisfy the same multiplication rule than e_i . Conversely, every automorphism of the octonions belong in this manner to at least one Cayley basis (that is, a basis generated by three different elements). The automorphism of the form given above are called canonical automorphism. Each canonical automorphism has a fixed element and three invariant planes

$$\begin{aligned} \Psi(e_1) &= \frac{1}{2} \begin{pmatrix} e_2 + ie_3 \\ e_4 + ie_7 \\ e_6 + ie_5 \end{pmatrix} = \frac{(1 + ie_1)}{2} \begin{pmatrix} e_6 \\ e_2 \\ e_4 \end{pmatrix} \\ \Psi(e_2) &= \frac{1}{2} \begin{pmatrix} e_4 + ie_6 \\ e_3 + ie_1 \\ e_5 + ie_7 \end{pmatrix} = \frac{(1 + ie_2)}{2} \begin{pmatrix} e_4 \\ e_3 \\ e_5 \end{pmatrix} \\ \Psi(e_3) &= \frac{1}{2} \begin{pmatrix} e_5 + ie_4 \\ e_1 + ie_2 \\ e_6 + ie_7 \end{pmatrix} = \frac{(1 + ie_3)}{2} \begin{pmatrix} e_5 \\ e_1 \\ e_6 \end{pmatrix} \\ \Psi(e_4) &= \frac{1}{2} \begin{pmatrix} e_3 + ie_5 \\ e_6 + ie_2 \\ e_7 + ie_1 \end{pmatrix} = \frac{(1 + ie_4)}{2} \begin{pmatrix} e_3 \\ e_6 \\ e_7 \end{pmatrix} \\ \Psi(e_5) &= \frac{1}{2} \begin{pmatrix} e_1 + ie_6 \\ e_4 + ie_3 \\ e_7 + ie_2 \end{pmatrix} = \frac{(1 + ie_5)}{2} \begin{pmatrix} e_1 \\ e_4 \\ e_7 \end{pmatrix} \\ \Psi(e_6) &= \frac{1}{2} \begin{pmatrix} e_2 + ie_4 \\ e_5 + ie_1 \\ e_7 + ie_3 \end{pmatrix} = \frac{(1 + ie_6)}{2} \begin{pmatrix} e_2 \\ e_5 \\ e_7 \end{pmatrix} \\ \Psi(e_7) &= \frac{1}{2} \begin{pmatrix} e_1 + ie_4 \\ e_2 + ie_5 \\ e_3 + ie_6 \end{pmatrix} = \frac{(1 + ie_7)}{2} \begin{pmatrix} e_1 \\ e_2 \\ e_3 \end{pmatrix} \end{aligned} \quad (2.193)$$

A canonical automorphism leaving invariant the element e_A is defined by

$$\Psi'(e_A) = e^{(a_A \lambda_3 + b_A \lambda_8) e_A} \begin{pmatrix} e_B + ie_C \\ e_D + ie_E \\ e_F + ie_g \end{pmatrix} = \begin{pmatrix} e'_B + ie'_C \\ e'_D + ie'_E \\ e'_F + ie'_g \end{pmatrix} \quad (2.194)$$

(no Einstein convention is understood). Because we have seven canonical transformations the group of automorphism of O is 14 dimensional.

Let us find the Lie algebra of the automorphisms. We will take the parameters a_A and b_A . Corresponding to a_A we have

$$\Psi'(e_A) = e^{a_A \lambda_3 e_A} \begin{pmatrix} e_B + ie_C \\ e_D + ie_E \\ e_F + ie_g \end{pmatrix} = \begin{pmatrix} e'_B + ie'_C \\ e'_D + ie'_E \\ e'_F + ie'_g \end{pmatrix}$$

which gives

$$\begin{pmatrix} e'_B \\ e'_C \end{pmatrix} = \begin{pmatrix} \cos(a_A) & \sin(a_A) \\ -\sin(a_A) & \cos(a_A) \end{pmatrix} \begin{pmatrix} e_B \\ e_C \end{pmatrix}$$

$$\begin{pmatrix} e'_D \\ e'_E \end{pmatrix} = \begin{pmatrix} \cos(a_A) & -\sin(a_A) \\ \sin(a_A) & \cos(a_A) \end{pmatrix} \begin{pmatrix} e_D \\ e_E \end{pmatrix}$$

This mean that it is induced a rotation on the planes (e_B, e_C) and (e_D, e_E) with angles a_A and $-a_A$. Therefore the generator of this group action is

$$J_{BC} - J_{DE} \quad (2.195)$$

being J_{BC} and J_{DE} the anti-hermitian rotation generators. In similar manner the generators for the transformation corresponding to b_A can be obtained, the result is

$$\frac{1}{\sqrt{3}}(J_{BC} + J_{DE} - 2J_{FG}). \quad (2.196)$$

The indices run from 1 to 7 and therefore we have 14 generators. It is also direct to check that they close under commutation and therefore they are a subalgebra of $SO(7)$. This is the G_2 subalgebra. If we add

$$J_{BC} + J_{DE} + J_{FG} \quad (2.197)$$

it can be seen that (2.195), (2.196) and (2.197) generate $SO(7)$ [169].

As we recalled before, the associator

$$(a, b, c) = (ab)c - a(bc) \quad (2.198)$$

for three arbitrary octonions a , b and c is not zero and therefore the algebra is non associative. But (2.198) is fully antisymmetric and therefore the octonion algebra is *alternating*. Is therefore obvious that

$$(e_i, e_i, e_j) = 0$$

which implies that

$$c_{acd}c_{bcd} = 6\delta_{ab}. \quad (2.199)$$

The octonions also obey the Moufang identity [125]

$$(xa)(xb) = x(ab)x. \quad (2.200)$$

which implies that

$$c_{paq}c_{qbs}c_{scp} = 3c_{abc}. \quad (2.201)$$

Let us define the dual octonion constants

$$c_{ijkl} = \frac{1}{3!}\epsilon^{abcdefg}c_{efg}, \quad (2.202)$$

which are explicitly

$$c_{4567} = c_{2374} = c_{1357} = c_{1276} = c_{2356} = c_{1245} = c_{1346} = 1. \quad (2.203)$$

It can be checked the fundamental identity

$$c_{abcd} = (\delta_{bc}\delta_{ad} - \delta_{ac}\delta_{bd}) + c_{abe}c_{cde}. \quad (2.204)$$

The constant c_{abc} and c_{abcd} can be assembled into a single object ψ^{abcd} , with $a, b, \dots = 1, \dots, 8$ defined by

$$\psi_{abc8} = \tilde{c}_{abc}, \quad \psi_{abcd} = \tilde{c}_{abcd} \quad (2.205)$$

Equation (2.202) translates into the eight-dimensional self-duality of ψ_{abcd} , that is

$$\psi_{abcd} = \frac{1}{4!} \epsilon^{abcdefgh} \psi_{efgh}. \quad (2.206)$$

If we consider the 8×8 matrices γ^a given by

$$(\gamma^a)_{bc} = (\tilde{c}_{abc} + \delta_{a[b} \delta_{c]8}) \quad (2.207)$$

where the constants $\tilde{c}_{abc}$ are equal to zero if one of the index is equal to 8 and to the octonion constants in other case, then is seen that $\{\gamma^a, \gamma^b\} = 4\delta^{ab}I$. This means that γ^a are a representation of the Dirac algebra in seven dimensions and therefore

$$(\gamma^{ab})_{cd} = \frac{1}{4} [\gamma^a, \gamma^b]_{cd} = (\tilde{c}_{abcd} + \tilde{c}_{ab[c} \delta_{d]8} + \delta_{a[c} \delta_{d]b}) \quad (2.208)$$

are also a representation of the Lie algebra of $SO(7)$. Consider the spinor η with components $\eta_a = \delta_{ab}$. Then by use of the identities (2.201) and (2.199) it can be proved that

$$c_{mnp} = i\bar{\eta}\gamma^{mnp}\eta, \quad \gamma^{mnp} = \gamma^{[m}\gamma^n\gamma^{p]}. \quad (2.209)$$

By use of Fierz rearrangement identities in seven dimensions together with (2.209) it is possible to show many identities relating c_{abc} and c_{abcd} [168], [128], they are the following

$$c_{abe}c_{cde} = -c_{abcd} + \delta_{ac}\delta_{bd} - \delta_{ad}\delta_{bc} \quad (2.210)$$

$$c_{acd}c_{bcd} = 6\delta_{ab} \quad (2.211)$$

$$c_{abp}c_{pcde} = 3c_{a[cd}\delta_{e]b} - 3c_{b[cd}c_{e]a} \quad (2.212)$$

$$c_{abcp}c_{defp} = -3c_{ab[de}\delta_{f]c} - 2c_{def[a}\delta_{b]c} - 3c_{ab[d}c_{ef]c} + 6\delta_a^{[d}\delta_b^e\delta_c^{f]} \quad (2.213)$$

$$c_{abpq}c_{pqe} = -4c_{abc} \quad (2.214)$$

$$c_{abpq}c_{pqcd} = -2c_{abcd} + 4(\delta_{ac}\delta_{bd} - \delta_{ad}\delta_{bc}) \quad (2.215)$$

$$c_{apqr}c_{bpqr} = 24\delta_{ab} \quad (2.216)$$

$$\begin{aligned} c_{abp}c_{pcq}c_{qde} &= c_{abd}\delta_{ce} - c_{abe}\delta_{cd} - c_{ade}\delta_{bc} + c_{bde}\delta_{ac} \\ &\quad - c_{acd}\delta^{be} + c_{ace}\delta_{bd} + c_{bcd}\delta_{ae} - c_{bce}\delta_{ad} \end{aligned} \quad (2.217)$$

$$c_{paq}c_{qbs}c_{scp} = 3c_{abc} \quad (2.218)$$

Formulas (2.210)-(2.218) can also be derived from the basic identity for the product of two ψ [105]

$$\psi_{abcd}\psi^{efgd} = 6\delta_a^{[e}\delta_b^f\delta_c^g] - 9\psi_{[ab}^{[ef}\delta_{c]}^g]. \quad (2.219)$$

Consider the operators

$$(P_{14})_{ab}^{cd} = \frac{2}{3}(\delta_{ab}^{cd} + \frac{1}{4}c_{ab}^{cd}), \quad (P_7)_{ab}^{cd} = \frac{1}{3}(\delta_{ab}^{cd} - \frac{1}{2}c_{ab}^{cd}), \quad (2.220)$$

with $\delta_{ab}^{cd} = \frac{1}{2}(\delta_a^c \delta_b^d - \delta_a^d \delta_b^c)$. It is straightforward to see from (2.212) that $(P_{14})_{ab}^{cd} c_{cde} = 0$ and $(P_7)_{ab}^{cd} c_{cde} = c_{abe}$. Also from (2.215) we obtain

$$\begin{aligned}(P_{14})_{ef}^{ab} (P_7)_{ab}^{cd} S_{cd} &= 0, \\ (P_{14})_{ef}^{ab} (P_7)_{ab}^{cd} S_{cd} &= 0, \\ [(P_{14})_{ab}^{cd} + (P_7)_{ab}^{cd}] S_{cd} &= S_{cd}\end{aligned}$$

and therefore (2.220) are orthogonal projectors. This means that for any tensor S^{ab} that transforms in the adjoint 21 representation of $SO(7)$ we have the decomposition

$$S_+^{ab} = \frac{2}{3}(S^{ab} + \frac{1}{4}c^{abcd}S^{cd}), \quad c^{abc}S_+^{bc} = 0, \quad (2.221)$$

$$S_-^{ab} = \frac{1}{3}(S^{ab} - \frac{1}{2}c^{abcd}S^{cd}) = c^{abc}\zeta^c, \quad \zeta^c = \frac{1}{6}c^{cab}S^{ab}, \quad (2.222)$$

$$S^{ab} = S_+^{ab} \oplus S_-^{ab}. \quad (2.223)$$

This is analogous to the fact in $D = 4$ that any tensor transforming under the adjoint representation $SO(4) \sim SU(2)_L \times SU(2)_R$ is the direct sum of its self-dual and anti-self-dual parts. We will say that S_+^{ab} and S_-^{ab} are the self-dual and antiself-dual parts of S^{ab} . From the facts stated above we see that

$$S_+^{ab} = \frac{1}{2}c_{abcd}S_+^{cd} \quad (2.224)$$

$$S_-^{ab} = -\frac{1}{4}c_{abcd}S_-^{cd} \quad (2.225)$$

that are analogous to the well known self-duality condition $F^{ab} = \pm \epsilon^{abcd}F^{cd}$ in $D = 4$ but with the octonion constants replacing the Kronecker symbols. By orthogonality of P_{14} and P_7 also holds the relation

$$S^{ab}U^{ab} = S_+^{ab}U_+^{ab} + S_-^{ab}U_-^{ab}. \quad (2.226)$$

being U^{ab} another $SO(7)$ tensor. The projectors (2.220) induce an isomorphism of the $SO(7)$ algebra given by

$$G^{ab} = \frac{2}{3}(\gamma^{ab} + \frac{1}{4}c_{abcd}\gamma^{cd}), \quad (2.227)$$

$$C^{ab} = \frac{1}{3}(\gamma^{ab} - \frac{1}{2}c_{abcd}\gamma^{cd}). \quad (2.228)$$

From (2.192) and (2.203) we can see that there are 14 generators G^{ab} and 7 generators C^{ab} . G^{ab} are a representation of the Lie algebra of G_2 .

The group G_2 is the subgroup of $SO(7)$ has a one dimensional invariant subspace generated by certain vector η . This mean that $A\eta = \eta$ being A an element of G_2 . This implies that $G^{ab}\eta = 0$, being G_{ab} the infinitesimal generators of the group G_2 . For the representation (2.227) this subspace is generated by a spinor with components $\eta_a = \delta_{a8}$. This is the decomposition $8 \rightarrow 7 + 1$ of $SO(7)$. The form of η depends on the representation, but its existence of course not. G_2 is also the group which leaves invariant the three-form

$$\Phi = \frac{1}{3!}c_{abc}e^a \wedge e^b \wedge e^c \quad (2.229)$$

and the 4-form

$$*\Phi = \frac{1}{3!}c_{abcd}e^a \wedge e^b \wedge e^c \wedge e^d \quad (2.230)$$

being $e_1, \dots, e_7$ a basis for a seven dimensional space V . The closure of (2.229) and (2.230) is one of the main facts of G_2 holonomy manifolds, as we will see in the following.

2.6.2 G_2 holonomy and self-duality

In the previous subsection we have defined the Lie group G_2 as the subgroup of $SO(7)$ that has one and only one stable spinor η , also as the automorphism group of the octonions, and as the group that leave the 3-form (2.229) invariant. We have also seen that the G_2 component of a tensor transforming under the adjoint representation 21 of $SO(7)$ is self-dual (2.221). In this section we will show that the analog holds for an space M with holonomy G_2 that is, the presence of one Killing spinor η globally defined on it, or that there is defined a three-form that is closed and co-closed over it. Moreover for a G_2 holonomy space it exist a rotation of the frame which take to zero the anti-self-dual part of the spin connection. Therefore the curvature of a G_2 holonomy metric is self-dual.

It is evident that the holonomy of an arbitrary seven dimensional manifold M is always included in $SO(7)$. From the discussion of the previous subsection it follows that is reduced to G_2 if and only if there exist only one Killing spinor η , that is, an spinor globally defined over M satisfying the condition

$$D_i \eta = 0, \quad (2.231)$$

being $D_i = I\partial_i + \omega_{ib}^a \gamma^{ab}$ the covariant derivative in spinor representation. In other words the existence of a globally covariant spinor allows the reduction $8 \rightarrow 7 + 1$ to 7 of the holonomy group.

If we choose a the representation of $SO(7)$ given by (2.227) and (2.228), and the spinor η such that $\eta_a = \delta_{ab}$ it follows that $(\gamma^{ab}\eta)_c = c_{abc}$. Then

$$D_i \eta = -\frac{1}{4} \omega_{iab} \gamma^{ab} \eta = -\frac{1}{4} \omega_{iab} c_{abc} = 0.$$

It can be checked by using (2.192) and (2.203) that the last condition is equivalent to

$$\omega_b^a = \frac{1}{2} c_{abcd} \omega_d^c. \quad (2.232)$$

A 7-dimensional space for which ω_b^a satisfies (2.232) is called self-dual. We conclude that self-dual spaces in $D = 7$ have G_2 holonomy. It can be proved that this result is independent of the representation of $SO(7)$ and that the converse is also true, that is, for G_2 holonomy there exists always a rotation of the frame for which the anti self-dual part ω_- goes locally to zero [128].

G_2 manifolds are always Ricci-flat. In fact, if we apply D_j to (2.231) and anti-symmetrize we will obtain

$$R_{bcd}^a \gamma^{ab} \eta = 0$$

where $R_b^a = R_{bcd}^a e^c \wedge e^d$ is the curvature tensor. Now if we multiply by γ^i this expression and using that $R_{abcd} = R_{cdab} = -R_{abdc}$ and that $R_{[ebc]}^a = 0$ we find that

$$R_{ij} \gamma^j \eta = 0 \quad \Leftrightarrow \quad R_{ij} \gamma^j = 0$$

But the matrices γ^i are linearly independent and therefore $R_{ij} = 0$. Indeed, in any dimension, the presence of a covariantly constant spinor implies Ricci-flatness by the reasoning given above.

It is well known that in $D = 4$ the self-duality of the spin connection implies the self-duality of the curvature tensor. This is also true in seven dimensions. In [128] a clear proof was given. It is based in the following identity for the octonion constants

$$c_{abcp} c_{defp} = -3c_{ab[de} \delta_{e]b} - 2c_{def[a} \delta_{b]c} + 6\delta_a^{[d} \delta_b^e \delta_c^{f]} - 2c_{def[a} \delta_{b]c}. \quad (2.233)$$

If (2.232) holds it is clear that $d\omega_b^a$ will be self-dual. The self-duality of $\omega_c^a \wedge \omega_b^c$ follows using that

$$\begin{aligned} \frac{1}{2}c_{abcd}\omega_e^c \wedge \omega_d^e &= \frac{1}{4}c_{abcd}c_{edfg}\omega_e^c \wedge \omega_g^f \\ &= -\frac{1}{2}c_{abfe}\omega_e^f \wedge \omega_c^e + \frac{1}{4}c_{aefg}\omega_e^b \wedge \omega_g^f - \frac{1}{4}c_{befg}\omega_e^a \wedge \omega_g^f + \omega_c^a \wedge \omega_b^c \\ &= -\frac{1}{2}c_{abcd}\omega_e^c \wedge \omega_d^e + 2\omega_c^a \wedge \omega_b^c. \end{aligned} \quad (2.234)$$

The identity (2.233) has been used here. From (2.234) is seen that

$$\frac{1}{2}c_{abcd}\omega_e^c \wedge \omega_d^e = \omega_c^a \wedge \omega_b^c$$

which implies the self-duality of $R = d\omega + \omega \wedge \omega$.

From (2.232), the antisymmetry of (2.203) and the Bianchi identity $R_{d[ebc]} = 0$ it follows that

$$R_{ab} = R_{acbc} = \frac{1}{2}c_{acde}R_{debc} = \frac{1}{2}c_{acde}R_{d[ebc]} = 0. \quad (2.235)$$

This is another proof that G_2 holonomy spaces are Ricci-flat. For such case the Weyl tensor W_{abcd} defined by

$$W_{abcd} = R_{abcd} + \frac{R}{(n-1)(n-2)}(g_{ac}g_{bd} - g_{ad}g_{bc}) - \frac{1}{(n-2)}(g_{ac}R_{bd} - g_{ad}R_{bc} - g_{bc}R_{ad} + g_{bd}R_{ac}) \quad (2.236)$$

will be equal to the Riemann tensor. The first one is traceless, in consequence Ricci-flat manifolds have traceless curvature tensor.

The condition to have holonomy G_2 is also equivalent to the existence of a closed and co-closed three form [27]. The natural candidates to consider are the G_2 equivariant 3-form Φ (2.229) and its dual $*\Phi$ [128]

$$\Phi = \frac{1}{3!}c_{abc}e^a \wedge e^b \wedge e^c \quad *\Phi = \frac{1}{4!}c_{abcd}e^a \wedge e^b \wedge e^c \wedge e^d. \quad (2.237)$$

Taking into account the identity

$$c_{abp}c_{pcde} = 3c_{a[cd}\delta_{e]b} - 2c_{b[cd}\delta_{e]a}$$

it is obtained

$$\begin{aligned} d\Phi &= -\frac{1}{3!2}c_{abc}e^a \wedge e^b \wedge \omega^{cd} \wedge e^d = -\frac{1}{3!}c_{abc}c_{cdef}e^a \wedge e^b \wedge \omega^{ef} \wedge e^d \\ &= \frac{1}{3!}c_{ade}e^a \wedge e^d \wedge \omega_{eb} \wedge e^b = -2d\Phi. \end{aligned}$$

From here follows

$$d\Phi = 0. \quad (2.238)$$

So, Φ is a closed form. Similarly, using (2.233) follows that

$$\begin{aligned} d*\Phi &= -\frac{1}{4!6}c_{abcd}\omega_{ae} \wedge e^e \wedge e^b \wedge e^c \wedge e^d = -\frac{1}{4!12}c_{aefg}c_{abcd}\omega^{fg} \wedge e^e \wedge e^b \wedge e^c \wedge e^d \\ &= \frac{1}{4!3}c_{fbcd}\omega^{fe} \wedge e^e \wedge e^b \wedge e^c \wedge e^d = -2d*\Phi, \end{aligned}$$

which implies the closure of $*\Phi$. This proves that G_2 holonomy implies the closure of Φ and $*\Phi$. But in [27] it has been proved that the closure and co-closure of Φ and the constance of η give the same number of independent equations and therefore, both conditions are indeed equivalent.

2.6.3 The Bryant-Salamon construction

We present in this subsection an extension of a four dimensional quaternionic Kahler space to a seven dimensional one with holonomy lying in G_2 . This construction was performed by Bryant and Salamon in [29] and considered recently in [99]. In this section we intend to present such construction not in the original form [29], but in a way more clear for physicists.

Consider a 7-manifold $M = R^3 \times M^4$ with metric

$$g = (du^1)^2 + (du^2)^2 + (du^3)^2 + g_h \quad (2.239)$$

being g_h 4-dimensional metric doesn't depending of the tree coordinates u_i . Condition (2.232) reduce for (2.239) to

$$\omega_j^i = \frac{1}{2} \epsilon_{ijkl} \omega_l^k$$

being ω_j^i the spin connection of g_h . Therefore if g_h has self-dual connection or at least there exists locally a frame in which the anti-self-dual part of the connection goes to zero (and therefore g_h is hyperkahler) then $g = g_{R^3} \oplus g_h$ has holonomy included in $SU(2) \subset G_2$. We can generalize the ansatz (2.239) and consider instead

$$g = e^{2g} [(D_- u^1)^2 + (D_- u^2)^2 + (D_- u^3)^2] + e^{2f} g_q, \quad (2.240)$$

where g_q is a quaternionic Kahler space in four dimension and we had replaced the usual differentials du^i by the covariant derivative

$$D_- u^i = du^i + \epsilon^{ijk} \omega_-^j u^k. \quad (2.241)$$

As usual ω_-^i is the anti-self-dual part of the spin connection. The metric (2.239) corresponds to the hyperkahler limit (in which locally we can select $\omega_-^i = 0$) with f and g reduced to constants. We select f and g as functions on the radius $|u| = \sqrt{u_i^2}$. Then condition (2.232) for (2.240) is a constraint to be satisfied by f and g the resulting metric will have holonomy $\Gamma \subset G_2$.

The explicit calculation of the connection ω gives

$$\begin{aligned} \omega^{ij} &= \frac{f'}{|u|} D_- u^{[i} u^{j]} - \epsilon^{ijk} \omega_-^k \\ \omega^{mn} &= \omega^{mn} - \kappa e^{2(f-g)} \epsilon^{ijk} u^k J_{mn}^j D_- u^i \\ \omega^{mi} &= \frac{g'}{|u|} e^{g-f} e^m u^i - \kappa e^{f-g} \epsilon^{ijk} u^k J_{mn}^j e^n \end{aligned} \quad (2.242)$$

where $(...)'$ denotes the derivative with respect to $|u|$. The self-duality condition (2.232) gives the equations

$$\begin{aligned} 0 &= c_{iab} \omega^{ab} = c_{ijk} \omega^{jk} + c_{imn} \omega^{mn} = \left[\frac{f'}{|u|} + \kappa e^{2(f-g)} \right] \epsilon^{ijk} D_- u^j u^k \\ 0 &= c_{mab} \omega^{ab} = c_{mni} \omega^{ni} = J_{mn}^i \omega^{ni} = \left[\frac{g'}{|u|} - \kappa e^{2(f-g)} \right] e^{g-f} J_{mn}^i e^n u^i. \end{aligned} \quad (2.243)$$

Equations (2.243) are invariant under $f \rightarrow f + \lambda$, $g \rightarrow g + \lambda$ which give a constant conformal scaling of the metric (2.245). Also we have the second symmetry $f \rightarrow f - \lambda$, $u \rightarrow e^\lambda u$ which leaves the metric invariant. Using these symmetries the solution result

$$e^{-4f} = 2\kappa |u|^2 + c, \quad e^{4g} = 2\kappa |u|^2 + c. \quad (2.244)$$

By introducing (2.244) into (2.245) we obtain the Bryant-Salamon family of G_2 metrics

$$g = \frac{1}{\sqrt{2\kappa|u|^2 + c}}(du^i + \epsilon^{ijk}\omega_-^j u^k)^2 + \sqrt{2\kappa|u|^2 + c} g_q. \quad (2.245)$$

Clearly in the limit $\omega_-^j = 0$ and $\kappa = 0$ we recover (2.239).

The Bryant-Salamon metrics (2.245) are non-compact, because the coordinates u_i can take any real value, and this represent a problem when consider applications to compactification of M -theory. Nevertheless they have the important property to be asymptotically conical. To see that they are conical let us write (2.245) introducing spherical coordinates

$$u_1 = |u| \cos(\theta),$$

$$u_2 = |u| \sin(\theta) \cos(\varphi),$$

$$u_3 = |u| \sin(\theta) \sin(\varphi),$$

and defining the new radial coordinate

$$r^2 = \sqrt{2\kappa|u|^2 + c}.$$

Then it is obtained

$$g = \frac{dr^2}{(1 - \frac{4c}{r^4})} + \frac{r^2}{4\kappa}(1 - \frac{4c}{r^4})g_{ab}(dx^a + \xi_i^a \omega_-^i)(dx^b + \xi_j^b \omega_-^j) + \frac{r^2}{2}g_q, \quad (2.246)$$

where g_{ab} and ξ are the metric and the killing vectors of S^2 respectively. The corresponding 7-bein is explicitly

$$e_i = \sqrt{\frac{\kappa}{2}} r V_i$$

$$e_5 = \frac{r}{2}(d\theta - \sin(\varphi)\omega_-^2 + \cos(\varphi)\omega_-^3)$$

$$e_6 = \frac{r}{2}(\sin(\theta)d\varphi + \sin(\theta)\omega_-^1 - \cos(\theta)\cos(\varphi)\omega_-^2 - \cos(\theta)\sin(\varphi)\omega_-^3)$$

$$e_7 = \frac{dr}{(1 - \frac{4c}{r^4})^{1/2}}$$

being V_i the einbeins of the quaternion base g_q . If we consider $r^2 \gg c$ then (2.246) is

$$g \approx dr^2 + r^2 \Omega, \quad (2.247)$$

being Ω the metric of a 6 manifold independent of the coordinate r . Thus the family (2.245) is conical. The explicit form of Ω is

$$\Omega = g_{ab}(dx^a + \xi_i^a \omega_-^i)(dx^b + \xi_j^b \omega_-^j) + g_q. \quad (2.248)$$

The metric (2.248) is of weak $SU(3)$ holonomy, we will return to this point below.

It is instructive to check also the existence of a closed and co-closed 3-form. It is seen by (2.229) that

$$\Phi = \frac{1}{6}(1 + |u|^2)^{-3/4} \epsilon_{ijk} D_- u_i \wedge D_- u_j \wedge D_- u_k + 2(1 + |u|^2)^{1/4} D_- u_i \wedge J^i \quad (2.249)$$

and the dual $*\Phi$ is

$$*\Phi = 4(1 + |u|^2) \Theta + \epsilon_{ijk} D_- u_i \wedge D_- u_j \wedge J^k. \quad (2.250)$$

The 4-form Θ is defined in (2.19) and in four dimensions is equal to the volume form of M

$$\Theta = \frac{1}{2} \bar{J}^1 \wedge \bar{J}^1 = \frac{1}{2} \bar{J}^2 \wedge \bar{J}^2 = \frac{1}{2} \bar{J}^3 \wedge \bar{J}^3.$$

The covariant derivative D_- is given by (2.241) and it follows that $(D_-)^2 u_i = \epsilon_{ijk} F^j u_k$. For a quaternionic base $d\Theta = 0$ and from (2.7) it follows that $D_- \bar{J}^a = 0$. It can be shown by use of this relations that Φ is closed and co-closed.

2.6.4 Weak G_2 and $Spin(7)$ holonomy spaces

As it follows from the previous sections G_2 holonomy manifolds admits a globally defined Killing spinor η . In this subsection we study eight dimensional spaces with the same property. Also we study seven dimensional ones that admits a conformal Killing spinor, that is, an spinor satisfying $D_j \eta \sim \eta$. For the former the holonomy is included in $Spin(7)$. The second have generically $SO(7)$ holonomy but are defined by certain conditions that are analog of the G_2 case. For this reason we call such them "weak G_2 holonomy manifolds" [177].

In seven dimension the presence of conformally Killing spinor η namely

$$D_i \eta = i\lambda \gamma_i \eta, \quad (2.251)$$

is a generalization of G_2 holonomy condition (2.231), which is obtained as the limit $\lambda \rightarrow 0$ of (2.251). If we act on (2.251) by D_i and antisymmetrize it is obtained

$$R_{ij}^{ab} \gamma_{ab} \eta = 8\lambda^2 \gamma_{ij} \eta.$$

Now if we multiply the last expression by γ^k and using that $R_{abcd} = R_{cdab} = -R_{abdc}$ and that $R_{[ebc]}^a = 0$ we get

$$R_{kj} \gamma^j \eta = 4(d-1) \tilde{\lambda}^2 \gamma_k \eta$$

which implies

$$R_{ij} = 4(d-1) \lambda^2 \delta_{ij}. \quad (2.252)$$

Therefore if there exist a globally defined η satisfying (2.251) the metric is Einstein $R_{ij} \sim g_{ij}$ but with cosmological constant different than zero. Spaces satisfying condition (2.251) have holonomy that lies in $SO(7)$ and just in the limit $\lambda \rightarrow 0$ we can state that the holonomy is reduced to a subgroup of G_2 . For $\lambda \neq 0$ the space are weak G_2 holonomy ones [177], [128].

As before, let us select the spinor $\eta_i = \delta_{i8}$, then

$$D_j \eta = i\lambda \gamma_j \eta$$

together with (2.207) and (2.208) implies that

$$c_{abc} \omega_j^{ab} = -\frac{\lambda}{2} \delta_{jc} \quad \Longleftrightarrow \quad c_{abc} \omega^{bc} = -\frac{\lambda}{12} e^a. \quad (2.253)$$

The relation (2.253) is equivalent to

$$\omega_-^{ab} = -\frac{\lambda}{12} c_{abc} e^c \quad \Longleftrightarrow \quad \omega^{ab} = \frac{1}{2} c_{abcd} \omega^{cd} - \frac{\lambda}{4} c_{abc} e^c. \quad (2.254)$$

Condition (2.254) is the weak self-duality condition in seven dimensions. If for a manifold there exists a frame for which (2.254) hold then it will be weak G_2 holonomy type.

The form Φ is defined as before in (2.237) and using (2.253)-(2.254) we obtain

$$\begin{aligned} d\Phi &= -\frac{1}{2}c_{abc}e^a \wedge e^b \wedge \omega^{cd} \wedge e^d = -\frac{1}{4}c_{abc}c_{cdef}e^a \wedge e^b \wedge \omega^{ef} \wedge e^d \\ &\quad + \frac{\lambda}{8}c_{abc}c_{cde}e^a \wedge e^b \wedge e^e \wedge e^d = -2d\Phi + 3\lambda * \Phi. \end{aligned} \quad (2.255)$$

where we used (2.210) in the last step. So we conclude that

$$d\Phi = \lambda * \Phi, \quad (2.256)$$

and the last condition implies that $d * \Phi = 0$. As expected, if the cosmological constant is zero (2.256) reduce to the closure and co-closure of Φ and the holonomy is again in G_2 .

Weak G_2 metrics can be constructed as the angular part of certain conical $Spin(7)$ holonomy spaces, that we will present in the following. This is analogous to the fact that weak $SU(3)$ manifolds are the angular part of certain conical G_2 holonomy manifolds. $Spin(7)$ spaces have a globally defined Killing spinor η as their seven dimensional Ricci flat counterpart.

$Spin(7)$ is the subgroup of $SO(8)$ preserving the 4-form

$$\hat{\Phi} = \psi_{abcd}e^a \wedge e^b \wedge e^c \wedge e^d \quad (2.257)$$

being ψ_{abcd} the self-dual constants associated to octonions defined in (2.205). Because we are in the middle dimension, we see that $*\Phi = \Phi$. Intuition could suggest that as in the G_2 case, the holonomy will be reduced from $SO(8)$ to $Spin(7)$ if $d\hat{\Phi} = 0$.

Let us consider the $SO(8)$ γ -matrices $\Gamma_a = (\Gamma_a, \Gamma_8)$, with $a = 1, \dots, 8$

$$\Gamma_a = \begin{pmatrix} 0 & i\gamma_a \\ -i\gamma_a & 0 \end{pmatrix}, \quad \Gamma_8 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \{\Gamma_a, \Gamma_b\} = 2\delta_{ab}, \quad (2.258)$$

that correspond to the standard embedding of $SO(7)_v$ in $SO(8)$. The $SO(8)$ generators $\Gamma^{ab} = \frac{1}{2}[\Gamma^a, \Gamma^b]$ satisfy the relations

$$\Gamma^{ab} = \epsilon^{abc}\Gamma^9\Gamma_{8c}, \quad c_{abcd}\Gamma^{ab} = -(4 + 2\Gamma^9)\Gamma_{cd}, \quad (2.259)$$

where $\Gamma^9 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ is the chirality matrix. Then it can be shown that the right handed part of an spinor η , namely $\eta_+ = (1 - \Gamma^9)\eta$ is invariant under the action of

$$G^{ab} = \frac{3}{8} \left(\Gamma^{ab} + \frac{1}{6} \psi_{cd}^{ab} \Gamma^{cd} \right), \quad (2.260)$$

that is

$$G^{ab}\eta_+ = 0. \quad (2.261)$$

We can define as before the associated projectors

$$(P_{21})_{ab}^{cd} = \frac{3}{8}(\delta_{ab}^{cd} + \frac{1}{6}\psi_{ab}^{cd}), \quad (P_7)_{ab}^{cd} = \frac{5}{8}(\delta_{ab}^{cd} - \frac{1}{10}\psi_{ab}^{cd}), \quad (2.262)$$

corresponding to the embedding of $Spin(7) \subset SO(8)$. Let us call as before A_+^{ab} and A_-^{ab} the part annihilated by P_7 , or P_{21} respectively of an $SO(8)$ tensor A^{ab} . As in the seven dimensional case the $Spin(7)$ part A_+^{ab} satisfies a self duality condition related to the constants ψ_{abcd} namely

$$A_+^{ab} = \frac{1}{2} \hat{c}_{abcd} A_+^{cd}. \quad (2.263)$$

Moreover, there is a Killing spinor η for which $A_+^{ab}\eta = 0$.

From the previous paragraph it follows that a manifold M has $Spin(7)$ holonomy if and only if there is a global Killing spinor η for which $D_i\eta = 0$ or equivalently, if its connection ω_b^a satisfies

$$\omega_b^a = \frac{1}{2} \hat{c}_{abcd} \omega_d^c. \quad (2.264)$$

The last condition is equivalent to the closure of $\hat{\Phi}$, namely

$$d\hat{\Phi} = 0. \quad (2.265)$$

We omit the proof because they are similar to those presented in the G_2 case. $Spin(7)$ manifolds are of course Ricci-flat as far as they admit a Killing spinor η . If we select one direction, say 8, we obtain from (2.264) that

$$\omega_r^8 = -\frac{1}{2} c_{rpq} \omega_q^p. \quad (2.266)$$

The rest of (2.264), namely

$$\omega_q^p = \frac{1}{2} c_{qrs} \omega_s^r - c_{qr}^p \omega^r{}^8$$

are automatically satisfied if eq.(2.266) holds.

There exists a Bryant-Salamon extension of a quaternionic Kahler space to one with $Spin(7)$ holonomy. It is possible to construct such spaces in a way similar to the presented for the Swann spaces (2.184). This is because it should be imposed $d\hat{\Phi} = 0$ for (2.257), which is the analog of the Swann 4-form (2.176) for the $Spin(7)$ case. We start with the ansatz (2.174) for the eight dimensional metric, namely

$$g_s = g\bar{g} + f|du + u\omega_-|^2.$$

The quaternion u and the anti-self-dual ImH valued 1-form ω_- are

$$u = u_0 + u_1I + u_2J + u_3K, \quad \bar{u} = u_0 - u_1I - u_2J - u_3K$$

$$\omega_- = \omega_-^1 I + \omega_-^2 J + \omega_-^3 K$$

$$|u|^2 = x\bar{x} = |u|^2 = (u_0)^2 + (u_1)^2 + (u_2)^2 + (u_3)^2,$$

being I, J, K the unit quaternions. The functions f and g depends only on the radius $|u|$. Condition $d\hat{\Phi} = 0$ gives a differential equation for f and g . The explicit expression of Φ is obtained from (2.257) and is

$$\begin{aligned} \hat{\Phi} = & 3fg[\alpha \wedge \bar{\alpha} \wedge \bar{e}^t \wedge e + \bar{e}^t \wedge e \wedge \alpha \wedge \bar{\alpha}] \\ & + g^2 \bar{e}^t \wedge e \wedge \bar{e}^t \wedge e + f^2 \alpha \wedge \bar{\alpha} \wedge \alpha \wedge \bar{\alpha} \end{aligned} \quad (2.267)$$

where $\alpha = du + u\omega_-$. Then is straightforward by use of (2.177)-(2.180) to impose $d\Phi = 0$ and find the system defining f and g . The final solution is

$$f = \frac{1}{(2\kappa|u|^2 + c)^{2/5}},$$

$$g = (2\kappa|u|^2 + c)^{3/5}.$$

with the corresponding metric

$$g_s = (2\kappa|u|^2 + c)^{3/5} \bar{g} + \frac{1}{(2\kappa|u|^2 + c)^{2/5}} |\alpha|^2. \quad (2.268)$$

Spaces defined by (2.268) are the Bryant-Salamon $Spin(7)$ ones.

Metrics (2.268) are non compact (because $|u|$ is not bounded), and asymptotically conical. Introducing spherical coordinates for u gives

$$\begin{aligned} u_1 &= |u| \cos(\theta), \\ u_2 &= |u| \sin(\theta) \sin(\varphi), \\ u_3 &= |u| \sin(\theta) \cos(\varphi) \sin(\phi), \\ u_4 &= |u| \sin(\theta) \cos(\varphi) \cos(\phi), \end{aligned}$$

and defining the radial variable

$$r^2 = \frac{9}{20} (2\kappa|u|^2 + c)^{3/5}$$

we obtain the spherical form of the metric

$$g = \frac{dr^2}{\kappa(1 - c/r^{10/3})} + \frac{9}{100\kappa} r^2 \left(1 - \frac{c}{r^{10/3}}\right) (\sigma^i - \omega_-^i)^2 + \frac{9}{20} r^2 g_q \quad (2.269)$$

being σ^i the left-invariant one-forms on $SU(2)$

$$\begin{aligned} \sigma_1 &= \cos(\varphi) d\theta + \sin(\varphi) \sin(\theta) d\phi \\ \sigma_2 &= -\sin(\varphi) d\theta + \cos(\varphi) \sin(\theta) d\phi \\ \sigma_3 &= d\varphi + \cos(\theta) d\phi. \end{aligned}$$

If we consider the limit $r \gg c$ we find the behavior

$$g \approx dr^2 + r^2 \Omega, \quad (2.270)$$

with Ω a seven dimensional metric independent of the coordinate r , namely

$$\Omega = (\sigma^i - \omega_-^i)^2 + g_q \quad (2.271)$$

We also see that the subfamilies of (2.269) with $c = 0$ are exactly conical and their angular part is (2.271). Such subfamilies are known as cohomogeneity one metrics.

There is a correspondence between cohomogeneity one $Spin(7)$ spaces and weak G_2 holonomy ones, and in particular we can prove that the cone (2.271) is of weak G_2 type. From (2.270) it is obtained the seven-bein $\tilde{e}^a$ and the connection $\tilde{\omega}^{ab}$ of an eight dimensional cohomogeneity one space, namely

$$\tilde{e}^8 = dr, \quad \tilde{e}^a = -\frac{\lambda}{4} r e^a. \quad (2.272)$$

being e^a the einbein of the angular part. The first Cartan structure

$$d\tilde{e}^a + \tilde{\omega}^{ab} \wedge \tilde{e}^b = 0$$

implies that

$$\omega^{ab} = \tilde{\omega}^{ab}, \quad \tilde{\omega}^{8a} = \frac{\lambda}{4} e^a. \quad (2.273)$$

From the einbeins (2.272) and (2.257) we obtain

$$\begin{aligned} \tilde{\Phi} &= \left(\frac{\lambda r}{4}\right)^3 dr \wedge \Phi + \left(\frac{\lambda r}{4}\right)^4 * \Phi, \\ d\tilde{\Phi} &= -\left(\frac{\lambda r}{4}\right)^3 dr \wedge (d\Phi - \lambda * \Phi) + \left(\frac{\lambda r}{4}\right)^4 d * \Phi, \end{aligned} \quad (2.274)$$

where Φ and $*\Phi$ are the usual seven-dimensional 3- and 4-form constructed from the e^a . We see from (2.274) that *Spin*(7) holonomy condition, namely $d\tilde{\Phi} = 0$ is equivalent to weak G_2 holonomy of the seven-dimensional base space, that is, to the condition

$$d\Phi - \lambda * \Phi = 0, \quad d * \Phi = 0.$$

and the converse is also true. Moreover equations (2.272) are exactly equivalent to

$$\tilde{\omega}^{ab} = \frac{1}{2} \psi_{abcd} \tilde{\omega}^{cd}.$$

which is the *Spin*(7) self-duality condition (2.264). Then we conclude that there is a one to one correspondence between co-homogeneity one *Spin*(7) and weak G_2 metrics [39].

2.7 G_2 holonomy spaces and M-theory compactifications

In this subsection we show that G_2 and weak G_2 holonomy manifolds are internal spaces of M-theory that preserve at least one supersymmetry after Kaluza-Klein reduction to four dimensions. The bosonic lagrangian of the SUGRA theory in $D = 11$, that is the low energy effective action of the M-theory is

$$\mathcal{L}_{bos} = \frac{1}{2\kappa^2} [eR - \frac{e}{48} F_{MNPQ} F^{MNPQ} - \frac{1}{3!4!4!} \epsilon^{m_1 \dots m_{11}} F_{m_1 \dots m_4} F_{m_5 \dots m_8} A_{m_9 m_{10} m_{11}}] \quad (2.275)$$

being $F_{MNPQ} = 24 \partial_{[M} A_{NPQ]}$ and $R = R_{MN}^{AB} e_A^M \wedge e_B^N$. We are interested in the bosonic part of the theory because we want to compactify it to obtain a four dimensional theory with maximal space-time symmetry [26]. This means that the vacuum should be invariant under $SO(1,4)$ or $SO(2,3)$ as the cosmological constant is positive, negative or zero, corresponding to the de Sitter, Minkowski or anti de Sitter spaces. The first requirement of maximal symmetry is that the vacuum expectation value of the Rarita-Schwinger fermion should vanish, therefore we set

$$\langle \Psi_M \rangle = 0.$$

The equations of motion in this case reduce to

$$R_{MN} - \frac{1}{2} g_{MN} R = \frac{1}{3} [F_{MNPQ} F_N^{PQR} - \frac{1}{8} g_{MN} F_{PQRS} F^{PQRS}] \quad (2.276)$$

$$\nabla_m F^{MNPQ} = -\frac{1}{576} \epsilon^{m_1 \dots m_8 MNPQ} F_{m_1 \dots m_4} F_{m_5 \dots m_8} \quad (2.277)$$

Following the Kaluza-Klein procedure we consider the product space $M_4 \times M_7$ with the eleven dimensional einbein

$$e_m^a(X^n) = \begin{pmatrix} e_\mu^m(x^\nu) & 0 \\ 0 & e_i^a(x^j) \end{pmatrix}, \quad (2.278)$$

the greek coordinates will correspond to M_4 and the latin to M_7 . This means that the metric is factorized as

$$g_{\alpha\beta} = g_{\alpha\beta}(x^\mu), \quad g_{ab} = g_{ab}(x^m), \quad ; \quad g_{a\alpha} = 0 \quad (2.279)$$

$$F_{\alpha\beta\gamma\delta} = F_{\alpha\beta\gamma\delta}(x^\mu) \quad F_{abcd} = F_{abcd}(x^m) \quad F_{abc\alpha}(x^m) = F_{ab\alpha\beta}(x^m) = F_{a\alpha\beta\gamma}(x^m) = 0 \quad (2.280)$$

The x^α independence of F_{abcd} and the x^a independence of $F_{\alpha\beta\gamma\delta}$ are a consequence of the Bianchi identity

$$\partial_{[M} F_{NPQR]} = 0, \quad (2.281)$$

together with (2.280). Let us introduce the following ansatz

$$F_{\mu\nu\rho\sigma} = A \det(e_\mu^m) \epsilon_{\mu\nu\rho\sigma}, \quad F_{ijkl} = B e_i^a e_j^b e_k^c e_l^d c^{abcd}, \quad (2.282)$$

into the equations (2.276), with A and B constants. If we set $B = 0$ then we obtain the well known Freund Rubin ansatz [178], but we will not assume this for the moment. The Einstein equations are

$$R_{\mu\nu} = -\frac{1}{3}(A^2 + \frac{7}{2}B^2)g_{\mu\nu} \quad R_{ij} = \frac{1}{6}(A^2 + 5B^2)g_{ij} \quad (2.283)$$

and we see that for any non trivial selection of A and B , i.e., non trivial backgrounds F_{MNPQ} the four dimensional part is Einstein with negative curvature. The equations of motion for F_{MNPQ} reduces to

$$B(d\Phi - A * \Phi) = 0, \quad (2.284)$$

where Φ is the three-form

$$\Phi = \frac{1}{3!4!} e_\mu^m \det(e_i^a) \epsilon_{ijklmnp} F^{lmnp} dx^i \wedge dx^j \wedge dx^k = e_i^a e_j^b e_k^c c^{abc} dx^i \wedge dx^j \wedge dx^k. \quad (2.285)$$

Now we turn our attention to the criterion for unbroken supersymmetry in the effective $D = 4$ theory. If the vacuum is supersymmetric, then condition $\langle \Psi_m \rangle = 0$ should be invariant under supersymmetry transformations, that is

$$\delta\psi_M = \mathcal{D}_M \epsilon - \frac{1}{288} F_{PQRS} \left(\Gamma^{PQRS}{}_M - 8 \Gamma^{QRS} \delta_M^P \right) \epsilon = 0, \quad (2.286)$$

In particular, invariance (2.286) impose restrictions over the internal manifold. Taking into account the $O(3,1) \times O(7)$ decomposition of the gravitino and the gamma matrices given by

$$\begin{aligned} \psi_M &\longrightarrow \psi_\mu \otimes \hat{\psi}_\alpha, & \psi &\otimes \hat{\psi}_{i,\alpha} \\ \Gamma^M &\longrightarrow \gamma^\mu \otimes I, & \gamma_5 &\otimes \gamma^i, \end{aligned}$$

we obtain that

$$\begin{aligned} \delta(\psi_\mu \otimes \hat{\psi}_\alpha) &= \mathcal{D}_\mu(\epsilon \otimes \hat{\epsilon}_\alpha) + \frac{1}{12} \tilde{B}(\gamma_\mu \epsilon) \otimes (\delta_{\alpha\beta} - 8\delta_{\alpha 8} \delta_{\beta 8}) \hat{\epsilon}_\beta - \frac{i}{6} A(\gamma_\mu \gamma_5 \epsilon) \otimes \hat{\epsilon}_\alpha \\ &= \tilde{\mathcal{D}}_\mu(\epsilon \otimes \hat{\epsilon}_\alpha) \end{aligned} \quad (2.287)$$

$$\begin{aligned}
\delta(\psi \otimes \hat{\psi}_{i,\alpha}) &= \mathcal{D}_i(\epsilon \otimes \hat{\epsilon}_\alpha) + \frac{1}{12}(A - i\tilde{B}\gamma_5)\epsilon \otimes (i\gamma_i\hat{\epsilon})_\alpha + \frac{i}{3}\tilde{B}(\gamma_5\epsilon) \otimes (e_{i\alpha}\hat{\epsilon}_8 + 3\delta_{\alpha 8}e_{i\beta}\hat{\epsilon}_\beta) \\
&= \check{\mathcal{D}}_i(\epsilon \otimes \hat{\epsilon}_\alpha).
\end{aligned} \tag{2.288}$$

Therefore in order to preserve a supersymmetry (2.287) and (2.288) should be zero. The compatibility conditions for this is given by

$$0 = [\check{\mathcal{D}}_\mu, \check{\mathcal{D}}_\nu] \epsilon \otimes \hat{\epsilon}_\alpha \tag{2.289}$$

Taking into account that $[D_\mu, D_\nu]\epsilon = R_{\mu\nu}^{mn}\gamma_{mn}\epsilon$ we obtain from (2.289) that

$$\mathcal{R}_{\mu\nu}(\epsilon \otimes \hat{\epsilon}_\alpha) = -\frac{1}{3}g_{\mu\nu} \left[A^2(\epsilon \otimes \hat{\epsilon}_\alpha) + \frac{1}{4}\tilde{B}^2(\epsilon \otimes \hat{\epsilon}_\alpha + 48\delta_{\alpha 8}\epsilon \otimes \hat{\epsilon}_8) \right]. \tag{2.290}$$

Equation (2.290) is compatible with the Einstein equations (2.283) only if $B = 0$ and then it follows that

$$\delta(\psi_\mu \otimes \hat{\psi}_\alpha) = \mathcal{D}_\mu(\epsilon \otimes \hat{\epsilon}_\alpha) - \frac{1}{4}\sqrt{\frac{|R|}{3}}(i\gamma_\mu\gamma_5\epsilon \otimes \hat{\epsilon}_\alpha),$$

being $R = R_{\mu\nu}g^{\mu\nu}$. Taking $B = 0$ we get $R = 5A^2/7$ and that ϵ is a conformal Killing spinor, that is

$$\mathcal{D}_i\hat{\epsilon} = -\frac{1}{2}\sqrt{\frac{R}{42}}(i\gamma_i\hat{\epsilon}) = -\frac{1}{12}A(i\gamma_i\hat{\epsilon}). \tag{2.291}$$

The variation (2.291) cancels (2.288) as should be and the internal space is of weak G_2 holonomy. From (2.251) and (2.256) we get that

$$d\Phi = -\frac{2}{3}A * \Phi. \tag{2.292}$$

This mean that $B \neq 0$ is incompatible with the requirement of supersymmetry. In other words, any non zero background field F_{abcd} broke susy. In absence of fluxes, i.e, $A = 0$ we have the product of Minkowski in four dimensions by a G_2 holonomy one, in other case we have the product of the AdS_4 by a weak G_2 holonomy space.

3. Heterotic geometry without isometries

We present some properties of hyperkahler torsion (or heterotic) geometry in four dimensions that makes it even more tractable than its hyperkahler counterpart. We show that in $d = 4$ hypercomplex structures and weak torsion hyperkahler geometries are the same. We present two equivalent formalisms describing such spaces, they are stated in the propositions of section 1. The first is reduced to solve a non linear system for a doublet of potential functions, which was found first Plebanski and Finley. The second is equivalent to find the solutions of a quadratic Ashtekar-Jacobson-Smolín like system, but without a volume preserving condition. Is in this second sense that heterotic spaces are simpler than usual hyperkahler ones. We also analyze strong the strong version of this geometry. Certain examples are presented, some of them are of the Callan-Harvey-Strominger type and others are not. We discuss the benefits and disadvantages of both formulations in detail.

3.1 Introduction

Supersymmetric σ models in two dimensions with $N = 2$ supersymmetry occur on Kahler manifolds while $N = 4$ occurs on hyperkahler spaces [44]. More general supersymmetric σ models can be constructed by including Wess-Zumino-Witten type couplings in the action [95]-[48]. These couplings can be interpreted as torsion potentials and the relevant geometry is a generalization of the Kahler and hyperkahler ones with closed torsion. Such spaces are known as strong Kahler and hyperkahler torsion (HKT) geometries. The first examples of $N=(4,4)$ SUSY sigma model with torsion (and the corresponding HKT geometry) were found in [180] and developed further in [181]-[183] by using of the harmonic superspace formalism.

Hyperkahler torsion geometry is also an useful mathematical tool in order to construct heterotic string models [49]-[50]. In particular heterotic $(4,0)$ supersymmetric models are those that lead to strong hyperkahler torsion geometry [51]-[53]. If the torsion is not closed we have a weak hyperkahler geometry and this case has also physical significance [55]-[57].

A direct way of classifying the possible HKT spaces is to find the most general weak hyperkahler spaces and after that to impose the strong condition, i.e, the closure of the torsion. Callan, Harvey and Strominger noticed that under a conformal transformation any usual hyperkahler geometry is mapped into one with torsion [55]; such examples are sometimes called minimal in the literature [68] and they are not the most general [136]. As there is a particular interest in constructing spaces with at least one tri-holomorphic Killing vector due to applications related to dualities [58], there were found heterotic extensions of the known hyperkahler spaces and, in particular, the Eguchi-Hanson and Taub-Nut ones. The heterotic Eguchi-Hanson geometry was shown to be conformal to the usual one, while heterotic Taub-Nut is a new geometry [136].

One of the main properties of hyperkahler torsion spaces is the integrability of the complex structures, that is, the annulation of their Nijenhuis tensor. In four dimensions this implies that the Weyl tensor of such manifolds is self-dual [59]-[61]. The converse of this statement is not true in general. On the other hand there exist a one to one correspondence between four dimensional self-dual structures with at least one isometry and 3-dimensional Einstein-Weyl structures [157]. The Einstein-Weyl condition is a generalization of the Einstein one to include conformal transformations, and weak hyperkahler spaces should correspond with certain special Einstein-Weyl metrics. In [67]-[68] it was shown that the self-dual spaces corresponding to the round three sphere and the Berger sphere (which are Einstein-Weyl) are of heterotic type.

Arguments related to the harmonic superspace formalism suggest that indeed there are more examples [69]-[70].

The present work is related to the construction of weak heterotic geometries in $d = 4$ without Killing vectors. This problem is of interest also because any weak space with isometries should arise as subcases of those presented here. For the sake of clarity we resume the result presented in this letter in the following two equivalent propositions.

Proposition 10 *Consider a metric g defined on a manifold M together with three complex structures J^i satisfying the algebra $J^i \cdot J^j = -\delta_{ij} + \epsilon_{ijk} J^k$ and for which the metric is quaternion hermitian, i.e, $g(X, Y) = g(J^i X, J^i Y)$. Define the conformal family of metrics $[g]$ consisting of all the metrics g' related to g by an arbitrary conformal transformation.*

a) *Then we have the equivalence*

$$d\overline{J}^i + \alpha \wedge \overline{J}^i = 0 \quad \Longleftrightarrow \quad N^i(X, Y) = 0, \quad (3.293)$$

where $\overline{J}^i$ and $N^i(X, Y)$ are the Kahler form and the Niejenhuis tensor associated to J^i , d is the usual exterior derivative and α is a 1-form. If any of (3.293) hold for g , then (3.293) is also satisfied for any g' of the conformal structure $[g]$, i.e, (3.293) is conformally invariant.

b) *Any four dimensional weak hyperkahler torsion metric is equivalent to one satisfying (3.293) and there exists a local coordinate system (x, y, p, q) for which the metric take the form*

$$g = (dx - \Phi_x dp + \Phi_x dq) \otimes dp + (dy + \Psi_y dp - \Psi_x dq) \otimes dq, \quad (3.294)$$

up to a conformal transformation $g \rightarrow \omega^2 g$. The potentials Ψ and Φ satisfy the non-linear system

$$[\Phi_y \partial_x \partial_x + \Psi_x \partial_y \partial_y - (\Phi_x + \Psi_y) \partial_x \partial_y + \partial_x \partial_p + \partial_y \partial_q] \begin{pmatrix} \Phi \\ \Psi \end{pmatrix} = 0. \quad (3.295)$$

c) *Conversely any metric (3.294) defines a conformal family $[g]$ in which all the elements g' are weak hyperkahler torsion metrics. The torsion T corresponding to (3.294) is given by*

$$\begin{aligned} T = & -\Xi_x dq \wedge (dy \wedge dx + \Phi_y dp \wedge dx - \Psi_y dy \wedge dp) \\ & + \Xi_y dp \wedge (dy \wedge dx + \Phi_x dy \wedge dq - \Psi_x dq \wedge dx), \end{aligned} \quad (3.296)$$

where $\Xi = \Phi_x - \Psi_y$. Under the conformal transformation $g \rightarrow \omega^2 g$ the torsion is transformed as $T \rightarrow T + *_g 2d \log(\omega)^4$.

Proposition 10 should not be considered as a generalization of the Kahler formalism for weak HKT spaces. Although (Φ, Ψ) is a doublet potential, the metric (3.294) is not written in complex coordinates. The use of holomorphic coordinates for such spaces is described in detail in [136]. The following is an Ashtekar-Jacobson-Smolín like formulation for the same geometry.

⁴The action of the Hodge star $*_g$ is defined by

$$*_g e^a = \epsilon_{abcd} e^b \wedge e^c \wedge e^d.$$

Proposition 11 *Consider a representative $g = \delta_{ab}e^a \otimes e^b$ of a conformal family $[g]$ defined on a manifold M as at the beginning of Proposition 1, e^a being tetrad 1-forms for which the metric is diagonal.*

a) Then all the elements g' of $[g]$ will be weak hyperkahler torsion iff

$$\begin{aligned} [e_1, e_2] + [e_3, e_4] &= -A_2e_1 + A_1e_2 - A_4e_3 + A_3e_4 \\ [e_1, e_3] + [e_4, e_2] &= -A_3e_1 + A_4e_2 + A_1e_3 - A_2e_4 \\ [e_1, e_4] + [e_2, e_3] &= -A_4e_1 - A_3e_2 + A_2e_3 + A_1e_4 \end{aligned} \tag{3.297}$$

where e_a is the dual tetrad of e^a and A_i are arbitrary functions on M .

b) Conversely any solution of (3.297) defines a conformal family $[g]$ in which all the elements g' are weak hyperkahler torsion metrics. The torsion T corresponding to (3.294) is given by

$$T = *_g(A_a - c_{ab}^b)e^a \tag{3.298}$$

where c_{ab}^c are the structure functions defined by the Lie bracket $[e_a, e_b] = c_{ab}^c e_c$. The transformation of the torsion under $g \rightarrow \omega^2 g$ follows directly from Proposition 1.

To conclude, we should mention that the properties of hyperkahler torsion geometry in higher dimension were considered for instance, in [84]-[90]. In particular, the quotient construction for HKT was achieved in [86]. Also it was found that when a sigma model is coupled to gravity the resulting target metric is a generalization of quaternionic Kahler geometry including torsion [88]. This result generalizes the classical one given by Witten and Bagger [?], who originally did not include a Wess-Zumino term to the action. A generalization of the Swann extension for quaternion Kahler torsion spaces was achieved [85]. To the knowledge of the authors the harmonic superspace description of quaternion Kahler geometry was already obtained in [184], [185] but the extension to the torsion case is still an open problem.

The present part is organized as follows. In section 3.2 we define what is weak and strong heterotic geometry. We show that the problem to finding weak examples is equivalent to solving a conformal extension of the hyperkahler condition (namely, the first (3.293) given above) together with the integrability condition for the complex structure. In section 3.3 we present the consequences of (3.293) following a work of Plebanski and Finley [61]. We found out that integrability and the conformal extension of the hyperkahler condition are equivalent in $d = 4$, which is the point a) of Proposition 10. Therefore weak heterotic geometry and hypercomplex structures are exactly the same concept. We also show that strong representatives of a given heterotic structure are determined by a conformal factor satisfying an inhomogeneous Laplace equation. Just in the case when the structure contains an hyperkahler metric it is possible to eliminate the inhomogeneous part by a conformal transformation. We discuss our results in the conclusions, together with possible applications. For completeness, we show in the appendix how this geometry arise in the context of supersymmetric sigma models.

3.2 Hyperkahler torsion manifolds

3.2.1 Main properties

In this section we define what is hyperkahler torsion geometry (for an explanation of its physical meaning see Appendix). We deal with $4n$ -dimensional Riemannian manifolds M with metric

$$g = \delta_{ab} e^a \otimes e^b, \quad (3.299)$$

being e^a a tetrad basis for which the metric is diagonal, which is defined up to an $SO(4n)$ rotation. Consider the $4n \times 4n$ matrices

$$J^1 = \begin{pmatrix} 0 & -I_{n \times n} & 0 & 0 \\ I_{n \times n} & 0 & 0 & 0 \\ 0 & 0 & 0 & -I_{n \times n} \\ 0 & 0 & I_{n \times n} & 0 \end{pmatrix}, \quad J^2 = \begin{pmatrix} 0 & 0 & -I_{n \times n} & 0 \\ 0 & 0 & 0 & I_{n \times n} \\ I_{n \times n} & 0 & 0 & 0 \\ 0 & -I_{n \times n} & 0 & 0 \end{pmatrix}$$

$$J^3 = J^1 J^2 = \begin{pmatrix} 0 & 0 & 0 & -I_{n \times n} \\ 0 & 0 & -I_{n \times n} & 0 \\ 0 & I_{n \times n} & 0 & 0 \\ I_{n \times n} & 0 & 0 & 0 \end{pmatrix}. \quad (3.300)$$

It is direct to check that (3.300) satisfy the quaternion algebra

$$J^i \cdot J^j = -\delta_{ij} + \epsilon_{ijk} J^k. \quad (3.301)$$

In particular from (3.301) it is seen that $J^i \cdot J^i = -I$, a property that resemble the imaginary condition $i^2 = -1$. Associated to (3.300) we define the $(1, 1)$ tensors

$$J^i = (J^i)^b_a e_b \otimes e^a \quad (3.302)$$

which are known as almost complex structures. Here e_a the dual of the 1-form e^a . Let us introduce the triplet $\overline{J}^i$ of $(0, 2)$ tensors by

$$\overline{J}^i(X, Y) = g(X, J^i Y). \quad (3.303)$$

It is direct to check that the metric (3.299) is quaternion hermitian with respect to any of the complex structures (3.302), that is

$$g(J^i X, Y) = -g(X, J^i Y) \quad (3.304)$$

for any X, Y in $T_x M$ (in this notation $J^i X$ denotes the contraction of J^i with X). By virtue of (3.304) the tensors (3.303) are skew-symmetric and define locally a triplet of 2-forms which in the vielbein basis take the form

$$\overline{J}^i = (\overline{J}^i)_{ab} e^a \wedge e^b. \quad (3.305)$$

The 2-forms (3.305) are known as the hyperkahler triplet.

Definition

Heterotic geometry (or torsion hyperkahler geometry) is defined by the following requirements.

(1) *Hypercomplex condition.* The almost complex structures (3.310) should be integrable. This is equivalent to say the Niejenhuis tensor

$$N^i(X, Y) = [X, Y] + J^i[X, J^i Y] + J^i[J^i X, Y] - [J^i X, J^i Y] \quad (3.306)$$

associated to the complex structure J^i is zero for every pair of vector fields X and Y in TM_x .

(2) *Existence of a torsion.* There exists a torsion tensor $T_{\nu\alpha}^\mu$ defined in terms of the metric for which the $T_{\mu\nu\alpha} = g_{\mu\xi} T_{\nu\alpha}^\xi$ is fully skew-symmetric and define a three form (we write it in the tetrad basis)

$$T = \frac{1}{3!} T_{\mu\nu\lambda} dx^\mu \wedge dx^\nu \wedge dx^\lambda. \quad (3.307)$$

Here greek indices denote the quantities related to a coordinate basis $x^{\mu 5}$. If (3.307) is closed, i.e, $dT = 0$, the geometry will be called strong, in other case weak.

(3) *Covariant constancy of J^i .* Let us define a derivative D_μ with the connection

$$\Upsilon_{\mu\nu}^\rho = \Gamma_{\mu\nu}^\rho - \frac{1}{2} T_{\mu\nu}^\rho, \quad (3.308)$$

being $\Gamma_{\mu\nu}^\rho$ the Christoffel symbols associated to the usual Levi-Civita connection. By definition the structures $(J^i)_a^b$ of an hyperkahler torsion space satisfy

$$D_\mu (J^i)_\nu^\rho = 0 \quad (3.309)$$

that is, they are covariantly constant with respect to D_μ .

We are concerned through this work with $d = 4$. In this case the explicit form of the (1, 1) tensors (3.302) is obtained by putting $n = 1$ into (3.300), the result is

$$\begin{aligned} J^1 &= -e_1 \otimes e^2 + e_2 \otimes e^1 - e_3 \otimes e^4 + e_4 \otimes e^3 \\ J^2 &= -e_1 \otimes e^3 + e_3 \otimes e^1 - e_4 \otimes e^2 + e_2 \otimes e^4 \\ J^3 &= -e_1 \otimes e^4 + e_4 \otimes e^1 - e_2 \otimes e^3 + e_3 \otimes e^2, \end{aligned} \quad (3.310)$$

and the corresponding hyperkahler triplet (3.305) is

$$\begin{aligned} \overline{J}^1 &= e^2 \wedge e^1 + e^4 \wedge e^3 \\ \overline{J}^2 &= e^3 \wedge e^1 + e^2 \wedge e^4 \\ \overline{J}^3 &= e^4 \wedge e^1 + e^3 \wedge e^2, \end{aligned} \quad (3.311)$$

Matrices (3.300) with $n = 1$ are a non unique 4×4 representation the algebra (3.301). In fact due to the $SO(4)$ freedom to select e^a it follows that (3.310) are defined up to an $SO(4)$ rotation. The multiplication rule (3.301) is unchanged under such rotation. It can also be checked from

⁵In the vienbein basis relation (3.307) will be

$$T = \frac{1}{3!} T_{abc} e^a \wedge e^b \wedge e^c.$$

(3.304) that the metric $g(X, Y)$ is quaternion hermitian with respect to any $SO(4)$ rotated complex structures.

As we have seen condition $N^i(X, Y) = 0$ for every J^i in (3.310) is equivalent to solve the quadratic system (3.297) which is invariant under a transformation

$$e_a \rightarrow \omega e_a \ (g \rightarrow \omega^2 g), \quad A_a \rightarrow A_a + e_a \log \omega.$$

This mean that weak hyperkahler spaces are some special kind of hypercomplex structures. In the limit $A_i = 0$ (3.297) reduce to a system equivalent to the Ashtekar-Jacobson-Smolín one [153] for hyperkahler spaces. Nevertheless one can not conclude that (3.297) always describe spaces which are conformally equivalent to hyperkahler ones, even in the case $A_a = 0$.⁶ In the special cases in which the resulting family is conformal hyperkahler we will call it of the Callan-Harvey-Strominger type [55].

It is important to recall that although the complex structures are defined up to certain $SO(4)$ automorphisms of (3.301), this does not affect the integrability condition $N^i(X, Y) = 0$. A $SO(4)$ rotation of the complex structures can be compensated by an $SO(4)$ rotation of the frame leaving (2.39) and therefore $N^a(e_i, e_j) = 0$ invariant. By the results that we will present below it will be clear that such rotations also do not affect conditions 2 and 3.

3.2.2 Relation with the Plebanski-Finley conformal structures

The requirements 1, 2 and 3 of the definition of HKT geometry are not independent because the last two implies the first. To see this clearly let us write the expression for the Nijenhuis tensor in a coordinate basis x^μ ,

$$N_{\mu\nu}^\rho = (J^i)_{\mu}^{\lambda} [\partial_{\lambda} (J^i)_{\nu}^{\rho} - \partial_{\nu} (J^i)_{\lambda}^{\rho}] - (\mu \leftrightarrow \nu) = 0. \quad (3.312)$$

Then (3.312) together with (3.309) implies that

$$T_{\rho\mu\nu} - (J^i)_{[\mu}^{\lambda} (J^i)_{\nu}^{\sigma} T_{\rho]\lambda\sigma} = 0. \quad (3.313)$$

It is simple to check that (3.313) is satisfied for any skew-symmetric torsion. Let us consider the tetrad basis e^a for which the complex structures take the form (3.300), then (3.313) is a direct consequence of the skew symmetry of $T_{\mu\nu\alpha}$. Since $N^i(X, Y)$ is a tensor then it is zero in any basis. Thus we have proved that the requirements 2 and 3 imply first.

We will show now that the task to find a weak HKT geometry is in fact equivalent to obtain the solutions of

$$d\bar{J}^i + \alpha \wedge \bar{J}^i = 0 \quad (3.314)$$

where α is an arbitrary 1-form. To prove this we note that the relation

$$2D_{\mu}g_{\nu\alpha} = -g_{\mu\xi}T_{\nu\alpha}^{\xi} - g_{\nu\xi}T_{\mu\alpha}^{\xi}$$

together with the skew symmetry of $T_{\mu\nu\alpha} = g_{\mu\xi}T_{\nu\alpha}^{\xi}$ yields directly that

$$D_{\mu}g_{\nu\alpha} = 0. \quad (3.315)$$

⁶In order to have an hyperkahler manifold the tetrad should also preserve certain volume form. If this condition hold the solutions are also called minimal.

Equation (3.315) implies the equivalence

$$D_\mu(J^i)_\alpha^\nu = 0 \quad \Longleftrightarrow \quad D_\mu(\bar{J}^i)_{\nu\alpha} = 0, \quad (3.316)$$

and therefore for an hyperkahler torsion space

$$D_\mu(\bar{J}^i)_{\nu\alpha} = 0.$$

From the identity [170]

$$D_\mu(\bar{J}^i)_{\nu\alpha} = D_{[\mu}(\bar{J}^i)_{\nu\alpha]} - 3D_{[\mu}(\bar{J}^i)_{\xi\rho]}(J^i)_\nu^\xi(J^i)_\alpha^\rho + (\bar{J}^i)_{\mu\xi}(N^i)_{\nu\alpha}^\xi,$$

is easily seen that

$$D_\mu(\bar{J}^i)_{\nu\alpha} = 0 \quad \Longleftrightarrow \quad D_{[\mu}(\bar{J}^i)_{\nu\alpha]} = 0, \quad N^i(X, Y) = 0. \quad (3.317)$$

where the brackets denote the totally antisymmetric combination of indices. But we have seen that $N^i(X, Y) = 0$ is satisfied by our hypotesis. An explicit calculation using (3.309),(3.308) and the skew-symmetry of $T_{\mu\nu\alpha}$ shows that

$$D_{[\mu}(\bar{J}^i)_{\nu\alpha]} = 0 \quad \Longleftrightarrow \quad d\bar{J}^i + \frac{1}{2}T_{dbc}(J^i)_a^d e^a \wedge e^b \wedge e^c = 0. \quad (3.318)$$

The second (3.318) is the generalization of the hyperkahler condition for torsion manifolds. If the torsion $T_{\mu\nu\alpha}$ is zero, then the second (3.318) implies that the hyperkahler triplet is closed, which is a well known feature of hyperkahler geometry. Consider now the explicit form of (3.318) for $\bar{J}^1$. Then (3.309) and the first (3.300) gives us that

$$\begin{aligned} d\bar{J}^1 + T_{234}e^1 \wedge e^3 \wedge e^4 - T_{134}e^2 \wedge e^3 \wedge e^4 + T_{312}e^4 \wedge e^1 \wedge e^2 - T_{412}e^3 \wedge e^1 \wedge e^2 \\ = d\bar{J}^1 + (T_{234}e^1 + T_{134}e^2 - T_{123}e^4 + T_{124}e^3) \wedge (e^1 \wedge e^2 + e^3 \wedge e^4) = 0 \end{aligned}$$

and therefore

$$d\bar{J}^1 + (*_g T) \wedge \bar{J}^1 = 0.$$

The same is obtained for $\bar{J}^2$ and $\bar{J}^3$, namely

$$d\bar{J}^i + (*_g T) \wedge \bar{J}^i = 0. \quad (3.319)$$

Let us define now the 1-form α given by

$$\alpha = *_g T \quad \Longleftrightarrow \quad *_g \alpha = T. \quad (3.320)$$

Then we have from (3.319) that (3.318) is equivalent to

$$d\bar{J}^i + \alpha \wedge \bar{J}^i = 0. \quad (3.321)$$

Therefore formulas (3.316)-(3.318) implies that hyperkahler torsion geometry is completely characterized by (3.314) which is what we wanted to show.

Condition (3.321) is the conformally invariant extension of the hyperkahler one and has been studied by Plebanski and Finley in the seventies [61]. A detailed calculation shows that (3.321) is invariant under

$$e^a \rightarrow \omega e^a \quad (g \rightarrow \omega^2 g) \quad \alpha \longrightarrow \alpha + 2d \log \omega, \quad (3.322)$$

in consequence it defines a conformal family $[g]$. The transformation law (3.322) implies in particular that if α is a gradient there exist a representative of $[g]$ for which

$$\alpha = 0 \quad \Longleftrightarrow \quad d\overline{\mathcal{J}}^i = 0.$$

Clearly such families will be conformal to an hyperkahler space and is of the Callan-Harvey-Strominger type. The main result of Plebanski and Finley is that in general (3.321) implies that the Weyl tensor of $[g]$ is self-dual [61]. From (3.320) we deduce that under $g \rightarrow \omega^2 g$ it follows that

$$T \longrightarrow T + *_g 2d \log(\omega), \quad (3.323)$$

which is a transformation found by Callan et all in [55].

Some more comments are in order. As we have seen the requirements 2 and 3 imply that $N^i(X, Y) = 0$ and also imply (3.321). Therefore we have

$$N^i(X, Y) = 0 \quad \Longleftarrow \quad d\overline{\mathcal{J}}^i + \alpha \wedge \overline{\mathcal{J}}^i = 0. \quad (3.324)$$

But is also known that [59]-[60]

$$N^i(X, Y) = 0 \quad \Longrightarrow \quad d\overline{\mathcal{J}}^i + \alpha \wedge \overline{\mathcal{J}}^i = 0. \quad (3.325)$$

To see (3.325) consider the connection ω given by the first structure equation

$$de^a + \omega_b^a \wedge e^b = 0.$$

It is well known that the antisymmetric part $\omega_{[bc]}^a$ is related to the structure functions defined by the Lie bracket $[e_a, e_b] = c_{ab}^c e_c$ by

$$\omega_{[bc]}^a = \frac{1}{2} c_{bc}^a.$$

Consider now the form

$$\alpha = A - \chi$$

where

$$A = A_a e^a, \quad \chi = c_{ab}^b e^a. \quad (3.326)$$

and the self-dual form $\overline{\mathcal{J}}^1 = e^1 \wedge e^2 + e^3 \wedge e^4$. Then by use of (3.297) one obtain that

$$\begin{aligned} d\overline{\mathcal{J}}^1 &= d(e^1 \wedge e^2 + e^3 \wedge e^4) \\ &= -\frac{1}{2} c_{ab}^{[1} e^{2]} \wedge e^a \wedge e^b - \frac{1}{2} c_{ab}^{[3} e^{4]} \wedge e^a \wedge e^b \\ &= e^1 \wedge e^2 \wedge (c_{ab}^a e^b + A_3 e^3 + A_4 e^4) + e^3 \wedge e^4 \wedge (c_{ab}^a e^b + A_1 e^1 + A_2 e^2) \\ &= (e^1 \wedge e^2 + e^3 \wedge e^4) \wedge (A - \chi) \end{aligned}$$

and therefore

$$d\overline{\mathcal{J}}^1 + \overline{\mathcal{J}}^1 \wedge (\chi - A) = 0.$$

This is equivalent to (3.335) with $\alpha = \chi - A$. The same formula holds for $\overline{\mathcal{J}}^2$ and $\overline{\mathcal{J}}^3$. Then we conclude that

$$N^i(X, Y) = 0 \quad \Longleftrightarrow \quad d\overline{\mathcal{J}}^i + \alpha \wedge \overline{\mathcal{J}}^i = 0, \quad (3.327)$$

that is, hypercomplex structures are *the same* as the conformal structures defined by (3.321).

3.3 The general solution

We have seen in the previous section that to find an hyperkahler torsion geometry is equivalent to find a metric which solve the conditions stated in (3.314). This task was solved by Plebanski and Finley and is stated in the following proposition [61].

Proposition 12 *For any four dimensional metric g for which*

$$d\bar{J}^i + \alpha \wedge \bar{J}^i = 0$$

hold there exist a local coordinate system (x, y, p, q) for which g take the form

$$g = (dx - \Phi_x dp + \Phi_x dq) \otimes dp + (dy + \Psi_y dp - \Psi_x dq) \otimes dq, \quad (3.328)$$

up to a conformal transformation $g \rightarrow \omega^2 g$. The functions Ψ and Φ depends (x, y, p, q) and satisfy the non-linear system

$$[\Phi_y \partial_x \partial_x + \Psi_x \partial_y \partial_y - (\Phi_x + \Psi_y) \partial_x \partial_y + \partial_x \partial_p + \partial_y \partial_q] \begin{pmatrix} \Phi \\ \Psi \end{pmatrix} = 0. \quad (3.329)$$

The converse of this assertion is also true.

For completeness we intend to give the proof with certain detail, following the original reference [61]. It is convenient to introduce the Penrose notation and write the metric g in the following complex form

$$g = \delta_{ab} e^a \otimes e^b = E^1 \otimes E^2 + E^3 \otimes E^4. \quad (3.330)$$

We have defined the new complex tetrad

$$\begin{aligned} E^1 &= \frac{1}{\sqrt{2}}(e^1 + ie^2), & E^2 &= \frac{1}{\sqrt{2}}(e^1 - ie^2) \\ E^3 &= \frac{1}{\sqrt{2}}(e^3 + ie^4), & E^4 &= \frac{1}{\sqrt{2}}(e^3 - ie^4). \end{aligned} \quad (3.331)$$

In term of this basis the complex structures (3.311) are expressed as

$$\begin{aligned} \bar{J}^1 &= 2E^4 \wedge E^1, & \bar{J}^3 &= 2E^3 \wedge E^2 \\ \bar{J}^2 &= -E^1 \wedge E^2 + E^3 \wedge E^4. \end{aligned} \quad (3.332)$$

The advantage of this notation is that allow us to directly apply the Frobenius theorem.

Frobenius theorem *In n -dimensions, if in a certain domain U there exist r 1-forms β^i ($i = 1, \dots, r$) such that*

$$\Omega = \beta^1 \wedge \dots \wedge \beta^r \neq 0$$

and there exist a 1-form γ such that

$$d\Omega = \gamma \wedge \Omega, \quad (3.333)$$

then there exists some functions f_j^i and g^j on U such that

$$\beta^i = \sum_{j=1}^r f_j^i dg^j. \quad (3.334)$$

The two conditions

$$d\overline{J}^1 + \alpha \wedge \overline{J}^1 = 0, \quad d\overline{J}^3 + \alpha \wedge \overline{J}^3 = 0 \quad (3.335)$$

of (3.321) are of the form (3.333) with E^i are playing the role of the 1-forms β^i , $\overline{J}^1$ and $\overline{J}^3$ are the analog of Ω and α play the role of γ . Then Frobenius theorem implies the existence of scalar functions $\tilde{A}, \dots, \tilde{H}$ and $p, \dots, s$ such that

$$\begin{aligned} E^1 &= \tilde{A}dp + \tilde{B}dq, \\ E^2 &= \tilde{E}dr + \tilde{F}ds, \\ E^3 &= -\tilde{G}dr - \tilde{H}ds \\ E^4 &= -\tilde{C}dp - \tilde{D}dq. \end{aligned} \quad (3.336)$$

The functions $\tilde{A}, \dots, \tilde{H}$ and $p, \dots, s$ are the analog of f_j^i and g^j in (3.334) respectively.

Consider the two functions ϕ and f defined by

$$\begin{aligned} \tilde{A}\tilde{D} - \tilde{B}\tilde{C} &= (\phi^{-1}e^f)^2, \\ \tilde{E}\tilde{H} - \tilde{F}\tilde{G} &= (\phi^{-1}e^{-f})^2. \end{aligned} \quad (3.337)$$

It is more natural to use the variables

$$\begin{aligned} (A, B, C, D) &= (\phi^{-1}e^f)^{-1}(\tilde{A}, \tilde{B}, \tilde{C}, \tilde{D}) \\ (E, F, G, H) &= (\phi^{-1}e^{-f})^{-1}(\tilde{E}, \tilde{F}, \tilde{G}, \tilde{H}) \end{aligned}$$

with the functions $A, \dots, H$ normalized as

$$AD - BC = 1, \quad EH - GF = 1. \quad (3.338)$$

Using (3.337) we can express (3.336) as

$$\begin{aligned} E^1 &= \phi^{-1}e^f(Adp + Bdq), \\ E^2 &= \phi^{-1}e^{-f}(Edr + Fds), \\ E^3 &= -\phi^{-1}e^{-f}(Gdr + Hds), \\ E^4 &= -\phi^{-1}e^f(Cdp + Ddq), \end{aligned} \quad (3.339)$$

Since

$$E^1 \wedge E^2 \wedge E^3 \wedge E^4 = \phi^{-4}dp \wedge dq \wedge dr \wedge ds \neq 0$$

one can consider the functions p, q, r and s as independent coordinates.

From (3.339), (3.332) and (3.335) it is obtained the following expression for α

$$\alpha = d \log(\phi) + f_p dp + f_q dq - f_r dr - f_s ds. \quad (3.340)$$

We now consider the resting condition (3.321), namely

$$d\overline{J}^2 + \alpha \wedge \overline{J}^2 = 0. \quad (3.341)$$

By use of (3.332) together with (3.339) we find that

$$\begin{aligned}\bar{\mathcal{J}}^2 = \phi^{-4}[(AE + CG)dp \wedge dr + (AF + CH)dp \wedge ds + (BE + DG)dq \wedge dr \\ + (BF + DH)dq \wedge ds].\end{aligned}\quad (3.342)$$

A direct calculation shows that (3.341) together with (3.340) implies

$$\begin{aligned}(AE + CG)2f_s - (AF + CH)2f_r &= (AE + CG)_s - (AF + CH)_r, \\ (BE + DG)2f_s - (BF + DH)2f_r &= (BE + DG)_s - (BF + DH)_r, \\ (AE + CG)2f_q - (BE + DG)2f_p &= -(AE + CG)_q + (BE + DG)_p, \\ (AF + CH)2f_q - (BF + DH)2f_p &= -(AF + CH)_q + (BF + DH)_p.\end{aligned}\quad (3.343)$$

The first two (3.343) shows that there exist certain functions x and y such that

$$\begin{aligned}AE + CG &= e^{2f}x_r, \\ BE + DG &= e^{2f}y_r, \\ AF + CH &= e^{2f}x_s, \\ BF + DH &= e^{2f}y_s.\end{aligned}\quad (3.344)$$

By multiplying the last two (3.343) by e^{2f} and using (3.344) we also obtain that

$$(e^{4f}x_r)_q = (e^{4f}y_r)_p, \quad (e^{4f}x_s)_q = (e^{4f}y_s)_p. \quad (3.345)$$

The solution of (3.344) and (3.345) are

$$\begin{aligned}G &= A(e^{2f}y_r) - B(e^{2f}x_r) \\ H &= A(e^{2f}y_s) - B(e^{2f}x_s) \\ E &= D(e^{2f}x_r) - C(e^{2f}y_r) \\ F &= D(e^{2f}x_s) - C(e^{2f}y_s)\end{aligned}\quad (3.346)$$

The normalization (3.338) and (3.346) implies that

$$J = \begin{vmatrix} x_r & x_s \\ y_r & y_s \end{vmatrix} = e^{-4f} \neq 0,$$

and therefore x and y can be used as independent coordinates instead of r and s . Moreover, the functions $A, \dots, D$ are not determined by the equations but just constrained by the first (3.338). This reflects that the tetrad and the vielbein are defined up to an $SO(4)$. This rotations leaves condition $AD - BC = 1$ invariant. Therefore without loss of generality we can select $A = D = 1$ and $B = C = 0$. The resulting tetrad is now

$$\begin{aligned}E^1 &= (\phi e^{-f})^{-1}dp, \\ E^2 &= (\phi e^{-f})^{-1}(dx + Kdp + Ldq), \\ E^3 &= -(\phi e^{-f})^{-1}(dy + Mdp + Ndq),\end{aligned}\quad (3.347)$$

$$E^4 = -(\phi e^{-f})^{-1} dq,$$

being $K = -x_p$, $L = -x_q$, $M = -y_p$ and $N = -y_q$. The dual basis of (3.347) is

$$\begin{aligned} E_2 &= \phi e^{-f} \partial_x \\ E_1 &= \phi e^{-f} (\partial_p - K \partial_x - M \partial_y), \\ E_4 &= -\phi e^{-f} (\partial_q - L \partial_x - N \partial_y), \\ E_3 &= -\phi e^{-f} \partial_y. \end{aligned} \tag{3.348}$$

Because we are working with a conformal structure is easy to check that we can make eliminate ϕe^{-f} in (3.348) by a conformal transformation. From (3.335) it is calculated that

$$2\alpha = (L - M)_x dq - (L - M)_y dp. \tag{3.349}$$

Also from (3.341) and (3.349) it follows that

$$\begin{aligned} \partial_x K + \partial_y L &= 0, & \partial_x M + \partial_y N &= 0, \\ (\partial_p - K \partial_x - M \partial_y) L - (\partial_q - L \partial_x - N \partial_y) K &= 0, \\ (\partial_p - K \partial_x - M \partial_y) N - (\partial_q - L \partial_x - N \partial_y) M &= 0 \end{aligned} \tag{3.350}$$

The first two (3.350) implies the existence of two functions Φ and Ψ such that

$$K = -\Phi_x, \quad L = \Phi_x, \quad M = \Psi_y, \quad N = -\Psi_x, \tag{3.351}$$

and the first two (3.350) together with (3.351) gives the following non-linear equations for the doublet (Φ, Ψ)

$$[\Phi_y \partial_x \partial_x + \Psi_x \partial_y \partial_y - (\Phi_x + \Psi_y) \partial_x \partial_y + \partial_x \partial_p + \partial_y \partial_q] \begin{pmatrix} \Phi \\ \Psi \end{pmatrix} = 0. \tag{3.352}$$

Then we conclude that the most general structure satisfying (3.321) is given in terms of two key functions Φ and Ψ satisfying (3.352). A simple calculation shows that the metric corresponding to (3.348) is (3.328) which is what we wanted to prove (Q.E.D).

It will be instructive we will show that for any structure of the proposition 12 it follows that

$$N^i(X, Y) = 0,$$

in accordance with (3.324). This is completely equivalent to solve the system (3.297), that is

$$\begin{aligned} [e_1, e_2] + [e_3, e_4] &= -A_2 e_1 + A_1 e_2 - A_4 e_3 + A_3 e_4, \\ [e_1, e_3] + [e_4, e_2] &= -A_3 e_1 + A_4 e_2 + A_1 e_3 - A_2 e_4, \\ [e_1, e_4] + [e_2, e_3] &= -A_4 e_1 - A_3 e_2 + A_2 e_3 + A_1 e_4, \end{aligned}$$

for the basis e_i corresponding to (3.328). From (3.331), (3.347) and (3.351) we find that

$$e^1 = \frac{1}{\sqrt{2}} [dp + (dx - \Phi_y dp + \Phi_x dq)],$$

$$\begin{aligned}
e^2 &= \frac{1}{i\sqrt{2}}[dp - (dx - \Phi_y dp + \Phi_x dq)] \\
e^3 &= -\frac{1}{\sqrt{2}}[dq + (dy + \Psi_y dp - \Psi_x dq)], \\
e^4 &= \frac{1}{i\sqrt{2}}[dq - (dy + \Psi_y dp - \Psi_x dq)].
\end{aligned} \tag{3.353}$$

The dual basis corresponding to (3.353) is

$$\begin{aligned}
e_1 &= \frac{1}{\sqrt{2}}[(\partial_p + \Phi_y \partial_x - \Psi_y \partial_y) + \partial_x], \\
e_2 &= -\frac{1}{i\sqrt{2}}[(\partial_p + \Phi_y \partial_x - \Psi_y \partial_y) - \partial_x] \\
e_3 &= -\frac{1}{\sqrt{2}}[\partial_y + (\partial_q - \Phi_x \partial_x + \Psi_x \partial_y)], \\
e_4 &= \frac{1}{i\sqrt{2}}[\partial_y - (\partial_q - \Phi_x \partial_x + \Psi_x \partial_y)].
\end{aligned} \tag{3.354}$$

It is simple to check that the first (3.297) for (3.354) implies that $A_i = 0$. After some tedious calculation one obtain from the full system (3.297) that

$$[\Phi_y \partial_x \partial_x + \Psi_x \partial_y \partial_y - (\Phi_x + \Psi_y) \partial_x \partial_y + \partial_x \partial_p + \partial_y \partial_q] \begin{pmatrix} \Phi \\ \Psi \end{pmatrix} = 0,$$

which is exactly (3.352).

We conclude that the general form of an HKT metric is

$$g = (dx - \Phi_y dp + \Phi_x dq) \otimes dp + (dy + \Psi_y dp - \Psi_x dq) \otimes dq$$

up to a conformal transformation. The form α in (3.349) can be expressed as

$$\alpha = \Xi_x E^1 - \Xi_y E^4, \tag{3.355}$$

where $\Xi = \Phi_x - \Psi_y$. Then it is direct to prove using (3.331) that

$$*E^1 = -E^2 \wedge E^3 \wedge E^4, \quad *E^4 = -E^1 \wedge E^2 \wedge E^3. \tag{3.356}$$

Formula (3.356) together with (3.320) and (3.355) gives explicitly the torsion, namely

$$T = -\Xi_x E^2 \wedge E^3 \wedge E^4 + \Xi_y E^1 \wedge E^2 \wedge E^3. \tag{3.357}$$

All the results of this section are stated in the propositions of the introduction.

The geometries presented till now are weak heterotic. Due to conformal invariance all the elements of the conformal family $[g]$ corresponding to (3.328) are also weak. Therefore in order to find an strong geometry we should not impose $dT = 0$ for the torsion (3.296) corresponding to (3.328) but instead, we must solve $d\tilde{T} = 0$ where $\tilde{T} = T + *_g 2d \log(\omega)$ is the torsion corresponding to the representative $\tilde{g} = \omega^2 g$. After some calculation we obtain the following *linear* equation defining ω

$$\Delta_g \log(\omega) = \partial_{[1} T_{234]}, \tag{3.358}$$

where Δ_g is the Laplace operator for (3.328). The left side (3.358) does not depend on ω and acts as an inhomogeneous source. Then the spaces $\tilde{g} = \omega^2 g$ will be a strong representatives of the family if ω solve (3.358). The spaces (3.328) will be strong only in the case that constant ω is a solution of the resulting equations, which implies that

$$\partial_{[1} T_{234]} = \Xi_{xp} - (\Xi_x \Phi_y)_y + (\Xi_x \Psi_y)_x + \Xi_{yq} - (\Xi_y \Phi_x)_x + (\Xi_y \Psi_x)_y = 0. \quad (3.359)$$

It also follows from the result of this section that if we deal with a minimal weak structure, then (3.358) is reduced to

$$\Delta_{\hat{g}} \log(\omega) = 0, \quad (3.360)$$

being $\hat{g}$ the hyperkahler representative of the family [55].

3.4 Discussion

In the present work we attacked the problem of constructing weak and strong 4-dimensional hyperkahler torsion geometries without isometries. We have shown that the most general local form for a weak metric is defined by two potential functions. Weak condition is conformally invariant and the corresponding metrics are determined up to a conformal transformation.

We argue that the present work have practical applications, although the system of equations for the potentials is highly non-linear and difficult to solve. We have seen that in four dimensions hypercomplex structures and weak torsion hyperkahler geometries are totally equivalent concepts. Concretely speaking, we have shown that integrability of the complex structures implies and is implied by the other weak HKT properties. Therefore the problem to find weak heterotic geometries is reduced to find the solutions of an Ashtekar-Jacobson-Smolín like system without the volume preserving condition, which has the advantage to be quadratic.

From this discussion is clear that a weak metric will be of the Callan-Harvey-Strominger type (i.e. conformal to an hyperkahler one) if there is a volume form preserved by the corresponding tetrad. Nevertheless not all the hypercomplex structures preserves a 4-form and the metrics of references [136]-[68] should be counterexamples. We have seen also that a space is of Callan-Harvey-Strominger type if the torsion is the dual of a gradient. Therefore both conditions should be equivalent.

The strong condition (i.e. the closure of the torsion form) is not invariant under general conformal transformation. Nevertheless the problem to find the strong representatives of a weak heterotic family $[g]$ is reduced to solve a *linear* non homogeneous laplacian equation over an arbitrary element g . The homogeneous part can be taken to zero if the metric is conformal to a hyperkahler one.

At first sight it looks that the "Ashtekar like" formulation for HKT is the most convenient formulation. Instead we suggest that the other variant could be more useful in the context dualities, in which the presence of isometries is needed [58]. To make this suggestion more concrete, let us recall that self-dual spaces are described in terms of a Kahler potential satisfying the heavenly equation [76]. Although this equation is non-linear and the general solution is not known, the analysis of the Killing equations and the possible isometries allowed to classify the hyperkahler metrics with one Killing vector [75]. In particular it was found that they are divided in two types, the first described for by a monopole equation and the second by some limit of a Toda system [72]-[75]. We propose that is worthy to generalize, if is possible, the mathematical machinery to classify the Killing equations of gravitational instantons [75]-[81] to the hyperkahler torsion case. Perhaps it will possible in this way to find the classification of

heterotic geometry with one or more Killing vectors and in particular, to identify the subcases that match for duality applications. This goal has not been achieved yet, and our suggestion could be an alternative point of view of the references [68]-[70].

4. D-instanton sums for matter hypermultiplets

We calculate some non-perturbative (D-instanton) quantum corrections to the moduli space metric of several ($n > 1$) identical matter hypermultiplets for the type-IIA superstrings compactified on a Calabi-Yau threefold, near conifold singularities. We find a non-trivial deformation of the (real) $4n$ -dimensional hypermultiplet moduli space metric due to the infinite number of D-instantons, under the assumption of n tri-holomorphic commuting isometries of the metric, in the hyperkahler limit (i.e. in the absence of gravitational corrections).

4.1 Ooguri-Vafa solution

The Ooguri-Vafa (OV) solution [162] describes the D-instanton corrected moduli space metric of a single matter hypermultiplet in type-IIA superstrings compactified on a Calabi-Yau threefold of Hodge number $\dim H^{2,1} = 1$, when both $N = 2$ supergravity and UH are switched off, while five-brane instantons are suppressed. The matter hypermultiplet low-energy effective action in that limit is given by the four-dimensional $N = 2$ supersymmetric non-linear sigma-model that has the four-dimensional OV metric in its target space.

Rigid (or global) $N = 2$ supersymmetry of the non-linear sigma-model requires a hyperkahler metric in its target space [164]. The OV metric has a (toric) $U(1) \times U(1)$ isometry by construction [162]. There always exist a linear combination of two commuting abelian isometries that is tri-holomorphic, i.e. it commutes with $N = 2$ rigid supersymmetry [165].

Given any four-dimensional hyperkahler metric with a tri-holomorphic isometry ∂_t , it can always be written down in the standard (Gibbons-Hawking) form

$$g = \frac{1}{V}(dt + A)^2 + V(dx^2 + dy^2 + dz^2), \quad (4.361)$$

that is governed by linear equations,

$$\Delta V = \vec{\nabla}^2 V \equiv \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) V = 0, \quad \text{almost everywhere}, \quad (4.362)$$

and

$$\vec{\nabla} V + \vec{\nabla} \times A = 0. \quad (4.363)$$

The 1-form $A = A_1 dx + A_2 dy + A_3 dz$ is fixed by the monopole equation (4.363) in terms of the real scalar potential $V(x, y, z)$. Equation (4.362) means that the function V is harmonic (away from possible isolated singularities) in three Euclidean dimensions $\mathbf{R}^3$. The singularities are associated with the positions of D-instantons.

Given extra $U(1)$ isometry, after being rewritten in the cylindrical coordinates ($\rho = \sqrt{x^2 + y^2}$, $\theta = \arctan(y/x)$, $\eta = z$), the hyperkahler potential $V(\rho, \theta, \eta)$ becomes independent upon θ . Equation (4.361) was used by Ooguri and Vafa [162] in their analysis of the matter hypermultiplet moduli space near a conifold singularity. The conifold singularity arises in the limit of the vanishing CY period,

$$\int_{\mathcal{C}} \mathbf{O} \rightarrow 0 \quad (4.364)$$

where the CY holomorphic (nowhere vanishing) 3-form $\mathbf{O}$ is integrated over a non-trivial 3-cycle $\mathcal{C}$ of CY. The powerful singularity theory [172] can then be applied to study the universal behaviour of the hypermultiplet moduli space near the conifold limit, by resolving the singularity.

In the context of the CY compactification of type IIA superstrings, the coordinate ρ represents the ‘size’ of the CY cycle $\mathcal{C}$ or, equivalently, the action of the D-instanton originating from the Euclidean D2-brane wrapped about the cycle $\mathcal{C}$. The physical interpretation of the η coordinate is just the expectation value of another (RR-type) hypermultiplet scalar. The cycle $\mathcal{C}$ can be replaced by a sphere S^3 for our purposes, since the D2-branes only probe the overall size of $\mathcal{C}$.

The OV potential V is periodic in the RR-coordinate η since the D-brane charges are quantized [159]. We normalize the corresponding period to be 1, as in reference [162]. The Euclidean D2-branes wrapped m times around the sphere S^3 couple to the RR expectation value on S^3 and thus should produce additive contributions to V , with the factor of $\exp(2\pi i m \eta)$ each.

In the classical hyperkahler limit, when both N=2 supergravity and all the D-instanton contributions are suppressed, the potential $V(\rho, \eta)$ of a single matter hypermultiplet cannot depend upon η since there is no perturbative superstring state with a non-vanishing RR charge. Accordingly, the classical pre-potential $V(\rho)$ can only be the Green function of the two-dimensional Laplace operator, i.e.

$$V_{\text{classical}} = -\frac{1}{2\pi} \log \rho + \text{const.} \quad (4.365)$$

whose normalization is also in agreement with the appearing in [162].

The calculation given in [162] to determine the exact D-instanton contributions to the hyperkahler potential V is based on the idea [159] that the D-instantons should resolve the singularity of the classical hypermultiplet moduli space metric at $\rho = 0$. A similar situation arises in the standard (Seiberg-Witten) theory of a quantized N=2 vector multiplet (see reference [164] for a review).

Equation (4.362) formally defines the electrostatic potential V of electric charges of unit charge in the Euclidean upper half-plane (ρ, η) , $\rho > 0$, which are distributed along the axis $\rho = 0$ in each point $\eta = n \in \mathbf{Z}$, while there are no two charges at the same point [162]. A solution to (4.362) obeying all these conditions is unique,

$$V_{\text{OV}}(\rho, \eta) = \frac{1}{4\pi} \sum_{n=-\infty}^{+\infty} \left(\frac{1}{\sqrt{\rho^2 + (\eta - n)^2}} - \frac{1}{|n|} \right) + \text{const.} \quad (4.366)$$

After Poisson summation (4.366) takes the desired form of singularity resolution [162]

$$V_{\text{OV}}(\rho, \eta) = \frac{1}{4\pi} \log\left(\frac{\mu^2}{\rho^2}\right) + \sum_{m \neq 0} \frac{1}{2\pi} e^{2\pi i m \eta} K_0(2\pi |m| \rho), \quad (4.367)$$

where the modified Bessel function K_0 of the 3rd kind has been introduced,

$$K_s(z) = \frac{1}{2} \int_0^{+\infty} \frac{dt}{t^{s-1}} \exp\left[-\frac{z}{2}\left(t + \frac{1}{t}\right)\right] \quad (4.368)$$

valid for all $\text{Re } z > 0$ and $\text{Re } s > 0$, while μ is a constant (modulus).

Inserting the standard asymptotical expansion of the Bessel function K_0 near $\rho = \infty$ into (4.367) yields [162]

$$\begin{aligned} V_{\text{OV}}(\rho, \eta) &= \frac{1}{4\pi} \log\left(\frac{\mu^2}{\rho^2}\right) + \sum_{m=1}^{\infty} \exp(-2\pi m \rho) \cos(2\pi m \eta) \\ &\quad \times \sum_{n=0}^{\infty} \frac{\Gamma(n + \frac{1}{2})}{\sqrt{\pi} n! \Gamma(-n + \frac{1}{2})} \left(\frac{1}{4\pi m \rho}\right)^{n+\frac{1}{2}} \end{aligned} \quad (4.369)$$

The string coupling constant g can be easily reintroduced into (4.369) by a substitution $\rho \rightarrow \rho/g$. The factors of $\exp(-2\pi m\rho/g)$ in (4.369) are the contributions due to the multiple D-instantons [162].

The OV potential (4.366) is given by a (regularized) T-sum over the T-duality transformations, $\eta \rightarrow \eta + 1$, being applied to the fundamental solution $V_0 \equiv \frac{1}{4\pi r} \equiv \frac{1}{4\pi\sqrt{\rho^2+\eta^2}}$ of (4.362),

$$V_{\text{OV}}(\rho, \eta) = A + \sum_{\text{T}} V_0(\rho, \eta) = A + \sum_{\text{T}} \frac{1}{4\pi\sqrt{\rho^2 + \eta^2}} \quad (4.370)$$

where A is a constant. The fundamental solution $V_0(\rho, \eta)$ is just the Green function of the three-dimensional Laplace operator Δ in (4.362).

4.2 Pedersen-Poon Ansatz

The natural generalization of the Gibbons-Hawking Ansatz (4.361) to higher-dimensional toric hyperkahler spaces are the *Pedersen-Poon* (PP) metrics (2.160). Namely, given n commuting tri-holomorphic isometries of a (real) $4n$ -dimensional hyperkahler space, there exists a coordinate system (x_a^i, t_i) , with $i, j = 1, 2, \dots, n$ and $a, b, c = 1, 2, 3$, where the hyperkahler metric takes the form [146]

$$g = U_{ij} dx^i \cdot dx^j + U^{ij} (dt_i + A_i)(dt_j + A_j) \quad (4.371)$$

Here the dot means summation over the a -type indices, all metric components are supposed to be independent upon all t_i (thus reflecting the existence of n isometries), $U^{ij} = (U_{ij})^{-1}$, while $U_i = (U_{i1}, \dots, U_{in})$ and A_i are to be solutions to the generalized monopole (or BPS) equations

$$\begin{aligned} F_{x_\mu^i x_\nu^j} &= \epsilon_{\mu\nu\lambda} \nabla_{x_\lambda^i} U_j, \\ \nabla_{x_\lambda^i} U_j &= \nabla_{x_\lambda^j} U_i, \\ U_i &= (U_{i1}, \dots, U_{in}), \end{aligned} \quad (4.372)$$

where the field strength F of the gauge fields A has been introduced. The PP metric (11) can be completely specified by its real PP-potential $P(x, w, \bar{w})$ that generically depends upon $3n$ variables,

$$x^j = x_3^j, \quad w^j = \frac{x_1^j + ix_2^j}{2}, \quad \bar{w}^j = \frac{x_1^j - ix_2^j}{2}, \quad (4.373)$$

because the BPS equations (4.372) allow a solution [23]

$$U_{ij} = P_{x^i x^j} \quad \text{and} \quad A_j = i \left(P_{w^k x^j} dw^k - P_{\bar{w}^k x^j} d\bar{w}^k \right), \quad (4.374)$$

provided P itself obeys (almost everywhere) a linear Laplace-like equation [146]

$$P_{x^i x^j} + P_{w^i \bar{w}^j} = 0. \quad (4.375)$$

The subscripts of P denote partial differentiation with respect to the given variables.

Equation (4.375) is apparently the multi-dimensional hyperkahler generalization of (4.362). The existence of the PP-potential F essentially amounts to integrability (or linearization) of the BPS equations (4.372), because the master equation (4.375) is linear, whose solutions are not difficult to find. For instance, a general solution to (4.375) can be written down as the contour (C) integral of an arbitrary potential $G(\eta^j(\xi), \xi)$ [146],

$$P = \text{Re} \oint_C \frac{dx}{2\pi i \xi} G(\eta^j(\xi), \xi), \quad \text{where} \quad \eta^j(\xi) = \bar{w}^j + x^j \xi - w^j \xi^2. \quad (4.376)$$

4.3 D-instanton sums

Our technical assumptions are essentially the same as in reference [162], namely,

- periodicity in all x^i with period 1, due to the D-brane charge quantization,
- the classical potential F near a CY conifold singularity should have a logarithmic behaviour (elliptic fibration), being independent upon all x^i (when all $w^i \rightarrow \infty$),
- the classical singularity of the metric should be removable, i.e. an exact metric should be complete (or non-singular),
- the metric should be symmetric under the permutation group of n sets of hypermultiplet coordinates $(x^j, w^j, \bar{w}^j)$, where $j = 1, 2, \dots, n$.

The last assumption means that we only consider *identical* matter hypermultiplets. Together with our main assumption about n tri-holomorphic isometries (see e.g., our Abstract and sect. 3), it is going to lead us to an explicit solution.

We first confine ourselves to the case of *two* identical matter hypermultiplets ($n = 2$), and then quote our result for an arbitrary $n > 2$. The problem amounts to finding a solution to the master equations (4.375) subject to the conditions above. The known OV solution $V_{\text{OV}}(w, x) \equiv V_{\text{OV}}(x, w)$, defined by equation (4.366) or (4.367), can be used to introduce another function $P(x, w)$ as a solution to two equations

$$P_{xx} = -P_{w\bar{w}} = V_{\text{OV}}(x, w). \quad (4.377)$$

We do not need an explicit solution to equation (4.377) in what follows. Since equations (4.375) are linear, the superposition principle applies to their solutions. It is now straightforward to verify that there exist a ‘trivial’ solution to equations (4.375), given by a PP potential

$$P_0 = c_0 [P(x_1, w_1) + P(x_2, w_2)] \quad (4.378)$$

where c_0 is a real constant. A particular non-trivial solution in the $n = 2$ case is given by

$$P_{\text{mixed}} = c_+ P(x_1 + x_2, w_1 + w_2) + c_- P(x_1 - x_2, w_1 - w_2) \quad (4.379)$$

where $c_{\pm}$ are two other real constants. Though the linear partial differential equation (4.375) has many other solutions, we argue at the end of this section that the most general solution, satisfying all our requirements, is actually given by

$$P = P_0 + P_{\text{mixed}}. \quad (4.380)$$

To this end, we continue with the particular solution (4.380). To write down our result for the hyperkahler moduli space metric of two matter hypermultiplets, we merely need the second derivatives of the PP potential, because of (4.374). Using equations (4.366), (4.374), (4.377), (4.378), (4.379) and (4.380), we find

$$4\pi U_{11} = c_+ \sum_{n=-\infty}^{+\infty} \left(\frac{1}{\sqrt{(x_1 + x_2 - n)^2 + (w_1 + w_2)(\bar{w}_1 + \bar{w}_2)/\lambda^2}} - \frac{1}{n} \right)$$

$$\begin{aligned}
& +c_- \sum_{n=-\infty}^{+\infty} \left(\frac{1}{\sqrt{(x_1 - x_2 - n)^2 + (w_1 - w_2)(\bar{w}_1 - \bar{w}_2)/\lambda^2}} - \frac{1}{n} \right) \\
& +c_0 \sum_{n=-\infty}^{+\infty} \left(\frac{1}{\sqrt{(x_1 - n)^2 + w_1 \bar{w}_1/\lambda^2}} - \frac{1}{n} \right)
\end{aligned} \tag{4.381}$$

with a modulus parameter λ . The component U_{22} has the the form as that of (4.381), but the indices 1 and 2 have to be exchanged. The remaining two components of the matrix U in (4.371) are given by

$$\begin{aligned}
4\pi U_{12} = 4\pi U_{21} = c_+ \sum_{n=-\infty}^{+\infty} \left(\frac{1}{\sqrt{(x_1 + x_2 - n)^2 + (w_1 + w_2)(\bar{w}_1 + \bar{w}_2)/\lambda^2}} - \frac{1}{n} \right) \\
+ c_- \sum_{n=-\infty}^{+\infty} \left(\frac{1}{\sqrt{(x_1 - x_2 - n)^2 + (w_1 - w_2)(\bar{w}_1 - \bar{w}_2)/\lambda^2}} - \frac{1}{n} \right).
\end{aligned} \tag{4.382}$$

The A -field components in the PP metric (4.371) are given by the second (4.374).

The physical interpretation of our solution as the infinite D-instanton sum is evident after rewriting it as an asymptotical expansion like that of (4.369), by using (4.367) and (4.368), now being applied to the two-hypermultiplet metric components (4.381) and (4.382). In particular, their classical behavior is given by

$$U_{11} \sim \frac{c_+}{4\pi} \ln \frac{\lambda^2}{|w_1 + w_2|^2} + \frac{c_-}{4\pi} \ln \frac{\lambda^2}{|w_1 - w_2|^2} + \frac{c_0}{4\pi} \ln \frac{\lambda^2}{|w_1|^2} \tag{4.383}$$

and similarly for U_{22} after exchanging the indices 1 and 2, and U_{12} after dropping the last term in (4.383).

As regards the case of an arbitrary $n > 2$, the easiest picture arises in terms of the PP potential P — see sect. 3. We define this potential as a linear combination of $P(x_1 + x_2 + \dots + x_n, w_1 + w_2 + \dots + w_n)$, $P(-x_1 + x_2 + \dots + x_n, -w_1 + w_2 + \dots + w_n)$, $\dots$, $P(-x_1 - x_2 + \dots - x_n, -w_1 - w_2 - \dots - w_n)$, $P(x_1, w_1)$, $P(x_2, w_2)$, $\dots$ and $P(x_n, w_n)$, each one being constructed out of the OV potential, as in (4.377). Next, we define the metric components of the matrix U_{ij} in (4.371) by the second derivatives of the total PP potential F , as in (4.374). The rest is fully straightforward, *cf.* equations (4.381) and (4.382).

Finally, some comments about the uniqueness of our solution (4.380) are in order. First, we notice that all functions U^{ij} also obey (4.375) by our construction (4.374). Given any other non-trivial solution, it should lead (after Poisson resummation) to *the same* classical behaviour (4.383), while it should also reduce to the OV solution (sect. 2) for a single hypermultiplet. According to the Poisson resummation formula,

$$\sum_{n=0}^{\infty} f(n) = \sum_{n=0}^{\infty} \tilde{f}(n) , \quad \text{where} \quad \tilde{f}(u) = \int_{-\infty}^{+\infty} dv f(v) e^{2\pi i uv} \tag{4.384}$$

the logarithmic terms can only come from the $n = 0$ term in the sum, e.g. when using (4.366), choosing $n = v$ in the Fourier integral (4.384), and then taking $u = 0$ after the Fourier transform.

Hence, a difference between ours and any other solution to the potentials F or U^{ij} cannot contribute to equation (4.383). Since equation (4.383) is x -independent, we first investigated the class of x -independent real potentials F , without any reference to the metric (4.371). By

solving (4.375) in this case we found an unique solution subject to our symmetries, namely, the one given by (4.383). Hence, the anticipated uniqueness of our solution boils down to the uniqueness of the periodic (of period 1) extension of a given w -dependent solution to a solution depending upon both w and x , and having a generic form $\sum_n G(x + n, w)$ with some basic function G . It seems to be quite plausible that the square root in (4.381) is the *only* basic function to yield the logarithmic terms out of the $n = 0$ term in the D-instanton sum.

4.4 Discussion

In our final results (4.381) and (4.382) the quantum corrections to the classical (logarithmic) terms (4.383) are exponentially suppressed by the D-instanton factors (in the semiclassical description). Unlike the case of a single matter hypermultiplet, the multi-hypermultiplet moduli space metric is also sensitive to the phases of w_i (i.e. not only to their absolute values). There are no perturbative (type IIA superstring) corrections, while all D-instanton numbers contribute to the solution.

Our results can also be applied to an explicit construction of *quaternionic* metrics in real $4(n - 1)$ dimensions out of known hyperkahler metrics in real $4n$ dimensions, which is of particular importance to a description of D-instantons in N=2 supergravity [22]. For instance, the four-dimensional quaternionic manifolds with toric isometry can be fully classified [132] in terms of the PP-potential F obeying equation (4.375).

It would be also interesting to connect our results to perturbative superstring calculations in the D-instanton background, which is still a largely unsolved problem.

5. Hyperkahler spaces with local triholomorphic $U(1) \times U(1)$ isometry and supergravity solutions

In this section some examples of toric hyperkahler metrics in eight dimensions are constructed. The Swann construction of hyperkahler metrics is applied to the Calderbank Pedersen spaces in order to find hyperkahler examples with $U(1) \times U(1)$ isometry. The connection with the Pedersen-Poon toric hyperkahler metrics is explained by finding the momentum map coordinate system. This hyperkahler examples are lifted to solutions of the D=11 supergravity with and without fluxes. Type IIA and IIB backgrounds are found by use of dualities.

5.1 Toric hyperkahler metrics of the Swann type

The Swann metric (2.184) is in general quaternion Kahler. In the special case with $b = 0$ corresponds to an hyperkahler metric with explicit form

$$g = |u|^2 \bar{g} + [(du_0 - x_i \omega_-^i)^2 + (du_i + u_0 \omega_-^i + \epsilon_{ijk} u_k \omega_-^k)^2]. \quad (5.385)$$

By use of (5.385) it is direct to extend the Calderbank-Pedersen metrics (2.126) to an hyperkahler one. The quaternionic base is (2.126) and the corresponding ω_-^i are [132]

$$\omega_-^1 = \frac{1}{F} \left(\left(\frac{1}{2} F + \rho F_\rho \right) \frac{d\eta}{\rho} - \rho F_\eta \frac{d\eta}{\rho} \right), \quad \omega_-^2 = -\frac{\alpha}{F}, \quad \omega_-^3 = \frac{\beta}{F}. \quad (5.386)$$

The resulting expression for the hyperkahler metric is

$$g = [(du_0 - x_i \omega_-^i)^2 + (du_i + u_0 \omega_-^i + \epsilon_{ijk} u_k \omega_-^k)^2] + |u|^2 \left[\frac{F^2 - 4\rho^2(F_\rho^2 + F_\eta^2)}{4F^2} \left(\frac{d\rho^2 + d\eta^2}{\rho^2} \right) + \frac{[(F - 2\rho F_\rho)\alpha - 2\rho F_\eta \beta]^2 + [(F + 2\rho F_\rho)\beta - 2\rho F_\eta \alpha]^2}{F^2[F^2 - 4\rho^2(F_\rho^2 + F_\eta^2)]} \right]. \quad (5.387)$$

The Calderbank-Pedersen metrics (2.126) have two commuting Killing vectors $\partial/\partial\theta$ and $\partial/\partial\varphi$. Clearly this are also Killing vectors of the spaces (5.387).

The Kahler triplet corresponding to (2.126) is

$$\begin{aligned} \bar{J}^1 &= \frac{1}{F^2} (F^2 - 4\rho^2(F_\rho^2 + F_\eta^2)) \left(\frac{d\rho \wedge d\eta}{\rho^2} + \alpha \wedge \beta \right), \\ \bar{J}^2 &= \frac{1}{F^2} (\rho F_\eta \beta + (\rho F_\rho - \frac{1}{2} F) \alpha) \wedge \frac{d\rho}{\rho}, \\ \bar{J}^3 &= \frac{1}{F^2} (\rho F_\eta \alpha - (\rho F_\rho + \frac{1}{2} F) \beta) \wedge \frac{d\eta}{\rho}. \end{aligned} \quad (5.388)$$

Then the hyperkahler triplet Ω^i of (5.385) can be expressed in compact form as

$$\Omega = u \bar{J} \bar{u} + (du + \omega_- u) \wedge \overline{(du + \omega_- u)}, \quad (5.389)$$

where we have defined

$$\bar{J} = \bar{J}^1 I + \bar{J}^2 J + \bar{J}^3 K, \quad \Omega = \Omega^1 I + \Omega^2 J + \Omega^3 K$$

$$\omega_- = \omega_-^1 I + \omega_-^2 J + \omega_-^3 K.$$

A direct calculation using (5.388) show us that

$$\mathcal{L}_{\frac{\partial}{\partial\varphi}}\Omega^i = 0, \quad \mathcal{L}_{\frac{\partial}{\partial\theta}}\Omega^i = 0. \quad (5.390)$$

This mean that $\frac{\partial}{\partial\varphi}$ and $\frac{\partial}{\partial\theta}$ are triholomorphic and therefore (5.387) is a toric hyperkahler metric of the type of the Proposition 9, where we should idenfity $t_1 = \theta$ and $t_2 = \varphi$.

For applications to supergravity we will write the metric (5.387) in the form (2.160). Therefore we should find the momentum maps

$$dx_\theta^k = i_\theta \overline{J}^k, \quad dx_\phi^k = i_\phi \overline{J}^k$$

for (5.385). The contraction of $\partial/\partial\theta$ with the hyperkahler form (5.389) gives

$$\begin{aligned} dx_\theta^1 &= \frac{1}{\sqrt{\rho}F} (2q_0 dq_2 + 2q_2 dq_0 - 2q_1 dq_3 - 2q_3 dq_1 - (\frac{1}{2\rho} + \frac{F_\rho}{F}) d\rho - \frac{F_\eta}{F} d\eta), \\ dx_\theta^2 &= \frac{1}{\sqrt{\rho}F} (2q_2 dq_3 + 2q_3 dq_2 - 2q_0 dq_1 - 2q_1 dq_0 - (\frac{1}{2\rho} + \frac{F_\rho}{F}) d\rho - \frac{F_\eta}{F} d\eta), \\ dx_\theta^3 &= \frac{1}{\sqrt{\rho}F} (2q_0 dq_0 + 2q_1 dq_1 - 2q_2 dq_2 + 2q_3 dq_3 - (\frac{1}{2\rho} + \frac{F_\rho}{F}) d\rho - \frac{F_\eta}{F} d\eta). \end{aligned}$$

The last expressions can be integrated to obtain

$$x_\theta^1 = \frac{2(q_0 q_2 + q_1 q_3)}{\sqrt{\rho}F}, \quad x_\theta^2 = \frac{2(q_2 q_3 - q_0 q_1)}{\sqrt{\rho}F}, \quad x_\theta^3 = \frac{q_0^2 - q_1^2 - q_2^2 + q_3^2}{\sqrt{\rho}F}. \quad (5.391)$$

Similarly for $\partial/\partial\varphi$ it is found

$$\begin{aligned} x_\varphi^1 &= \eta x_\theta^1 + \frac{2\sqrt{\rho}(q_1 q_2 - q_0 q_3)}{F}, \quad x_\varphi^2 = \eta x_\theta^2 + \frac{\sqrt{\rho}(q_0^2 - q_1^2 + q_2^2 - q_3^2)}{F}, \\ x_\varphi^3 &= \eta x_\theta^3 + \frac{2\sqrt{\rho}(q_0 q_1 + q_2 q_3)}{F}, \end{aligned} \quad (5.392)$$

in accordance with [132].

The next step is to determine the matrix U_{ij} for (2.175). This is easily found by noticing that from (2.160) it follows that

$$U^{ij} = \overline{g}(\frac{\partial}{\partial t^i}, \frac{\partial}{\partial t^j}), \quad U^{ij} A_j = \overline{g}(\frac{\partial}{\partial t^i}, \cdot). \quad (5.393)$$

Then introducing the expression for (5.387) into the first (5.393) gives

$$U^{ij} = \frac{|q|^2}{F(\frac{1}{4}F^2 - \rho^2(F_\rho^2 + F_\eta^2))} \begin{pmatrix} \frac{1}{2}F - \rho F_\rho & -\rho F_\eta \\ -\rho F_\eta & \frac{1}{2}F + \rho F_\rho \end{pmatrix}, \quad (5.394)$$

with inverse

$$U_{ij} = \frac{F}{|q|^2} \begin{pmatrix} \frac{1}{2}F + \rho F_\rho & \rho F_\eta \\ \rho F_\eta & \frac{1}{2}F - \rho F_\rho \end{pmatrix}. \quad (5.395)$$

To find A_i one should obtain from (5.387) and (2.126) that

$$\begin{aligned}\bar{g}\left(\frac{\partial}{\partial\theta}, \cdot\right) &= \frac{1}{F}[(q_2\sqrt{\rho}+q_3\frac{\eta}{\sqrt{\rho}})(2q_1\tilde{A}^1-dq_0)+2(q_2\frac{\eta}{\sqrt{\rho}}+q_3\sqrt{\rho})(dq_1+q_0\tilde{A}^1)-2(q_0\sqrt{\rho}+q_1\frac{\eta}{\sqrt{\rho}})(dq_2+q_3\tilde{A}^1) \\ &\quad +2(q_0\frac{\eta}{\sqrt{\rho}}-q_1\sqrt{\rho})(dq_3-q_2\tilde{A}^1)], \\ \bar{g}\left(\frac{\partial}{\partial\varphi}, \cdot\right) &= \frac{1}{F}[\frac{q_3}{\sqrt{\rho}}(2q_1\tilde{A}^1-dq_0)+\frac{2q_2}{\sqrt{\rho}}(dq_1+q_0\tilde{A}^1)-\frac{2q_1}{\sqrt{\rho}}(dq_2+q_3\tilde{A}^1)+\frac{2q_0}{\sqrt{\rho}}(dq_3-q_2\tilde{A}^1)],\end{aligned}$$

where $\tilde{A}^1$ is given in (5.386) in terms of F . Then from the second (5.393) it is obtained

$$A_1 = \frac{F}{|q|^2}[(\frac{F}{2} + \rho F_\rho)\bar{g}(\frac{\partial}{\partial\theta}, \cdot) + \rho F_\eta \bar{g}(\frac{\partial}{\partial\varphi}, \cdot)], \quad (5.396)$$

$$A_2 = \frac{F}{|q|^2}[\rho F_\eta \bar{g}(\frac{\partial}{\partial\theta}, \cdot) + (\frac{F}{2} - \rho F_\rho)\bar{g}(\frac{\partial}{\partial\varphi}, \cdot)]. \quad (5.397)$$

Therefore we have reduced (5.387) with this data to the form (2.160). Formulas (5.395), (5.396) and (5.397) define a class of solutions of the Pedersen Poon equations (2.161) and an hyperkahler metric (2.160) simply described in terms of an unknown function F satisfying (2.127). We must recall however that this simplicity is just apparent because U_{ij} depends explicitly on the coordinates $(\rho, \eta, |q|^2)$, which depends implicitly on the momentum maps $(x_\varphi^i, x_\theta^i)$ by (5.391) and (5.392). Therefore U_{ij} is given only as an implicit function of the momentum maps.

For physical applications it is important to find solutions which in this limit tends to

$$\bar{g} = U_{ij}^\infty dx^i \cdot dx^j + U_{ij}^\infty dt_i dt_j, \quad (5.398)$$

for a constant invertible matrix U_{ij}^∞ [138]. Formulas (5.391) and (5.392) shows that the asymptotic limit $x_\theta \rightarrow \infty$ or $x_\varphi \rightarrow \infty$ corresponds to $q \rightarrow \infty$ or $\sqrt{\rho}F \rightarrow 0$. In consequence from (5.395) and (5.394) it follows that $U^{ij} \rightarrow \infty$ and $U_{ij} \rightarrow 0$ asymptotically, which is not the desired result. This problem can be evaded defining a new metric (2.160) with

$$\bar{U}_{ij} = U_{ij} + U_{ij}^\infty, \quad (5.399)$$

and with the same one-forms (5.396) and (5.397) and coordinate system (5.391) and (5.392). Clearly to add this constant do not affect the solution and this data is again a solution of the Pedersen-Poon equation (2.161) for which $\bar{U}_{ij} \rightarrow U_{ij}^\infty$ and $\bar{U}^{ij} \rightarrow U_{ij}^\infty$. Explicitly we have

$$\bar{U}_{ij} = \frac{F}{|q|^2} \begin{pmatrix} \frac{1}{2}F + \rho F_\rho + \frac{U_{11}^\infty |q|^2}{F} & \rho F_\eta + \frac{U_{12}^\infty |q|^2}{F} \\ \rho F_\eta + \frac{U_{12}^\infty |q|^2}{F} & \frac{1}{2}F - \rho F_\rho + \frac{U_{22}^\infty |q|^2}{F} \end{pmatrix}, \quad (5.400)$$

with inverse

$$\bar{U}^{ij} = \frac{1}{\det(\bar{U}_{ij})} \begin{pmatrix} \frac{1}{2}F - \rho F_\rho + \frac{U_{22}^\infty |q|^2}{F} & -\rho F_\eta - \frac{U_{12}^\infty |q|^2}{F} \\ -\rho F_\eta - \frac{U_{12}^\infty |q|^2}{F} & \frac{1}{2}F + \rho F_\rho + \frac{U_{11}^\infty |q|^2}{F} \end{pmatrix}. \quad (5.401)$$

This modified metric is more suitable for physical purposes, but do not correspond to a Calderbank-Pedersen base and this show that the converse of the Swann theorem is not necessarily true.

5.2 Supergravity solutions related to hyperkahler manifolds

The hyperkahler spaces defined by (5.391), (5.392), (5.395), (5.396) and (5.397) can be extended to 11-dimensional supergravity solutions and to IIA and IIB backgrounds by use of dualities. This section presents them following mainly [138], more details can be found there and in references therein.

We consider $D = 11$ supergravity solutions with vanishing fermion fields and $F_{\mu\nu\alpha\beta}$. Such solutions are of the form

$$g = g(E^{2,1}) + U_{ij}dx^i \cdot dx^j + U^{ij}(dt_i + A_i)(dt_j + A_j), \quad (5.402)$$

that is, a direct sum of the flat metric on R^3 and an hyperkahler space in eight dimensions. If all the fields in 11 dimensional supergravity are invariant under the action of an $U(1)$ Killing vector then the 11-dimensional supergravity can be reduced to the type IIA one along the isometry. In order to make such reduction we should write the 11-dimensional metric (5.402) as [179]

$$g = e^{-\frac{2}{3}\varphi(x)} g_{\mu\nu}(x) dx^\mu dx^\nu + e^{\frac{4}{3}\varphi(x)} (dy + C_\mu(x) dx^\mu)^2, \quad (5.403)$$

$$A_{11} = A(x) + B(x) \wedge dy. \quad (5.404)$$

The field A_{11} is the 3-form potential and x^μ are the coordinates of the D=10 spacetime. The $NS \otimes NS$ sector of the IIA supergravity is $(\phi, g_{\mu\nu}, B_{\mu\nu})$ and the $R \otimes R$ sector is $(C_\mu, A_{\mu\nu\rho})$. After reduction of (5.402) along the isometry $t_1 = \varphi$ we find that the non-vanishing fields are

$$g_{10} = \left(\frac{U_{11}}{\det U}\right)^2 [g(E^{2,1}) + U_{ij}dx^i \cdot dx^j] + \left(\frac{1}{U_{11} \det U}\right)^{1/2} (d\theta + A_1)^2, \quad (5.405)$$

$$\phi = \frac{3}{4} \log(U_{11}) - \frac{3}{4} \log(\det U), \quad (5.406)$$

$$C = A_2 - \frac{U_{12}}{U_{11}} (d\theta + A_1). \quad (5.407)$$

The Killing spinor of the 11 dimensional supergravity was also independent of t_1 and the reduced spinor is again a IIA Killing spinor.

The field ϕ is independent of θ and C satisfies $\mathcal{L}_k C = 0$. Then one can use T-duality rules to construct a IIB supergravity solution [51]

$$g = [g_{mn} - g_{\theta\theta}^{-1}(g_{m\theta}g_{n\theta} - B_{m\theta}B_{n\theta})]dx^m dx^n + 2g_{\theta\theta}^{-1}B_{\theta n}d\theta dx^n + g_{\theta\theta}^{-1}d\theta^2 \quad (5.408)$$

$$\tilde{B} = \frac{1}{2}dx^m \wedge dx^n [B_{mn} + 2g_{\theta\theta}^{-1}(g_{m\theta}B_{n\theta})] + g_{\theta\theta}^{-1}g_{\theta m}d\theta \wedge dx^m \quad (5.409)$$

$$\tilde{\phi} = \phi - \log(g_{\theta\theta}) \quad (5.410)$$

where the tilde indicates the transformed fields. The restrictions

$$B = 0, \quad i_k A = 0,$$

gives the T-dual fields

$$\begin{aligned} l &= C_\theta \\ B' &= [C_{mn} - (g_{\theta\theta})^{-1}C_\theta g_{\theta m}]dx^m \wedge d\theta, \\ i_k D &= A, \end{aligned} \quad (5.411)$$

where l is the IIB pseudoscalar, B' is the Ramond-Ramond 2-form potential and D is the IIB 4-form potential. The non vanishing IIB fields resulting from the application of the T-duality are

$$g_{10} = (\det U)^{3/4} [(\det U)^{-1} g(E^{2,1}) + (\det U)^{-1} U_{ij} dx^i \cdot dx^j + d\theta^2], \quad (5.412)$$

$$B_i = A_i \wedge d\theta, \quad (5.413)$$

$$\tau = -\frac{U_{12}}{U_{11}} + i \frac{\sqrt{\det(U)}}{U_{11}}, \quad (5.414)$$

where

$$\tau = l + ie^{-\phi_B}, \quad B_1 = B, \quad B_2 = B',$$

and g_{10} is the Einstein frame metric satisfying

$$g_{10} = e^{-\phi_B/2} g_{IIB}.$$

The examples that we have presented till now are obtained by use of the isometries of the internal spaces. But more backgrounds can be obtained by reducing (5.402) along one of the space directions $E^{2,1}$ and it is obtained the IIA solution

$$g = g(E^{1,1}) + U_{ij} dx^i \cdot dx^j + U^{ij} (dt_i + A_i)(dt_j + A_j), \quad (5.415)$$

with the other fields equal to zero. After T-dualizing in both angular directions it is obtained

$$\begin{aligned} g &= g(E^{1,1}) + U_{ij} dX^i \cdot dX^j, \\ B &= A_i \wedge dt_i, \\ \phi &= \frac{1}{2} \log(\det U), \end{aligned} \quad (5.416)$$

where $X^i = x^i, t^i$. This solution can be lifted to a D=11 supergravity solution

$$g_{11} = (\det U)^{2/3} [(\det U)^{-1} g(E^{1,1}) + (\det U)^{-1} U_{ij} dX^i \cdot dX^j + d\theta^2], \quad (5.417)$$

$$F = F_i \wedge dt_i \wedge d\theta. \quad (5.418)$$

It is possible to generalize (5.402) to include a non vanishing 4-form F . The result is the membrane solution

$$g = H^{-2/3} g(E^{2,1}) + H^{1/3} [U_{ij} dx^i \cdot dx^j + U^{ij} (dt_i + A_i)(dt_j + A_j)], \quad (5.419)$$

$$F = \pm \omega(E^{2,1}) \wedge dH^{-1}, \quad (5.420)$$

where H is an harmonic function on the hyperkahler manifold, i.e, satisfies

$$U^{ij} \partial_i \cdot \partial_j H = 0.$$

We have seen in the last section that every entry of U_{ij} is an harmonic function and so such H can be generated with an hyperbolic eigenfunction F . After reduction along φ it is found the following IIB solution

$$g_{10} = (\det U)^{3/4} H^{1/2} [H^{-1} (\det U)^{-1} g(E^{2,1}) + (\det U)^{-1} U_{ij} dx^i \cdot dx^j + H^{-1} d\theta^2], \quad (5.421)$$

$$B_i = A_i \wedge d\theta, \quad (5.422)$$

$$\tau = -\frac{U_{12}}{U_{11}} + i\frac{\sqrt{\det U}}{U_{11}}, \quad (5.423)$$

$$i_k D = \pm \omega(E^{2,1}) \wedge dH^{-1}. \quad (5.424)$$

If instead (5.419) is dimensionally reduced along a flat direction it is obtained the IIA solution

$$g = g(E^{1,1}) + U_{ij} dx^i \cdot dx^j + U^{ij} (dt_i + A_i)(dt_j + A_j), \quad (5.425)$$

$$B = \omega(E^{1,1}) H^{-1}, \quad (5.426)$$

$$\phi = -\frac{1}{2} \log(H). \quad (5.427)$$

A double dualization gives a new IIA solution

$$g = H^{-1} g(E^{1,1}) + U_{ij} dX^i \cdot dX^j, \quad (5.428)$$

$$B_i = A_i \wedge d\varphi^i + \omega(E^{1,1}) H^{-1}, \quad (5.429)$$

$$\phi = \frac{1}{2} \log(\det U) - \frac{1}{2} \log(H). \quad (5.430)$$

The lifting to eleven dimensions gives

$$g_{11} = H^{1/3} (\det U)^{2/3} [H^{-1} (\det U)^{-1} g(E^{1,1}) + (\det U)^{-1} U_{ij} dX^i \cdot dX^j + H^{-1} d\theta^2], \quad (5.431)$$

$$F = (F_i \wedge dt_i + \omega(E^{1,1}) \wedge dH^{-1}) \wedge d\theta. \quad (5.432)$$

All the backgrounds presented in this section can be constructed with a single F satisfying (2.127), but the dependence on (x_θ, x_φ) remains implicit.

6. G_2 toric metrics and supergravity backgrounds

If the Bryant-Salamon extension is applied to the Calderbank-Pedersen spaces the $U(1) \times U(1)$ isometry is preserved and the result is a toric G_2 holonomy metric. In this subsection the G_2 holonomy metrics corresponding to the examples A and B of subsection 2.4.7 will be constructed. We select such cases by simplicity, but the procedure could be applied to any quaternion Kahler space as well. By use of the results of the preceding section we extend this toric G_2 holonomy metrics to vacuum configurations of the eleven dimensional supergravity preserving at least one supersymmetry. Also type IIA backgrounds will be found by reduction along one of the isometries.

It is straightforward to show that the explicit form of (2.245) when the base space (and, in consequence, the total one) has torus symmetry is

$$g = \frac{dr^2}{h(r)} + \frac{r^2}{2}[U_{\phi\phi}d\phi^2 + U_{\phi\psi}d\phi d\psi + U_{\psi\psi}d\psi^2 + Q_\phi d\phi + Q_\psi d\psi + g_{\rho\rho}(d\eta^2 + d\rho^2) + H], \quad (6.433)$$

where it has been defined

$$\begin{aligned} h(r) &= 1 - 4c/r^4 \\ U_{11} &= g_{\phi\phi} + h(r) \frac{u_1^2(\rho^2 + \eta^2) + (u_2\eta + u_3\rho)^2}{2\rho F^2}, \\ U_{22} &= g_{\psi\psi} + h(r) \frac{u_1^2 + u_2^2}{2\rho F^2}, \\ U_{12} = U_{21} &= g_{\phi\psi} + h(r) \frac{(u_1^2 + u_2^2)\eta + u_2u_3\rho}{2\rho F^2}, \\ Q_\phi &= h(r) \frac{1}{\sqrt{\rho}F} [u_1(\eta du_2 + \rho du_3) - (u_2\eta + u_3\rho)du_1 - u_1(u_3\eta - u_2\rho)A^1], \\ Q_\psi &= h(r) \left[\frac{u_1 du_2 - u_2 du_1 - u_1 u_3 A^1}{\sqrt{\rho}F} \right], \\ H &= h(r) [|d\vec{u}|^2 + (u_1^2 + u_2^2)(A^1)^2 - 2A^1(u_3 du_2 - u_2 du_3)]. \end{aligned}$$

The second rank tensor g_{ab} is the metric of the base manifold. The product metric of (6.433) with M^4

$$g_{11} = ds_M^2 + \frac{dr^2}{h(r)} + \frac{r^2}{2}[U_{\phi\phi}d\phi^2 + U_{\phi\psi}d\phi d\psi + U_{\psi\psi}d\psi^2 + Q_\phi d\phi + Q_\psi d\psi + g_{\rho\rho}(d\eta^2 + d\rho^2) + H]. \quad (6.434)$$

is a vacuum configuration of the eleven dimensional supergravity [26]. With the help of the quantities

$$\begin{aligned} \alpha_1 &= \frac{U_{22}Q_\phi - U_{12}Q_\psi}{\det U}, \quad \alpha_2 = \frac{U_{11}Q_\psi - U_{12}Q_\phi}{\det U}, \\ h &= H + U_{11}\alpha_1^2 + 2U_{12}\alpha_1\alpha_2 + U_{22}\alpha_2^2, \\ \phi_1 &= \phi, \quad \phi_2 = \psi, \end{aligned}$$

the metric (6.434) is expressed in more simple manner as

$$g_{11} = ds_M^2 + \frac{dr^2}{h(r)} + \frac{r^2}{2}[U_{ij}(d\phi_i + \alpha_i)(d\phi_j + \alpha_j) + h].$$

The last expression takes the usual form of the Kaluza-Klein ansatz

$$g_{11} = e^{-\frac{2}{3}\varphi_D} G_{\mu\nu} dx^\nu dx^\mu + e^{\frac{4}{3}\varphi_D} (d\phi + dx^\mu C_\mu(x))^2. \quad (6.435)$$

with the dilaton field and the RR 1-form defined by

$$\varphi_D = \frac{3}{4} \log\left(\frac{r^2 U_{11}}{2}\right),$$

$$C = \frac{U_{12} d\psi + Q_\phi}{U_{11}}.$$

The usual reduction procedure [179] for (6.435) along ϕ_1 gives the following IIA metric in the string frame

$$g_A = \left(\frac{r^2 U_{11}}{2}\right)^{1/2} \left\{ ds_{M^4}^2 + \frac{dr^2}{h(r)} + \frac{r^2}{2U_{11}} [\det U d\psi^2 + 2(U_{11} Q_\psi - U_{12} Q_\phi) d\psi - Q_\phi^2 + U_{11} H] \right\}. \quad (6.436)$$

The components of (6.436) are

$$g_{\psi\psi} = \left(\frac{r^2 U_{11}}{2}\right)^{1/2} \frac{r^2}{4U_{11} F \sqrt{\rho}} h(r) [U_{11} \sin^2(\theta) - U_{12} \left(\frac{\rho}{2} \sin(2\theta) \sin(\varphi) + \eta \sin^2(\theta)\right)]$$

$$g_{\psi\theta} = \left(\frac{r^2 U_{11}}{2}\right)^{1/2} \frac{r^2}{4U_{11} F} h(r) U_{12} \sqrt{\rho} \cos(\varphi)$$

$$g_{\theta\theta} = \left(\frac{r^2 U_{11}}{2}\right)^{1/2} \left[\frac{r^2 h(r)}{4} - \frac{r^2 h^2(r)}{8U_{11} \rho F^2} \rho^2 \cos^2(\varphi) \right]$$

$$g_{\varphi\varphi} = \left(\frac{r^2 U_{11}}{2}\right)^{1/2} \left[\frac{r^2 h(r)}{4} \sin^2(\theta) - \frac{r^2 h^2(r)}{8U_{11} \rho F^2} \sin^2(\theta) (\eta \sin(\theta) + \rho \cos(\theta) \sin(\varphi))^2 \right]$$

$$g_{\varphi\theta} = \left(\frac{r^2 U_{11}}{2}\right)^{1/2} \frac{r^2 h^2(r)}{4U_{11} F^2} \sin(\theta) \cos(\varphi) (\eta \sin(\theta) + \rho \cos(\theta) \sin(\varphi)),$$

where it have been introduced the spherical coordinates θ, φ through the relations

$$u_1 = \sin(\theta) \cos(\varphi), \quad u_2 = \sin(\theta) \sin(\varphi), \quad u_3 = \cos(\theta).$$

The range of this coordinates is $\theta \in [0, \pi]$ and $\varphi \in [0, 2\pi]$; the other components of the metric are identically zero.

The base metric (2.129) have a singularity at $\rho \rightarrow 0$. For this case it is obtained

$$U_{11} = \frac{\rho^2(1+3\rho) + \eta^2(9+7\rho)}{2\rho^5} + 2h(r) \left[\frac{u_1^2(\rho^2 + \eta^2) + (u_2\eta + u_3\rho)^2}{\rho^4} \right] \sim \frac{f(x^i)}{\rho^5},$$

$$U_{22} = \frac{8}{\rho^4} + 2h(r) \left(\frac{u_1^2 + u_2^2}{\rho^4} \right) \sim \frac{f(x^i)}{\rho^4},$$

$$U_{12} = U_{21} = \frac{8\eta}{\rho^4} + h(r) \left[\frac{(u_1^2 + u_2^2)\eta + u_2 u_3 \rho}{\rho^4} \right] \sim \frac{f(x^i)}{\rho^4},$$

$$Q_\phi = h(r) \frac{2}{\rho^2} [u_1(\eta du_2 + \rho du_3) - (u_2\eta + u_3\rho) du_1]$$

$$\begin{aligned}
& -u_1(u_3\eta - u_2\rho)\frac{2d\eta}{\rho}] \sim \frac{f(x^i, dx^i)}{\rho^3}, \\
Q_\psi &= h(r)\frac{1}{\rho^2}(u_1du_2 - u_2du_1 - u_1u_3\frac{2d\eta}{\rho}) \sim \frac{f(x^i, dx^i)}{\rho^3}, \\
H &= h(r)[|d\vec{u}|^2 + 4(u_1^2 + u_2^2)\frac{d\eta^2}{\rho^2} - 4\frac{d\eta}{\rho}(u_3du_2 - u_2du_3)] \sim \frac{f(x^i, dx^i)}{\rho^2}.
\end{aligned}$$

where x^i denotes all the coordinates except ρ and the behaviour for short distances was evaluated. The dilaton field is given explicitly as

$$\varphi_A = \frac{3}{4} \log\left\{\frac{r^2}{2}\left[\frac{\rho^2(1+3\rho) + \eta^2(9+7\rho)}{2\rho^5}\right] + r^2 h(r)\left[\frac{u_1^2(\rho^2 + \eta^2) + (u_2\eta + u_3\rho)^2}{\rho^4}\right]\right\} \sim \log\left(\frac{f(x^i)}{\rho^5}\right).$$

The expression for the R-R one form is

$$\begin{aligned}
C &= \frac{2\rho\{8\eta + h(r)[(u_1^2 + u_2^2)\eta + u_2u_3\rho]\}d\psi}{\rho^2(1+3\rho) + \eta^2(9+7\rho) + 4\rho h(r)[u_1^2(\rho^2 + \eta^2) + (u_2\eta + u_3\rho)^2]} \\
&+ \frac{4\rho^2 h(r)[\rho u_1(\eta du_2 + \rho du_3) - \rho(u_2\eta + u_3\rho)du_1 - 2u_1(u_3\eta - u_2\rho)d\eta]}{\rho^2(1+3\rho) + \eta^2(9+7\rho) + 4\rho h(r)[u_1^2(\rho^2 + \eta^2) + (u_2\eta + u_3\rho)^2]} \sim f(x^i)\rho.
\end{aligned}$$

The components of (6.436) diverges in this case for short ρ ,

$$\begin{aligned}
g_{\psi\varphi} &\sim \frac{f(x^i)}{\rho^5}, \quad g_{\psi\theta} \sim \frac{f(x^i)}{\rho^4}, \quad g_{\theta\theta} \sim \frac{f(x^i)}{\rho^2} \\
g_{\varphi\varphi} &\sim \frac{f(x^i)}{\rho^2}, \quad g_{\varphi\theta} \sim \frac{f(x^i)}{\rho}.
\end{aligned}$$

The quaternionic space (2.130) is singular too in the limit $\rho \rightarrow 0$. Using it as a base space gives

$$\begin{aligned}
U_{11} &= \frac{8\eta^4\rho^2(19+5\rho) + 16\eta^6(9+7\rho) + \rho^6(1+35\rho) + 3\eta^2\rho^4(35+61\rho)}{9\rho^5(\rho^2 - 4\eta^2)^2(8\eta^4 + 6\eta^2\rho^2 + 3\rho^4)} \\
&+ 8h(r)\left[\frac{u_1^2(\rho^2 + \eta^2) + (u_2\eta + u_3\rho)^2}{9\rho^4(\rho^2 - 4\eta^2)^2}\right] \sim \frac{f(x^i)}{\rho^5}, \\
U_{22} &= \frac{64(4\eta^4 + \rho^4)}{9\rho^4(\rho^2 - 4\eta^2)^2(8\eta^4 + 6\eta^2\rho^2 + 3\rho^4)} + 8h(r)\left[\frac{u_1^2 + u_2^2}{9\rho^4(\rho^2 - 4\eta^2)^2}\right] \sim \frac{f(x^i)}{\rho^4}, \\
U_{12} = U_{21} &= \frac{32(8\eta^5 - 4\eta^3\rho^2 + 3\eta\rho^4)}{9\rho^4(\rho^2 - 4\eta^2)^2(8\eta^4 + 6\eta^2\rho^2 + 3\rho^4)} + 8h(r)\left[\frac{(u_1^2 + u_2^2)\eta + u_2u_3\rho}{9\rho^4(\rho^2 - 4\eta^2)^2}\right] \sim \frac{f(x^i)}{\rho^4}, \\
Q_\phi &= 4h(r)\frac{1}{3\rho^2(4\eta^2 - \rho^2)}\{u_1(\eta du_2 + \rho du_3) - (u_2\eta + u_3\rho)du_1 \\
&- u_1(u_3\eta - u_2\rho)\left[\frac{8\eta}{\rho^2 - 4\eta^2}d\rho + \frac{4(\rho^2 - 2\eta^2)}{\rho(\rho^2 - 4\eta^2)}d\eta\right]\} \sim \frac{f(x^i, dx^i)}{\rho^3}, \\
Q_\psi &= h(r)\frac{4}{3\rho^2(4\eta^2 - \rho^2)}\{u_1du_2 - u_2du_1 - u_1u_3\left[\frac{8\eta}{\rho^2 - 4\eta^2}d\rho + \frac{4(\rho^2 - 2\eta^2)}{\rho(\rho^2 - 4\eta^2)}d\eta\right]\} \sim \frac{f(x^i, dx^i)}{\rho^3}, \\
H &= h(r)\{|d\vec{u}|^2 + (u_1^2 + u_2^2)\left[\frac{8\eta}{\rho^2 - 4\eta^2}d\rho + \frac{4(\rho^2 - 2\eta^2)}{\rho(\rho^2 - 4\eta^2)}d\eta\right]^2 - 2\left[\frac{8\eta}{\rho^2 - 4\eta^2}d\rho\right.
\end{aligned}$$

$$+\frac{4(\rho^2-2\eta^2)}{\rho(\rho^2-4\eta^2)}d\eta](u_3du_2-u_2du_3)\}\sim\frac{f(x^i,dx^i)}{\rho}.$$

The dilaton field is expressed through the relation

$$e^{\frac{4}{3}\varphi_D}=\frac{r^2}{2}\{\frac{8\eta^4\rho^2(19+5\rho)+16\eta^6(9+7\rho)+\rho^6(1+35\rho)+3\eta^2\rho^4(35+61\rho)}{9\rho^5(\rho^2-4\eta^2)^2(8\eta^4+6\eta^2\rho^2+3\rho^4)}\\+8h(r)[\frac{u_1^2(\rho^2+\eta^2)+(u_2\eta+u_3\rho)^2}{9\rho^4(\rho^2-4\eta^2)^2}]\},$$

from where follows that

$$\varphi_D\sim\log(\frac{f(x^i)}{\rho^5}),\quad\rho\rightarrow 0.$$

The behavior of the RR one-form at short distances results

$$C\sim f(x^i,dx^i)\rho.$$

and the components of the IIA metric diverges as

$$g_{\psi\varphi}\sim\frac{f(x^i)}{\rho^5},\quad g_{\psi\theta}\sim\frac{f(x^i)}{\rho^4},\quad g_{\theta\theta}\sim\frac{f(x^i)}{\rho^2}\\g_{\varphi\varphi}\sim\frac{f(x^i)}{\rho^2},\quad g_{\varphi\theta}\sim\frac{f(x^i)}{\rho}.$$

A detailed analysis of the singularities of the backgrounds corresponding to the m -pole solutions and their physical interpretation was given in [97] and [98], the interested reader may consult those references. To finish we recall that the toric G_2 spaces presented here are not unique, other toric examples were also constructed in [176] and match for application to M-theory as well by an procedure analog of those presented here.

7. Conclusion

The main results presented in this dissertation are the following

- We have shown that in $d = 4$ weak hyperkahler torsion structures are the same that hypercomplex structures and the same that the Plebanski-Finley conformally invariant heavens. With the help of this identification we have found the most general local form of an hyperkahler torsion space in four dimensions. We also presented an Ashtekar like formulation for them in which to finding an hyperkahler torsion metric is reduced to solve a quadratic differential system. We have shown that there exists more solutions than the Callan-Harvey-Strominger ones, this are the structures for which it does not exist a preserved volume form. Some examples with isometries have been constructed.

- It is found the most general form for the target space metric to the moduli space metric of several ($n > 1$) identical matter hypermultiplets for the type-IIA superstrings compactified on a Calabi-Yau threefold, near conifold singularities, even taking into account non-perturbative D-instanton quantum corrections. The metric in consideration is "toric hyperkahler" if we do not take into account the gravitational corrections. This generalize the Ooguri-Vafa solution and has been obtained by using essentially the same assumptions than them. Some new effects related to interference between hypermultiplets have been presented, which do not hold in the presence of a single hypermultiplet.

- It is constructed a family of toric hyperkahler spaces in eight dimensions by use of a mathematical relation called Swann extension, which relates quaternion Kahler and hyperkahler spaces in different dimensions. The resulting hyperkahler space is expressed in a coordinate system, for which it takes the generalized Gibbons-Hawking type. The solution was lifted to a background of the eleven dimensional supergravity preserving certain ammount of supersymmetry, with and without the presence of fluxes. Several type IIA and IIB backgrounds have been found by reduction along a circle and by use of T-duality rules.

- It is constructed a family of $Spin(7)$, G_2 and weak G_2 holonomy metrics. The result has been lifted to a supergravity solution preserving one supersymmetry. The presence of a toric symmetry allows a reduction to type IIA backgrounds by usual reduction along one of the Killing vectors.

I acknowledge specially to my director A.P.Isaev. I have been benefited during my studies with discussions with A.Pashnev, S.Ketov, M.Tsulaia, C.Sochichiu, E.Ivanov, D. Mladenov, B.Dimitrov, B.Saha and M.Paucar Acosta. Finally I would like to acknowledge the support of CLAF and to dedicate this dissertation to Professor L.Maspero.

A Supersymmetric sigma models

In this section we sketch why strong torsion hyperkahler geometry is the target space metric of $N = 4$ supersymmetric sigma models [44]. Sigma models in dimension two are described by Lagrangians of the form

$$I_b = -\frac{1}{2\pi} \int d^2x [g_{AB}(\varphi) \partial_\mu \varphi^A \partial^\mu \varphi^B + b_{AB}(\varphi) \epsilon^{\mu\nu} \partial_\mu \varphi^A \partial_\nu \varphi^B], \quad (1.437)$$

being g_{AB} an arbitrary metric tensor⁷. The fields φ^A are bosonic ($A = 1, \dots, d$) depending on the two coordinates x^i ($i = 1, 2$). This mean that φ^A parameterize two-cycles on a d-dimensional target manifold M with a metric g_{AB} . The second term is a Wess-Zumino-Witten type term.

If we want to make the supersymmetric extension of (1.437) we need to include a number of fermion degrees equal to the number of the boson ones. Thus we need to introduce a set of spinors ψ'^A and consider a supersymmetry transformation which on general grounds should be of the form

$$\delta \varphi^A = \bar{\varepsilon}_1 J_B^A \psi'^B \quad (1.438)$$

being J some $(1, 1)$ tensor. We still have the freedom to make the redefinition $\psi'^B = (J^{-1})_A^B \psi^A$ in (1.438) and consider the transformation

$$\delta \varphi^A = \bar{\varepsilon}_1 \psi^A \quad (1.439)$$

The $N = (1, 0)$ supersymmetric extension of (1.437) is [53]

$$I_{sb} = -\frac{1}{2\pi} \int d^2x [g_{AB}(\varphi) \partial_\mu \varphi^A \partial^\mu \varphi^B + b_{AB}(\varphi) \epsilon^{\mu\nu} \partial_\mu \varphi^A \partial_\nu \varphi^B + \bar{\psi}^A \gamma^\mu D_\mu \psi^B g_{AB}], \quad (1.440)$$

where the spinors ψ^A are right-handed and ε is left-handed and we have defined the covariant derivative

$$D_\mu \psi^A = \partial_\mu \psi^A + \Upsilon_{BC}^A \psi^B \partial_\mu \varphi^C, \\ \Upsilon_{BC}^A = \Gamma_{BC}^A + \frac{1}{2} T_{BC}^A$$

being Γ_{BC}^A the Christoffel symbols of the Levi-Civita connection for g_{AB} . The torsion form T_{ABC} is entirely defined in terms of the Wess-Zumino-Witten potential b_{AB} as

$$2T_{ABC} = -3b_{[AB,C]} \iff dT = 0. \quad (1.441)$$

The indices A, B are lowered and raised through the metric g_{AB} . The action (1.440) is invariant under (1.439) and

$$\delta \psi^A = \gamma^\mu \partial_\mu \varphi^A \varepsilon - \psi^B \bar{\varepsilon} S_{BC}^A \psi^C. \quad (1.442)$$

The tensor S_{ABC} is constrained by supersymmetry arguments to be skew symmetric and covariantly constant

$$S_{ABC;D} = S_{ABC,D} - 3S_{E[AB} \Upsilon_{C]D}^E = 0.$$

⁷In the main text we have used latin indices (a,b,..) as flat and greek indices ($\alpha, \beta, \dots$) as curved. Instead in this appendix capital latin indices (A,B,..) are curved, we use capital latin indices and not greek in order to not confuse them with the greek indices used to denote space-time derivatives ∂_μ or gamma matrices γ^μ .

The presence of a covariantly constant 3-form gives a new restriction on the manifold. On group manifolds this condition has solution and S can be choose proportional to T [53]. We find the commutator

$$[\delta, \delta']\varphi^A = \bar{\varepsilon}'\gamma^\mu\varepsilon(2\partial_\mu\varphi^A - \bar{\psi}^B\gamma_\mu S_{BC}^A\psi^C), \quad (1.443)$$

and correspond to the usual supersymmetry algebra without the presence of central charges.

Let us consider now the problem to construct an $N = (2, 0)$ supersymmetric extension of (1.437). For this purpose we should impose a new supersymmetry transformation to (1.440) without spoiling (1.439) and (1.442). As we saw the general supersymmetry variation for φ^A is of the form

$$\delta^1\varphi^A = \bar{\varepsilon}_1(J^1)_B^A\psi^B, \quad (1.444)$$

being J^1 certain $(1, 1)$ tensor. Due to the invariance of (1.440) with respect to the other symmetry (1.442) we have to impose

$$I_{sb}(\varphi, J^1\psi) = I_{sb}(\varphi, \psi). \quad (1.445)$$

Formula (1.445) implies that

$$g_{AB}(J^1)_C^B + (J^1)_A^B g_{BC} = 0, \quad (1.446)$$

and then the metric is hermitian with respect to the tensor J^1 . Condition (1.445) also implies that

$$D_A(J^1)_C^B = 0 \quad \implies \quad (J^1)_{[AB,C]} = -2(J^1)_{[A}^D T_{B,C]D}. \quad (1.447)$$

This mean that J^1 is covariantly constant with respect to the connection with torsion. The compatibility condition of (1.447) is

$$(J^1)_E^A R_{BCD}^E = R_{ECD}^A (J^1)_B^E, \quad (1.448)$$

and more generally the tensor J^1 commute with all the generators of the holonomy group. The new supersymmetry transformations result

$$\delta^1\varphi^A = \bar{\varepsilon}_1(J^1)_B^A\psi^B.$$

$$\delta^1\psi^A = \gamma^\mu\partial_\mu\varphi^B(J^1)_B^A\varepsilon_1 - [(J^1)_D^A\Upsilon_{BE}^D(J^1)_C^E + (S_1)_{BC}^A]\psi^B\bar{\varepsilon}_1\psi^C. \quad (1.449)$$

The tensor $(S_-)_{BC}^A$ is not determined by other quantities. The commutator of the two supersymmetries acting over φ^A is

$$\begin{aligned} [\delta, \delta^1]\varphi^A &= \bar{\varepsilon}_1\gamma_\mu\varepsilon_1[(J^1)_B^A + (J^1)_A^B]\partial_\mu\varphi^B \\ &\quad + \bar{\varepsilon}_1\gamma_\mu\varepsilon_1\bar{\psi}^B\gamma_\mu\psi^C(J^1)_D^A N_{BC}^D, \end{aligned} \quad (1.450)$$

plus terms depending on $(S_1)_{BC}^A$. Then the usual supersymmetric algebra holds if (1.450) is zero and this implies that

$$(J^1)_B^A + (J^1)_A^B = 0, \quad N_{AB}^C = 0, \quad (S_1)_{BC}^A = 0. \quad (1.451)$$

The first equation (1.451) together with (1.446) implies that J^1 is an almost complex structure and therefore the dimension of the target manifold should be even ($d=2n$). The second (1.451) show that J^1 is integrable. The metrics g_{AB} for which (1.447) and (1.451) hold are Kahler torsion. We conclude that $N = 2$ supersymmetric sigma models occur on *Kahler torsion manifolds*.

If there is a third supersymmetry corresponding to another complex structure J^2 and to a parameter ε_2 then it is obvious that the previous reasoning is true and the properties (1.446), (1.447) and (1.451) hold for J^2 . But we obtain new restrictions by requiring that the transformation corresponding to ε_1 and ε_2 close to a supersymmetry. The algebra

$$[\delta(\varepsilon_1), \delta(\varepsilon_2)]\varphi^A = 2\bar{\varepsilon}_2^i \gamma^\mu \varepsilon_1^j \delta_{ij} \partial_\mu \varphi^A \quad (1.452)$$

(which correspond to (1.443) with $S_{BC}^A = 0$) is obtained if and only if

$$J^i \cdot J^j + J^j \cdot J^i = \delta_{ij} I \quad (1.453)$$

$$N^{ij}(X, Y) = [X, Y] + J^i[X, J^j Y] + J^j[J^i X, Y] - [J^i X, J^j Y] = 0. \quad (1.454)$$

$N^{ij}(X, Y)$ is known as the mixed Niejenhuis tensor.

Let us define the tensor $J^3 = J^1 \cdot J^2$ for a $N = (3, 0)$ supersymmetric sigma model. It follows from (1.453) that

$$(J^3)^2 = -I$$

and therefore J^3 is also an almost complex structure. It is direct to check

$$J^i \cdot J^j = -\delta_{ij} + \epsilon^{ijk} J^k. \quad (1.455)$$

The integrability of J^1 and J^2 implies the integrability of J^3 and (1.454). Also (1.447) for J^1 and J^2 implies that

$$D_A(J^3)_C^B = 0. \quad (1.456)$$

Therefore $N = (3, 0)$ supersymmetry implies $N = (4, 0)$ supersymmetry for sigma models and from (1.454), (1.441), (1.447) and (1.455) it follows that the target metric g_{AB} of a $N = (4, 0)$ supersymmetric sigma model is always *strong hyperkahler torsion*. If the form b_{AB} is zero, then the geometry presented here reduce to the usual Kahler and hyperkahler ones.

The weak cases are also of interest in the context of low energy heterotic string actions because they contain a three form that is not closed due to a Green-Schwarz anomaly. For more details of this assertions see [55]-[56], and for more applications of this geometry in the construction of sigma models see [186]-[188].

References

- [1] M.Berger, Bull.Soc.Math.France. 83 (1955) 279.
- [2] E.Cartan Acta.Math 48 (1926) 1-42.
- [3] A.Wolf J.Math.Mech.14 (1965) 1033; D.Alekseevskii Func.Anal.Appl.2 (1968) 11.
- [4] J.Ishihara J.Diff.Geom.9 (1974) 483; S.Salamon Invent.Math.67 (1982) 143.
- [5] B.de Wit and A.Van Proeyen Commun.Math.Phys.149 (1992) 307.
- [6] A. Besse "Einstein manifolds", Springer Verlag, second edition 2002.
- [7] L. Landau and E. Lifshitz "The classical theory of fields", Pergamon Press, fourth edition 1975.
- [8] A. Ambrose and I. Singer Trans.Am.Math.Soc 75 (1953) 428.
- [9] D.Joyce, J.Diff.Geom. 43 (1996) 2; D.Joyce, J.Diff.Geom. 43 (1996) 329.
- [10] M. Atiyah and N.J.Hitchin The geometry and dynamic of magnetic monopoles, Princeton University Press 1988.
- [11] M.Atiyah, V.Drinfeld, N.Hitchin and Y.Manin Phys.Lett.A 65 (1978) 185.
- [12] N.J.Hitchin, A.Karlhede, U.Lindstrom and M.Rocek Comm.Math.Phys.108 (1987) 535.
- [13] K.Galicki Comm.Math.Phys.108 (1987) 117.
- [14] E.Witten and J.Bagger Nucl.Phys.B 222 (1983) 1.
- [15] P.Fre Nucl. Phys. Proc. Suppl. 45BC (1996) 59.
- [16] L. Andrianopoli, M. Bertolini, A. Ceresole, R. D'Auria, S. Ferrara, P. Fre' and T. Magri J.Geom.Phys.23 (1997) 111.
- [17] I.Y. Park and R. von Unge JHEP 0003 (2000) 037; U.Lindstrom, M.Rocek and R.von Unge JHEP 0001 (2000) 022.
- [18] K. Behrndt and G. Dall'Agata Nucl.Phys.B 627 (2002) 357.
- [19] I.Antoniadis and B.Pioline Int.J.Mod.Phys. A12 (1997) 4907.
- [20] B.de Wit, M.Rocek, S.Vandoren Phys.Lett.B 511 (2001) 302.
- [21] P.B.Kronheimer J.Diff.Geometry 29 (1989) 665; P.B.Kronheimer J.Diff.Geometry 29 (1989) 685.
- [22] A.Swann Math.Ann.289 (1991) 421.
- [23] L.Anguelova, M.Rocek and S.Vandoren JHEP 0205 (2002) 064.
- [24] L. Jarv, T. Mohaupt and F. Saueressig JHEP 0312 (2003) 047; L. Jarv, T. Mohaupt and F. Saueressig hep-th/0310174.

- [25] J. Gutowski, G. Papadopoulos, Nucl.Phys. B615 (2001) 237.
- [26] M.Duff, B.Nilsson and C.Pope, Phys.Rep. 130 (1986) 1.
- [27] R.Bryant, Ann.Math.126 (1987) 525.
- [28] G.Gibbons, D.Page and C.Pope, Commun.Math.Phys. 127 (1990) 529.
- [29] R. Bryant and S.Salamon, Duke Math. Journal 58 (1989) 829.
- [30] A. Brandhuber, J. Gomis, S. Gubser and S. Gukov, Nucl.Phys. B611 (2001) 179.
- [31] A.Brandhuber, Nucl.Phys. B629 (2002) 393.
- [32] D.Joyce, Compact manifolds with special holonomy, first edition (Oxford University Press 2000).
- [33] Y.Konishi and M.Naka, Class.Quant.Grav. 18 (2001) 5521.
- [34] M.Cvetic, G.Gibbons, H.Lu and C.Pope, Phys.Lett. B534 (2002) 172.
- [35] M.Cvetic, G.Gibbons, H.Lu and C.Pope Annals Phys. 310 (2004) 265.
- [36] M. Cvetic, G.W. Gibbons, H. Lu and C.N. Pope, Phys.Rev. D65 (2002) 106004.
- [37] Z.W. Chong, M. Cvetic, G.W. Gibbons, H. Lu, C.N. Pope and P. Wagner, Nucl.Phys. B638 (2002) 459.
- [38] M. Cvetic, G.W. Gibbons, H. Lu and C.N. Pope Class.Quant.Grav. 20 (2003) 4239.
- [39] N. Hitchin, Stable forms and special metrics, math.DG/0107101.
- [40] E.Witten Nucl.Phys.B186 (1981) 412.
- [41] B.Acharya, Adv.Theor.Math.Phys. 3 (1999) 227.
- [42] B.Acharya and E.Witten, Chiral Fermions from Manifolds of G_2 Holonomy, hep-th/0109152.
- [43] A.Bilal and S.Metzger Nucl.Phys. B 663 (2003) 343.
- [44] L.Alvarez-Gaume and D.Freedman Commun.Math.Phys.80 (1981) 443.
- [45] E.Witten Commun.Math.Phys. 92 (1984) 455.
- [46] T.Curtright and C.Zachos Phys.Rev.Lett 53 (1984) 1799.
- [47] S.Gates C.Hull and M.Rocek, Nucl.Phys.B 248 (1984), 157.
- [48] P.Howe and G.Sierra Phys.Lett.148B (1984) 451.
- [49] A.Sen Phys.Rev.D 32 (1985) 2102.
- [50] C.Hull and E.Witten Phys.Lett 160 B (1985) 398.

- [51] E.Bergshoeff and E.Sezgin Mod.Phys.Lett A 1 (1986) 191.
- [52] P.Howe and G.Papadopoulos Class.Quant.Grav. 4 (1987) 1794; 5 (1987) 501.
- [53] P.Spindel, A.Servin, W.Troost and A.Van Proeyen Nucl.Phys.B 308 (1988) 662.
- [54] C.Callan Instantons and Solitons in Heterotic String Theory hep-th/9109052.
- [55] C.Callan J.Harvey and A.Strominger Nucl.Phys.B 359 (1991) 611.
- [56] A.Strominger Nucl.Phys.B274 (1986) 253.
- [57] G.W. Gibbons, G. Papadopoulos and K.S. Stelle Nucl.Phys. B508 (1997) 623-658.
- [58] A.Giveon and M.Rocek Nucl.Phys. B421 (1994) 173-190, Introduction to Duality hep-th/9406178.
- [59] J.Grant and I.Strachan, Hypercomplex Integrable Systems, solv-int/9808019.
- [60] C.P.Boyer Proceedings of the American Mathematical Society 102, 157 (1988).
- [61] J. Plebanski and Finley Journal of Mathematical Physics, vol 17, number 4 (1976), 585.
- [62] N.Hitchin Complex manifolds and Einstein's equations in Twistor geometry and Non-linear Systems H.D.Doebner and T.D.Palev eds. Springer 1982.
- [63] M.Atiyah, N.Hitchin and I.Singer Proc.R.Soc.London.A 362 (1978) 425.
- [64] A.Ashtekar T.Jacobson and L.Smolin Comm.Math.Phys.115 (1988) 631; T.Jacobson and E.Newman Comm.Math.Phys.121 (1989) 659.
- [65] P.Jones and K.Tod Class.Quant.Grav.2 (1985) 565.
- [66] G.Bonneau and G.Valent Class.Quant.Grav. 11 (1994) 1133-1154.
- [67] G.Papadopoulos Phys.Lett B 356 (1995) 249.
- [68] T.Chave, K.Tod and G.Valent Phys.Lett.B383 (1996) 262.
- [69] E.Ivanov Phys.Lett. B 356 (1995) 239.
- [70] F.Delduc, S.Kalitzin and E.Sokatchev Class.Quant.Grav. 7 (1990) 1567.
- [71] P.S. Howe and G. Papadopoulos Commun.Math.Phys. 151 (1993) 467-480.
- [72] G.Gibbons and S.Hawking Phys.Lett.B 78 (1978) 430.
- [73] S.Kloster, M.Som and A.Das J.Math.Phys 23 (1974) 1096.
- [74] J. Gegenberg and A.Das Gen.Rel.Grav 16 (1982) 817.
- [75] C.Boyer and J.Finley J.Math.Phys.23 (1982) 1126.
- [76] J.Plebanski J.Math.Phys.16 (1975) 2395-2402.

- [77] M.Dunajski J.Geom.Phys.30 (1989) 266.
- [78] M. Przanoskwi Acta Phys.Polon.B14 (1983) 625-627.
- [79] M. Przanoskwi Acta Phys.Polon.B18 (1987) 879-889.
- [80] M. Przanoskwi Phys.Lett.A110 (1985) 295-297.
- [81] M. Przanoskwi J.Math.Phys.32 (1991) 1004-1010.
- [82] C.LeBrun Comm.Math.Phys 118 (1988) 591.
- [83] D. Joyce Duke Mathematical Journal 77 (1995) 519.
- [84] G.Gantcharov and Y.S.Poon Commun.Math.Phys. 213 (2000) 19-37.
- [85] Y.S.Poon and A.Swann Commun.Math.Phys. 241 (2003) 177-189.
- [86] G.Grantcharov, G.Papadopoulos and Y.S.Poon J.Math.Phys. 43 (2002) 3766-3782
- [87] L.Ornea, Y.S.Poon and A.Swann Class.Quant.Grav. 20 (2003) 1845-1856.
- [88] A. Opfermann and G. Papadopoulos math-ph/9807026.
- [89] P.S. Howe, G. Papadopoulos Phys.Lett. B379 (1996) 80.
- [90] A. Opfermann Phys.Lett. B416 (1998) 101-107.
- [91] C. Le Brun J.Diff.Geom 34 (1991) 223.
- [92] N.Hitchin, Stable forms and special metrics, math.DG/0107101.
- [93] B.Acharya and M.O'Loughlin, Phys.Rev.D 55 (1997) 4521.
- [94] D.Page and C.Pope, Class.Quantum.Grav. 3 (1986) 249.
- [95] E.Witten and M.Atiyah Adv. Theor. Math. Phys. 6 (2003) 1.
- [96] A.Brandhuber and P.Berglund, Nucl.Phys. B641 (2002) 351.
- [97] L.Anguelova and C.Lazaroiu, JHEP 0210 (2002) 038; L.Anguelova and C.Lazaroiu JHEP 0209 (2002) 053.
- [98] L.Anguelova and C.Lazaroiu JHEP 0301 (2003) 066.
- [99] K.Behrdnt, G. Dall'Agata, D. Lust and S. Mahapatra, JHEP 0208 (2002) 027; K.Behrdnt, Nucl.Phys. B635 (2002) 158.
- [100] M.Aganagic and C.Vafa, Mirror Symmetry and a G_2 Flop, hep-th/0105225.
- [101] M.Atiyah, J.Maldacena and C.Vafa, J.Math.Phys. 42 (2001) 3209.
- [102] R.Gopakumar and C.Vafa, Adv.Theor.Math.Phys. 5 (1999) 413.
- [103] Shamit Kachru and John McGreevy, JHEP 0106 (2001) 027.

- [104] C.Vafa, J.Math.Phys. 42 (2001) 2798.
- [105] I. Bakas, E. Floratos, A. Kehagias, Phys.Lett. B445 (1998) 69.
- [106] H. Partouche, B. Pioline, JHEP 0103 (2001) 005; P. Kaste, A. Kehagias, H. Partouche, JHEP 0105 (2001) 058.
- [107] R. Blumenhagen and V.Braun JHEP 0112 (2001) 006.
- [108] T. Eguchi, Y. Sugawara, Nucl.Phys. B630 (2002) 132; K. Sugiyama, S.Yamaguchi, Phys.Lett. B538 (2002) 173.
- [109] A. Belhaj, J.Phys. A35 (2002) 8903.
- [110] S. Acharya and S. Gukov Phys.Rept. 392 (2004) 121-189.
- [111] V.Apostolov and S.Salamon Comm. Math. Phys. 246 (2004), no. 1, 43-61.
- [112] P. Kaste, R. Minasian, M. Petrini and A. Tomasiello, JHEP 0209 (2002) 033
- [113] R. Hernandez, K. Sfetsos and D.Zoakos JHEP 0409 (2004) 010.
- [114] H. Nishino and S.Rajpoot Phys.Lett. B569 (2003) 102-112.
- [115] S. Gukov, Shing-Tung Yau, E. Zaslow "Duality and Fibrations on G_2 Manifolds", hep-th/0203217.
- [116] N.Grandi and A. Lugo Phys.Lett. B553 (2003) 293-300.
- [117] J.D. Edelstein, A. Paredes and A.V. Ramallo JHEP 0301 (2003) 011; J. Edelstein and C. Nunez JHEP 0104 (2001) 028.
- [118] M. J. Duff and James T. Liu Nucl.Phys. B674 (2003) 217-230.
- [119] G. Curio JHEP 0303 (2003) 024.
- [120] S.Gubser hep-th/0201114.
- [121] U. Gursoy, C. Nunez and M. Schwelling JHEP 0206 (2002) 015; S. Hartnoll and C. Nunez JHEP 0302 (2003) 049.
- [122] Aalok Misra, JHEP 0210 (2002) 056.
- [123] S.L. Shatashvili and C. Vafa, Superstrings and Manifolds of Exceptional Holonomy, hep-th/940702.
- [124] B.Acharya Nucl.Phys. B524 (1998) 262.
- [125] R.D.Schafer, An Introduction to Nonassociative Algebras, (Academic Press Inc., New York ,1966).
- [126] T.Eguchi and A.Hanson, Ann.Phys. 120 (1979) 82; S.Hawking, Phys.Lett. 60A (1979) 81; V.Belinskii, G.Gibbons, D.Page and C.Pope, Phys.Lett. B76 (1978) 433.

- [127] T.Eguchi, P.Gilkey and A.Hanson, Phys.Rep.66, 5 (1980) 214.
- [128] A.Bilal, J.Derendinger and K.Stefesos, Nucl.Phys. B628 (2002) 112.
- [129] R.S.Ward, Class.Quant.Grav 7 (1990) L95; N.Woodhouse, Class.Quant.Grav 6 (1989) 933.
- [130] D.Calderbank and P.Tod Differ.Geom.Appl. 14 (2001) 199.
- [131] D.Calderbank J. Geom. Phys. 36 (2000) 152.
- [132] D.Calderbank and H.Pedersen, Selfdual Einstein metrics with torus symmetry, math.DG/0105263; D.Calderbank and M.Singer, Einstein metrics and complex singularities, math.DG/0206229.
- [133] D.Calderbank and H.Pedersen, Ann.Inst.Fourier 50 (2000) 921.
- [134] R. Bielawski and A.Dancer, Comm.Anal.Geom.8 (2000) 727.
- [135] H.Pedersen, Math.Ann. 274 (1986) 35.
- [136] P.Casteill, E.Ivanov, G.Valent, Phys.Lett. B508 (2001) 354-364; Nucl.Phys. B627 (2002) 403.
- [137] B.de Wit, M. Rocek and S.Vandoren JHEP 0102 (2001) 039.
- [138] J.P. Gauntlett, G.W. Gibbons, G. Papadopoulos and P.K. Townsend Nucl.Phys. B500 (1997) 133.
- [139] R. Portugues JHEP 0403 (2004) 051
- [140] G.Gibbons and N.Manton Phys.Lett.B 356 (1995) 32.
- [141] G.W. Gibbons and P. Rychenkova Commun.Math.Phys. 186 (1997) 585.
- [142] G. Papadopoulos and J. Gutowski Nucl.Phys. B551 (1999) 650.
- [143] T.Kraan Comm.Math.Phys. 212 (2000) 503.
- [144] J.Gauntlett, D.Tong and P.Townsend Phys.Rev.D 63 (2001) 085001.
- [145] J.Gauntlett, D.Tong and P.Townsend Phys.Rev.D 64 (2001) 025010.
- [146] H.Pedersen and Y.S.Poon Comm.Math.Phys.117 (1988) 569.
- [147] D.Joyce Duke.Math.Journal 77 (1995) 519.
- [148] M.Przanowski J.Math.Phys.32 (1991) 1004.
- [149] K.P.Tod The $SU(\infty)$ Toda equation and special four dimensional metrics in Geometry and Physics Aarhaus J.E.Andersen J.Dupont H.Pedersen and A.Swann eds. 1995; Lecture Notes in Pure and Appl.Maths 184 Marcel Dekker New York 1997.
- [150] N.Hitchin Complex manifolds and Einstein's equations in Twistor geometry and Non-linear Systems H.D.Doebner and T.D.Palev eds. Springer 1982.

- [151] I.M.Singer and J.A.Thorpe The curvature of 4-dimensional Einstein spaces in Global Analysis, Papers in honor of K.Kodaira D.C.Spencer and S.Iyanaga eds.
- [152] M.Atiyah, N.Hitchin and I.Singer Proc.R.Soc.London.A 362 (1978) 425.
- [153] A.Ashtekar T.Jacobson and L.Smolin Comm.Math.Phys.115 (1988) 631.
- [154] G.Gibbons and S.Hawking Phys.Rev.D 15 (1977) 2752; K.P.Tod and R.S.Ward Selfdual metrics with selfdual Killing vectors Proc.R.Soc.London.A 368 (1979) 411.
- [155] R. Bielawski and A.Dancer Comm.Anal.Geom.8 (2000) 727.
- [156] K. Behrndt and S. Mahapatra hep-th/0312063.
- [157] P.E.Jones and K.P.Tod Class.Quant.Grav.2 (1985) 565.
- [158] P.Nurowski Class.Quant.Grav 14 (1997) A261.
- [159] K. Becker, M. Becker and A. Strominger Nucl.Phys.B 456 (1995) 130; A. Strominger Phys.Lett. 421 (1998) 139 ; K. Becker and M. Becker, Nucl.Phys.B 551(1999) 102.
- [160] M. Gutperle and M. Spalinski, JHEP **0006** (2000) 037 ; Nucl. Phys. B 598 (2001) 509.
- [161] U. Theis and S. Vandoren, JHEP **0209** (2002) 059 ; L. Anguelova, M. Rocek and S. Vandoren hep-th/0402132.
- [162] H. Ooguri and C. Vafa, Phys. Rev. Lett. 77 (1996) 3298.
- [163] E. Witten, Nucl. Phys. B 474 (1996) 343.
- [164] S. V. Ketov, *Quantum Non-linear Sigma-Models*, Springer-Verlag, 2000
- [165] G.W. Gibbons, D. Olivier, P. J. Ruback, G. Valent, Nucl. Phys. **B296** (1988) 679.
- [166] S. V. Ketov, *Selfinteracting $N=2$ multiplets in $4d$, and UV-finiteness of $2d$, $N=4$ non-linear sigma-models*, Tomsk preprint 1984; published in the Proceedings of the International Colloquium ‘Group-Theoretical Methods in Physics’, Nauka: Moscow, 1985, Vol. 1, p. 87;
S. V. Ketov, B. B. Lokhvitsky and I. V. Tyutin, Theor. Math. Phys. **71** (1987) 496.
- [167] E. Bergshoeff, C.M. Hull and T. Ortin Nucl.Phys. B451 (1995) 547.
- [168] B. De Wit and H. Nicolai Nucl.Phys.B 231 (1984) 506.
- [169] M.Gunaydin and F.Gursey J.Math.Phys 14 (1973) 1651.
- [170] A. Futaki, Kahler-Einstein metrics and integral invariants, Lec.Notes Math.1314 (Springer Verlag, Berlin, 1980).
- [171] U. Lindstrom and M. Rocek Nucl.Phys.B 222 (1983) 288.
- [172] V. Arnold, A. Gusein-Zade and A. Varchenko, *Singularities of Differentiable Maps*, Birkhauser, 1985.

- [173] S.V.Ketov Nucl.Phys. B649 (2003) 365.
- [174] S.V.Ketov Nucl.Phys. B656 (2003) 63; S.V.Ketov Phys.Lett. B558 (2003) 119.
- [175] I. Antoniadis, R. Minasian, S. Theisen and P. Vanhove Class.Quant.Grav. 20 (2003) 5079.
- [176] V. Apostolov, S. Salamon Kahler reduction of metrics with holonomy G_2 , math.DG/0303197.
- [177] A. Gray Math Z 123 (1971) 290.
- [178] P. Freund and M.Rubin Phys.Lett.B 97 (1980) 233.
- [179] P. West and P.West, Nucl. Phys. B 243 (1984) 112; M. Huq and N. Namazie Class. Quant. Grav 2 (1985) 293; F. Giani and M. Pernici Phys. Rev. D 30 (1984) 325.
- [180] E.Ivanov and S.Krivonos J.Phys A Math. Gen 17 (1984) L671; Theor. Math.Phys 63 (1985) 260.
- [181] E.A. Ivanov, S.O. Krivonos and V.M. Leviant, Nucl. Phys. B 304 (1988) 601; Phys. Lett.B 215 (1989) 689; Phys. Lett.B 221 (1989) 432E.
- [182] O.E. Gorovoy and E.A. Ivanov, Nucl. Phys. B 381 (1992) 394.
- [183] E. Ivanov and A. Sutulin, Nucl. Phys. B 432 (1994) 246.
- [184] E.Ivanov Nucl.Phys.B 576 (2000) 543.
- [185] A. Galperin, E.Ivanov and O.Ogievetsky, Annals Phys. 230 (1994) 201 A. Galperin and O.Ogievetsky Phys.Lett.B 30 (1993) 67.
- [186] G.W. Gibbons, G. Papadopoulos, K.S. Stelle Nucl.Phys. B508 (1997) 623.
- [187] C.Burdik, S.Krivonos and A.Shcherbakov "N=4 supersymmetric Eguchi-Hanson sigma model in d=1", hep-th/0508165.
- [188] S. Bellucci, S. Krivonos, A. Marrani and E. Orazi "Root Action for N=4 Supersymmetric Mechanics Theories", hep-th/0511249.
- [189] O.P.Santillan Nucl.Phys.B 660 (2003) 169.
- [190] O. P. Santillan and A. Z. Zorin Commun.Math.Phys 255 (2005) 33.
- [191] S. Ketov, O. Santillan and A. Zorin Mod.Phys.Lett. A19 (2004) 2645.
- [192] A.Isaev and O.P.Santillan JHEP 0905 (2005) 045.